

# Optimization

## Introduction to optimization problems

- minima, maxima, feasible sets, equivalent reformulations

## Criteria for optimality

- differentiable case; sufficient & necessary conditions

## Duality

- Lagrange function, dual problem

## Convex analysis and optimization

- convex sets, convex functions, optimality criteria

## Optimization algorithms

- simplex method
- primal-dual methods
- smooth unconstrained optimization
- constrained optimization

# Introduction: Idea and motivation

## Optimization problem

$$\text{minimize } f(x), \quad x \in \mathbb{R}^n \quad \text{subject to } c_E(x) = b_E \text{ and } c_I(x) \leq b_I$$

$x$

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$

$$c_E: \mathbb{R}^n \rightarrow \mathbb{R}^p$$

$$c_I: \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$b_E \in \mathbb{R}^p$$

$$b_I \in \mathbb{R}^m$$

optimisation variable

objective function

equality constraint function

inequality constraint function

$c_I(x) \leq b_I$  is meant component-wise

## Classes of optimization problems

linear program :  $f \in C_E \cup C_I$  linear / affine

nonlinear program :  $f \in C_E, C_I$  nonlinear

convex optimisation problem :  $f \in C_E$  linear,  $C_I$  convex (comp.-wise)

$g: \mathbb{R}^n \rightarrow \mathbb{R}^m$  linear  $\Leftrightarrow g(ax + by) = ag(x) + bg(y) \forall x, y \in \mathbb{R}^n, a, b \in \mathbb{R}$

$g: \mathbb{R}^n \rightarrow \mathbb{R}$  convex  $\Leftrightarrow g(ax + (1-a)y) \leq ag(x) + (1-a)g(y) \forall x, y \in \mathbb{R}^n, a \in [0, 1]$

other types not covered in this lecture (at least initially)

discrete optimisation problem :  $f \in C_E, C_I$  defined on  $\mathbb{Z}^n$

combinatorial optimisation pb. :  $f \in C_E, C_I$  defined on  $\{1, \dots, k\}^n$

optimisation in Banach spaces :  $f \in C_E, C_I$  def. on Banach sp.

optimisation on manifolds :  $f \in C_E, C_I$  def. on Riemann. manif.

## Applications

### portfolio optimisation

- optimally invest money in  $n$  stocks;  $x_i =$  investment in stock  $i$
- constraint on budget, positivity, ...
- objective fcn: total risk, variance of income, ...

### process- or product-optimisation


- $x_i =$  production amount of product  $i$
- constraints on resources, capacity, ...
- objective fcn: profit

### data fitting

- $x =$  parameter in a model
- constraints: a-priori information (positivity etc.)
- objective fcn: data discrepancy

# Introduction: Idea and motivation

## Example 1: Optimization of high pressure gas network

- network 

$q_i = \text{flows}$   
 $p_i = \text{pressures}$   
 $d_i = \text{demands}$
- mass constraint for node 3:  
in general  $Aq - d = 0$   
*sparse matrix*
- pipe constraint for pipe 1:  
in general  $p^T B p + g(q) = 0$   
*quadratic / nonlinear constraint*
- compressor constraint:  
in general  $Cq + z \cdot h(p, q) \geq 0$   
*discrete / nonlinear constraint*
- bounds  $p \leq p_{\max}$   

- objectives: minimise supply / compressor costs, etc.

# Introduction: Idea and motivation

## Example 2: Least squares fitting

$$\min \|Ax - b\|_2^2 = \sum_{i=1}^k (a_i^T x - b_i)^2, \quad A \in \mathbb{R}^{k \times n}, \quad b \in \mathbb{R}^k, \quad k \geq n$$

*rows of A*

seek minimum by setting derivative to zero:

$$0 = A^T(Ax - b)$$

if  $A^T A$  is invertible, then

$$x = (A^T A)^{-1} A^T b$$

pseudoinverse  $A^+ = (A^T A)^{-1} A^T$

classical image processing application: deblurring

$x$  = pixel-wise vector of grey values of true image

$b$  = observed (blurred) grey values

$A$  = blurring matrix

## Difference to Calculus of Variations

- both fields tightly connected
- CoV tries to establish existence of minimisers and their properties, typically via "energy methods"
- Optimisation tries to find minimisers and criteria for optimality; emphasises numerical methods

## Overall plan of lecture

- Introduction to optimisation problems  
(minima, maxima, feasible sets, equivalent reformulations)
- Criteria for optimality  
(differentiable case; sufficient & necessary conditions)
- Duality  
(Lagrange function, dual problem)
- Convex optimisation  
(convex sets, convex functions, optimality criteria)
- Optimisation algorithms
  - simplex method
  - primal-dual methods
  - smooth unconstrained optimisation
  - constrained optimisation

# Introduction: Optimization problems

## Basic notation

$$\min f(x) \quad \text{subject to} \quad c_E(x) = 0, \quad c_I(x) \leq 0 \quad (\text{pt-wise}) \quad (*)$$

(\*) describes the problem to find  $x \in \mathbb{R}^n$

that minimises  $f$  among all  $x$  satisfying the constraints

notation:  $\cdot f \equiv f_0 : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$

$$\cdot c_I \equiv (f_1, \dots, f_m)^T : \mathbb{R}^n \rightarrow (\mathbb{R} \cup \{\infty\})^m$$

$$\cdot c_E \equiv (h_1, \dots, h_p)^T : \mathbb{R}^n \rightarrow (\mathbb{R} \cup \{\infty\})^p$$

Def: domain of  $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$  :  $\text{dom } f = \{x \in \mathbb{R}^n \mid f(x) < \infty\}$

Def: domain of optimisation problem (\*) :  $\mathcal{D} = \bigcap_{i=0}^m \text{dom } f_i \cap \bigcap_{i=1}^p \text{dom } h_i$

# Introduction: Optimization problems

## Basic terminology

Def:  $x \in \mathcal{D}$  is feasible if all constraints  $f_1(x), \dots, f_m(x) \leq 0$  and  $h_1(x) = \dots = h_p(x) = 0$  are fulfilled.

$(*)$  is feasible if there is a feasible  $x \in \mathcal{D}$

Def:  $p^* = \inf \{ f_0(x) \mid x \in \mathcal{D} \text{ is feasible} \}$  is called optimal value

if  $(*)$  is not feasible, we set  $p^* := \infty$

if  $\exists x_k \in \mathcal{D}$  feasible with  $f_0(x_k) \xrightarrow{k \rightarrow \infty} -\infty$ , set  $p^* = -\infty$ ;  
then  $(*)$  is called unbounded from below

Def:  $x^*$  is called (globally) optimal point if  $x^*$  is feasible and  $f(x^*) = p^*$

$X_{\text{opt}} = \{ x \in \mathcal{D} \mid x \text{ is optimal point} \}$

A feasible point  $x$  with  $f_0(x) \leq p^* + \varepsilon$  is called  $\varepsilon$ -suboptimal

A feasible point  $x$  is called locally optimal if  $\exists R > 0$  s.t.

$$f_0(x) = \inf \{ f_0(z) \mid z \in \mathcal{D} \text{ is feasible}, \|z - x\| \leq R \}$$

A locally/globally optimal point  $x$  is called strictly optimal if  $\exists R > 0$  s.t.  $f_0(x) < f_0(y)$  for all feasible  $y \in \mathcal{D}$  with  $y \neq x, \|x - y\| \leq R$

# Introduction: Optimization problems

## Examples

Optimisation problems on  $\mathbb{R}_{++} := \{x \in \mathbb{R} \mid x > 0\}$

1)  $f_0(x) = \frac{1}{x}$        $p^* = 0$ ,  $X_{opt} = \emptyset$

2)  $f_0(x) = -\log x$        $p^* = -\infty$ ,  $X_{opt} = \emptyset$

3)  $f_0(x) = x \log x$        $p^* = -\frac{1}{e}$ ,  $x^* = \frac{1}{e}$ ,  $X_{opt} = \{\frac{1}{e}\}$

Thm ; Let  $(*)$  be a convex optimisation problem,  
i.e.  $f_0, \dots, f_m$  are convex,  $h_1, \dots, h_p$  affine.  
Then  $x^*$  locally optimal  $\Leftrightarrow x^*$  globally optimal

proof : homework

## Equivalent optimization problems

- (\*) is the standard form of an optimisation problem.

It can always be generated, e.g. the constraint  $g_i(x) = \tilde{g}_i(x)$  is equivalent to  $h_i(x) := g_i(x) - \tilde{g}_i(x) = 0$  or  $\tilde{f}_i(x) \geq 0$  is equivalent to  $f_i(x) := -\tilde{f}_i(x) \leq 0$ .

- Two optimisation problems are called equivalent if the solution of one can be calculated from the solution of the other

Example:  $\min f_0(x) = \alpha_0 f_0(x)$

subject to  $\tilde{f}_i(x) = \alpha_i f_i(x) \leq 0, \tilde{h}_j(x) = \beta_j h_j(x) = 0, i=1, \dots, m, j=1, \dots, p$

is for  $\alpha_i > 0, \beta_j \neq 0$  equivalent to

$\min f_0(x)$  subject to  $f_i(x) \leq 0, h_j(x) = 0$

# Introduction: Transformations of optimization problems

## Basic transformations I

$$(P_1) \quad \min f_0(x) \quad \text{s. t.} \quad f_i(x) \leq 0, \quad i=1, \dots, m, \quad h_i(x) = 0, \quad i=1, \dots, p$$

$$(P_2) \quad \min \tilde{f}_0(x) \quad \text{s. t.} \quad \tilde{f}_i(x) \leq 0, \quad i=1, \dots, m, \quad \tilde{h}_i(x) = 0, \quad i=1, \dots, p$$

- Substitution of variables

Let  $\phi: \mathbb{R}^n \rightarrow \mathbb{R}^n$  bijective,

$$\tilde{f}_i = f_i \circ \phi \quad (i=0, \dots, m) \quad \tilde{h}_i = h_i \circ \phi \quad (i=1, \dots, p)$$

If  $x$  solves  $(P_1)$ , then  $\phi^{-1}(x)$  solves  $(P_2)$  and vice versa.

• Transformation of objective and constraint functions

Let  $\psi_0: \mathbb{R} \rightarrow \mathbb{R}$  strictly monotonically increasing

$$\psi_1, \dots, \psi_m: \mathbb{R} \rightarrow \mathbb{R} \quad \text{with} \quad \psi_i(u) \leq 0 \Leftrightarrow u \leq 0$$

$$\phi_1, \dots, \phi_p: \mathbb{R} \rightarrow \mathbb{R} \quad \text{with} \quad \phi_i(u) = 0 \Leftrightarrow u = 0$$

$$\tilde{f}_i = \psi_i \circ f_i, \quad i=0, \dots, m, \quad \tilde{h}_i = \phi_i \circ h_i, \quad i=1, \dots, p$$

$$\text{We have} \quad X_{\text{opt}}^{(P_1)} = X_{\text{opt}}^{(P_2)}$$

## Basic transformations II

$$(P_1) \quad \min f_0(x) \quad \text{s.t.} \quad f_i(x) \leq 0, \quad i=1, \dots, m, \quad h_i(x) = 0, \quad i=1, \dots, p$$

• slack variables  $s_i$ : exploit  $f_i(x) \leq 0 \Leftrightarrow \exists s_i \geq 0 : f_i(x) + s_i = 0$

$$(P_2) \quad \min f_0(x) \quad \text{s.t.} \quad s_i \geq 0, \quad i=1, \dots, m, \quad f_i(x) + s_i = 0, \quad h_j(x) = 0, \quad i=1, \dots, m, \quad j=1, \dots, p$$

$(x, s)$  feasible (optimal) for  $(P_2) \Leftrightarrow x$  feasible (optimal) for  $(P_1)$  &  $s_i = -f_i(x)$

• Elimination of equality constraints

Assume  $C = \{x \in \mathbb{R}^n \mid h_1(x) = \dots = h_p(x) = 0\}$  can be parametrised

by  $\phi : \mathbb{R}^k \rightarrow \mathbb{R}^n$ , i.e.  $x \in C \Leftrightarrow \exists z \in \mathbb{R}^k : x = \phi(z)$

$$\tilde{f}_i = f_i \circ \phi, \quad i=0, \dots, m$$

$$(P_2) \quad \min \tilde{f}_0(x) \quad \text{s.t.} \quad \tilde{f}_1(x), \dots, \tilde{f}_m(x) \leq 0$$

$$X_{\text{opt}}^{(P_1)} = \phi(X_{\text{opt}}^{(P_2)})$$

Example:  $h_1(x) = \dots = h_p(x) = 0 \Leftrightarrow Ax = b$  (linear constraints)

Set  $\phi(z) = Fz + x_0$  for  $F \in \mathbb{R}^{n \times k}$  with  $\text{range } F = \ker A$ ,  $x_0$  feasible

## Basic transformations III

$$(P_1) \min f_0(x) \quad \text{s.t.} \quad f_i(x) \leq 0, \quad i=1, \dots, m, \quad h_i(x) = 0, \quad i=1, \dots, p$$

• implicit constraints

$$\text{Let } F(x) = \begin{cases} f_0(x) & \text{if } f_1(x), \dots, f_m(x) \leq 0, \quad h_1(x) = \dots = h_p(x) = 0 \\ \infty & \text{else} \end{cases}$$

$$(P_2) \min F(x)$$

$$X_{\text{opt}}^{(P_2)} = X_{\text{opt}}^{(P_1)}$$

$$\text{Example: } \min \begin{cases} \|x\|^2 & \text{if } Ax = b \\ \infty & \text{else} \end{cases} \Leftrightarrow \min \|x\|^2 \quad \text{s.t.} \quad Ax = b$$

• optimisation over one variable

$$\text{Let } x = (x_1, x_2) \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2}, \quad \tilde{F}(x_1) = \inf \left\{ f_0(x_1, x_2) \mid f_1(x_1, x_2), \dots, f_m(x_1, x_2) \leq 0, h_1(x_1, x_2) = \dots, h_p(x_1, x_2) = 0 \right\}$$

$$(P_2) \min \tilde{F}(x_1)$$

$$x_1 \in X_{\text{opt}}^{(P_2)} \quad \Leftrightarrow \quad \exists x_2 \in \mathbb{R}^{n_2} : (x_1, x_2) \in X_{\text{opt}}^{(P_1)}$$

useful if minimisation over  $x_2$  can be done explicitly

## Basic transformations IV

$$(P_1) \min f_0(x) \text{ s.t. } f_i(x) \leq 0, i=1, \dots, m, \quad h_i(x)=0, i=1, \dots, p$$

• epigraph formulation

$$(P_2) \min_{t, x} t \text{ s.t. } f_0(x) - t \leq 0, f_1(x), \dots, f_m(x) \leq 0, \quad h_1(x), \dots, h_p(x) = 0$$

$$(x, t) \in X_{\text{opt}}^{(P_2)} \Leftrightarrow x \in X_{\text{opt}}^{(P_1)} \text{ and } f_0(x) = t$$

• generalised epigraph formulation

$$\text{Let } f_0(x) = \sum_{j=1}^M g_j(x)$$

$$(P_2) \min_{t_1, \dots, t_M, x} \sum_{j=1}^M t_j \text{ s.t. } g_j(x) - t_j \leq 0, j=1, \dots, M + \text{other constraints}$$

$$(x, t_1, \dots, t_M) \in X_{\text{opt}}^{(P_2)} \Leftrightarrow x \in X_{\text{opt}}^{(P_1)} \text{ and } g_j(x) = t_j, j=1, \dots, M$$

# Optimality criteria: Unconstrained optimization

## Motivation and notation

### Optimality Conditions

- indicate when a point is not optimal (necessary conditions)
- guarantee that a candidate solution is indeed (locally) optimal (sufficient condition)
- guide the design of algorithms

### Assumption and notation

- in the following, we assume that  $f, \dots, f_{n_1}, \dots, f_{n_p}$  are continuously differentiable up to the order needed in each result (e.g., for first order optimality conditions up to first order)
- $Df = \left( \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right)$ ,  $\nabla f = Df^T$
- $D^2f = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \dots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \dots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$

Optimality criteria: Unconstrained optimization *case  $m = p = 0$*

## 1st order necessary conditions

Thm: If  $x^*$  is a local optimum of  $(X)$ , then  $Df_0(x^*) = 0$ .

proof: By contradiction: suppose  $Df_0(x^*) \neq 0$ .

By Taylor's thm, for  $\alpha > 0$ ,

$$\begin{aligned} f_0(x^* - \alpha \nabla f(x^*)) &= f_0(x^*) + Df_0(x^*)(-\alpha \nabla f_0(x^*)) + \underbrace{o(|\alpha \nabla f_0(x^*)|)}_{\text{"little o-notation"}} \\ &= f_0(x^*) - \underbrace{\alpha |\nabla f_0(x^*)|^2}_A + \underbrace{o(\alpha |\nabla f_0(x^*)|)}_B \end{aligned}$$

For all  $\alpha > 0$  small enough,  $B < A$  so that

$$f_0(x^* - \alpha \nabla f(x^*)) < f_0(x^*)$$

$\Rightarrow x^*$  cannot be local optimum.  $\square$

"little-o notation": We write  $o(g(s))$  if we want to refer to a function  $f(s)$  with  $\lim_{s \rightarrow 0} \frac{f(s)}{g(s)} = 0$ .

# Optimality criteria: Unconstrained optimization

## 2nd order necessary conditions

Thm: If  $x^*$  is a local optimum of  $(X)$ , then  $Df_0(x^*) = 0$  and  $D^2f_0(x^*)$  is positive semi-definite, i.e.

$$s^T D^2f_0(x^*) s \geq 0 \quad \forall s \in \mathbb{R}^n.$$

proof: By contradiction: suppose  $\exists s \in \mathbb{R}^n : s^T D^2f_0(x^*) s < 0$ .  
We know  $Df_0(x^*) = 0$  already.

By Taylor's theorem, for  $\alpha > 0$ ,

$$f_0(x^* + \alpha s) = f_0(x^*) + \underbrace{\frac{1}{2} (\alpha s)^T D^2f_0(x^*) (\alpha s)}_{A < 0} + \underbrace{o(|\alpha s|^2)}_B$$

For all  $\alpha > 0$  small enough,  $B < |A|$  so that

$$f_0(x^* + \alpha s) < f_0(x^*)$$

$\Rightarrow x^*$  cannot be local optimum. □

# Optimality criteria: Unconstrained optimization

## 2nd order sufficient conditions

Thm: If  $Df_0(x^*) = 0$  and  $D^2f_0(x^*)$  is positive definite, i.e.

$$s^T D^2f_0(x^*) s > 0 \quad \forall s \in \mathbb{R}^n,$$

then  $x^*$  is a strictly locally optimal point.

proof: By continuity,  $D^2f_0(x)$  is pos. def. for all  $x$  in an open ball  $B_\delta(x^*)$  around  $x^*$ .

Let  $x \in B_\delta(x^*)$ ,  $x \neq x^*$ . By Taylor's thm.,  $\exists z$  between  $x$  and  $x^*$  s.t.

$$f_0(x) = f_0(x^*) + \underbrace{Df_0(x^*)}_{0} (x-x^*) + \underbrace{\frac{1}{2}(x-x^*)^T D^2f_0(z)}_{>0} (x-x^*)$$

$$> f_0(x^*)$$

□

# Optimality criteria: Unconstrained optimization

## Example in $\mathbb{R}^2$

Find local minima of  $f(x_1, x_2) = x_1^3 + 3x_1^2x_2 + 3x_1x_2^2 + x_2^3 - 2x_1^2 - 8x_1x_2 - 2x_2^2$

1<sup>st</sup> order necessary cond.:

$$0 = \nabla f(x_1, x_2) = \begin{pmatrix} 3x_1^2 + 6x_1x_2 + 3x_2^2 - 4x_1 - 8x_2 \\ 3x_1^2 + 6x_1x_2 + 3x_2^2 - 4x_2 - 8x_1 \end{pmatrix}$$

$$\Rightarrow x = (0, 0) \quad \text{or} \quad x = (1, 1)$$

2<sup>nd</sup> order necessary cond.:

$$D^2f(x_1, x_2) = \begin{pmatrix} 6x_1 + 6x_2 - 4 & 6x_1 + 6x_2 - 8 \\ 6x_1 + 6x_2 - 8 & 6x_1 + 6x_2 - 4 \end{pmatrix}$$

Local maximum or saddle?

$$\begin{array}{l} \swarrow \\ D^2f(0, 0) = \begin{pmatrix} -4 & -8 \\ -8 & -4 \end{pmatrix} \end{array} \quad \begin{array}{l} \text{not pos. semi-def.} \\ \text{(determinant } < 0) \end{array}$$

2<sup>nd</sup> order sufficient cond.:

$$D^2f(1, 1) = \begin{pmatrix} 8 & 4 \\ 4 & 8 \end{pmatrix} \quad \begin{array}{l} \text{pos def.} \\ \text{(all minors pos.)} \end{array}$$




$\Rightarrow (1, 1)$  is only locally optimal point

Optimality criteria: Equality constraints

## Tangent plane & regular points

A set of equality constraints  $0 = c_E(x) = \begin{pmatrix} h_1(x) \\ \vdots \\ h_p(x) \end{pmatrix}$  defines a hypersurface  $S$  in  $\mathbb{R}^n$ .

In general:  $c_E$  smooth  $\Rightarrow S$  smooth  $\leftarrow$  what can happen at non-regular points?



Def: A curve on  $S$  is a continuous function  $x: [a, b] \rightarrow S$ .

•  $x$  "passes through  $x^* \in S$ " if  $\exists t^* \in [a, b]: x(t^*) = x^*$ .

• The tangent plane  $T_{x^*} S$  to  $S$  at  $x^*$  is the vector space

$$T_{x^*} S = \{ \dot{x}(t^*) \mid x \text{ is differentiable curve on } S \text{ with } x(t^*) = x^* \}$$

Def:  $x^* \in \mathbb{R}^n$  with  $c_E(x^*) = 0$  is called regular if

$\nabla h_1(x^*), \dots, \nabla h_p(x^*)$  are linearly independent.

Optimality criteria: Equality constraints

## Tangent plane at regular points

Thm: At a regular point  $x^*$  of the surface  $S = \{x \in \mathbb{R}^n \mid c_E(x) = 0\}$ ,  
 $T_{x^*} S = \{y \mid Dc_E(x^*)y = 0\}$ .

proof: " $\subset$ ": Let  $y \in T_{x^*} S$ , i.e.  $\exists x: [a, b] \rightarrow S, t \in [a, b], x(t) = x^*, \dot{x}(t) = y$ .

$$c_E(x(t)) = 0 \Rightarrow 0 = [c_E(x(t))] \Big|_{t=t^*}' = Dc_E(x(t^*)) \dot{x}(t^*)$$

" $\supset$ ": Let  $Dc_E(x^*)y = 0$ .

idea: find a curve  $t \mapsto x^* + ty + Dc_E(x^*)^T u(t)$  on  $S$

Consider  $F: \mathbb{R}^{p+1} \rightarrow \mathbb{R}^p, F(t, u) = c_E(x^* + ty + Dc_E(x^*)^T u)$

$F(0, 0) = 0, D_u F(0, 0) = Dc_E(x^*) Dc_E(x^*)^T$  is invertible

implicit function  $\Rightarrow \exists$  continuous  $u: [-a, a] \rightarrow \mathbb{R}^p$  s.t.  $F(t, u(t)) = 0$

$\Rightarrow x(t) = x^* + ty + Dc_E(x^*)^T u(t)$  is curve on  $S$

$\Rightarrow 0 = \frac{d}{dt} c_E(x(t)) \Big|_{t=0} = Dc_E(x^*)y + Dc_E(x^*) Dc_E(x^*)^T \dot{u}(0)$

$\Rightarrow \dot{u}(0) = 0 \Rightarrow \dot{x}(0) = y + Dc_E(x^*)^T \dot{u}(0) = y$

□

Optimality criteria: Equality constraints

Case  $m=0$

## 1st order necessary conditions I

Lemma: If  $x^*$  is a local optimum of  $(\star)$  and regular w.r.t.  $c_E$ ,  
then  $Df_0(x^*)y = 0$   
for all  $y \in \mathbb{R}^n$  with  $Dc_E(x^*)y = 0$ .

proof: Let  $Dc_E(x^*)y = 0 \xrightarrow{\text{previous thm}} y \in T_{x^*}S$  for  $S = \{x \in \mathbb{R}^n \mid c_E(x) = 0\}$   
 $\Rightarrow \exists x: [-a, a] \rightarrow S$  differentiable,  $\dot{x}(0) = y$ ,  $x(0) = x^*$ .

•  $x^*$  is local optimum of  $(\star)$

$\Rightarrow t=0$  is local minimizer of  $t \mapsto f_0(x(t))$

$\Rightarrow 0 = [f_0(x(t))]'|_{t=0} = Df_0(x^*)y$  □

interpretation:  $\nabla f_0(x^*)$  is orthogonal to tangent plane

# Optimality criteria: Equality constraints

## 1st order necessary conditions II

Thm: If  $x^*$  is a local optimum of  $(X)$  and regular w.r.t.  $C_E$ , then  
 $\exists$  Lagrange multiplier  $\lambda \in \mathbb{R}^p$  s.t.  $Df_0(x^*) + \lambda^T DC_E(x^*) = 0$ .

proof: Abbreviate  $A = DC_E(x^*)$  and  $g = \nabla f_0(x^*)$

Let the columns of  $N$  be a basis of  $\ker A = (\text{range } A)^\perp$

$\Rightarrow g = -A^T \lambda + Nz$  for some  $\lambda \in \mathbb{R}^p, z \in \mathbb{R}^{n-p}$

The previous Lemma implies  $N^T g = 0$ , i.e.

$$0 = -N^T A^T \lambda + N^T N z = N^T N z.$$

However,  $N$  has full rank so that  $z = 0$ .

$$\Rightarrow g = -A^T \lambda$$

□

interpretation:  $\nabla f_0(x^*)$  is lin. comb. of  $\nabla h_1(x^*), \dots, \nabla h_p(x^*)$

Optimality criteria: Equality constraints

## 2nd order necessary conditions

Thm: If  $x^*$  is a local optimum of  $(*)$  and regular w.r.t.  $c_E$ , then  
 $\exists \lambda \in \mathbb{R}^p$  with  $Df_0(x^*) + \lambda^T Dc_E(x^*) = 0$  and

$$D^2 f_0(x^*) + \sum_{k=1}^p \lambda_k D^2 h_k(x^*)$$

is positive semidefinite on  $T_{x^*} S$  for  $S = \{x \in \mathbb{R}^n \mid c_E(x) = 0\}$

proof: Let  $x(t)$  be a curve on  $S$  with  $x(0) = x^*$ .

By 2<sup>nd</sup> order condition for unconstrained problems,

$$0 \leq \frac{d^2}{dt^2} f_0(x(t)) \Big|_{t=0} = \dot{x}(0)^T D^2 f_0(x(0)) \dot{x}(0) + Df_0(x(0)) \ddot{x}(0).$$

Also,  $0 = \lambda^T c_E(x(t))$  implies

$$0 = \frac{d^2}{dt^2} \lambda^T c_E(x(t)) = \sum_{k=1}^p \lambda_k \dot{x}(0)^T D^2 h_k(x(0)) \dot{x}(0) + \lambda^T Dc_E(x(0)) \ddot{x}(0)$$

Adding both eqs. and using the 1<sup>st</sup> order condition,

$$0 \leq \dot{x}(0)^T \left[ D^2 f_0(x^*) + \sum_{k=1}^p \lambda_k D^2 h_k(x^*) \right] \dot{x}(0),$$

where  $\dot{x}(0) \in T_{x^*} S$  was arbitrary.  $\square$

Optimality criteria: Equality constraints

## 2nd order sufficient conditions

Thm: If there are  $x^* \in \mathbb{R}^n$ ,  $\lambda \in \mathbb{R}^p$  with

$$c_E(x^*) = 0, \quad Df_0(x^*) + \lambda^T Dc_E(x^*) = 0,$$

$$D^2 f_0(x^*) + \sum_{i=1}^p \lambda_i D^2 h_i(x^*) \text{ pos. def. on } T_{x^*} S,$$

then  $x^*$  is a strict local optimum of  $(\mathcal{P})$ .

proof: Assume the opposite, i.e.  $\exists$  sequence  $y_k = x^* + \delta_k s_k$ ,  $\delta_k > 0$ ,  $\delta_k \rightarrow 0$ ,  $|s_k| = 1$ ,  
with  $c_E(y_k) = 0$  and  $f_0(y_k) \leq f_0(x^*)$ .

$s_k$  bounded  $\Rightarrow$  subsequence converges. wlog,  $s_k \rightarrow s^*$ .

$0 = c_E(y_k) - c_E(x^*)$ ; divide this by  $\delta_k$  and let  $k \rightarrow \infty \Rightarrow Dc_E(x^*)s^* = 0$ .

Taylor's thm:  $0 \geq f_0(y_k) - f_0(x^*) = \delta_k Df_0(x^*)s_k + \frac{\delta_k^2}{2} s_k^T D^2 f_0(\eta_0) s_k \quad (E_0)$

$$0 = h_i(y_k) - h_i(x^*) = \delta_k Dh_i(x^*)s_k + \frac{\delta_k^2}{2} s_k^T D^2 h_i(\eta_i) s_k \quad (E_i)$$


for some  $\eta_i$  between  $x^*$  and  $y_k$ .

$$(E_0) + \sum_{i=1}^p \lambda_i (E_i) \Rightarrow 0 \geq \frac{\delta_k^2}{2} s_k^T \left[ D^2 f_0(\eta_0) + \sum_{i=1}^p \lambda_i D^2 h_i(\eta_i) \right] s_k \quad \forall k \quad \square$$

# Optimality criteria: Equality constraints

## Example

Solve  $\min_{x \in \mathbb{R}^3} |x - (0, 2, 0)|^2$  s. t.  $h_1(x) = |x|^2 - 1 = 0$ ,  $h_2(x) = x_1 = 0$



1<sup>st</sup> order necessary condition:  $0 = \nabla f(x) + \lambda_1 \nabla h_1(x) + \lambda_2 \nabla h_2(x) = 2(x - \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}) + 2\lambda_1 x + \lambda_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

$$0 = h_1(x)$$

$$0 = h_2(x)$$

$$\Rightarrow x_1 = 0, \quad x_3 = 0, \quad x_2 = \pm 1, \quad \lambda_2 = 0, \quad \lambda_1 = \begin{cases} 1 \\ -3 \end{cases}$$

2<sup>nd</sup> order necessary condition:  $\mathcal{D}^2 f \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda_1 \mathcal{D}^2 h_1 \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda_2 \mathcal{D}^2 h_2 \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = 2\mathbf{I} + 2\lambda_1 \mathbf{I} \ll 0$

$\Rightarrow$  no local minimum

2<sup>nd</sup> order sufficient condition:  $\mathcal{D}^2 f \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda_1 \mathcal{D}^2 h_1 \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda_2 \mathcal{D}^2 h_2 \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = 2\mathbf{I} + 2\lambda_1 \mathbf{I} \gg 0$

$\Rightarrow$  local minimum

## Active set &amp; regular points

Def: • Let  $x^*$  be feasible, i.e.  $h_1(x^*) = \dots = h_p(x^*) = 0, f_1(x^*), \dots, f_m(x^*) \leq 0$ .

Let  $J$  be the set of indices  $j$  with  $f_j(x^*) = 0$ .

The set of constraints  $f_j(x^*) \leq 0, j \in J$ , is called active set.

$f_j(x^*) \leq 0, j \notin J$ , is called inactive set.

Sometimes we count the equality constraints to the active set.

- $x^*$  is called regular point if  $\nabla h_1(x^*), \dots, \nabla h_p(x^*), \nabla f_j(x^*), j \in J$ , are linearly independent.

In a neighbourhood of  $x^*$ , the inactive set may be completely ignored!

Optimality criteria: Inequality constraints

## Karush-Kuhn-Tucker (KKT) conditions

Thm: If  $x^*$  is a locally optimal point of  $(X)$  and regular, then there are

$\lambda \in \mathbb{R}^p$ ,  $\mu \in \mathbb{R}^m$  with

$$c_E(x^*) = 0, \quad c_I(x^*) \leq 0, \quad \mu \geq 0, \quad \mu^T c_I(x^*) = 0 \text{ (comp.-wise)}$$

$$Df_0(x^*) + \lambda^T Dc_E(x^*) + \mu^T Dc_I(x^*) = 0$$

*"complementary slackness"*  
i.e. at least one of  $\mu_i, (c_I(x^*))_i$  is zero!

proof: • Complementary slackness

$\Leftrightarrow \mu_i \neq 0$  only if  $i \in J$  for the active set  $J$

•  $x^*$  is locally optimal

$\Rightarrow$  it is also locally optimal for the constraints  $c_E(x) = 0, f_i(x) = 0, i \in J$

$\Rightarrow 0 = Df_0(x^*) + \lambda^T Dc_E(x^*) + \mu^T Dc_I(x^*)$  for some  $\lambda \in \mathbb{R}^p, \mu \in \mathbb{R}^m$  with  $\mu_i = 0$  if  $i \notin J$

• assume  $\mu_k < 0$ ; let  $S = \{x \in \mathbb{R}^n \mid c_E(x) = 0, f_i(x) = 0 \forall i \in J \setminus \{k\}\}$ ;

by regularity,  $\exists Y \in T_{x^*} S : Df_k(x^*) y < 0$ ; let  $x$  be curve on  $S$  with  $x(0) = x^*, \dot{x}(0) = y$

$\Rightarrow x(t)$  feasible for  $t \geq 0$  small and  $\frac{d}{dt} f_0(x(t))|_{t=0} = Df_0(x^*) y = -\mu_k Df_k(x^*) y < 0 \quad \square$

# Optimality criteria: Inequality constraints

## 2nd order necessary conditions

Thm: If  $x^*$  is a locally optimal point of  $(*)$  and regular, then  $\exists \lambda \in \mathbb{R}^p, \mu \in \mathbb{R}^m$  such that in addition to the KKT conditions

$$D^2 f_0(x^*) + \sum_{k=1}^p \lambda_k D^2 h_k(x^*) + \sum_{i=1}^m \mu_i D^2 f_i(x^*)$$

is pos. semi-definite on the tangent space to the active constraints.

proof: •  $x^*$  is locally optimal

$\Rightarrow$  it is also locally optimal for the constraints  $c_E(x)=0, f_i(x)=0, i \in J$

$\Rightarrow$  use 2<sup>nd</sup> order necessary condition for equality constraints  $\square$

# Optimality criteria: Inequality constraints

## 2nd order sufficient conditions

Thm:  $x^*$  is a strictly locally optimal point of  $(*)$  if  $\exists \lambda \in \mathbb{R}^p, \mu \in \mathbb{R}^m$  s.t.

$$c_E(x^*) = 0, \quad c_I(x^*) \leq 0, \quad \mu \geq 0, \quad \mu c_I(x^*) = 0 \text{ (comp.-wise)}$$

$$Df_0(x^*) + \lambda^T Dc_E(x^*) + \mu^T Dc_I(x^*) = 0$$

$$\text{and} \quad D^2f_0(x^*) + \sum_{k=1}^p \lambda_k D^2h_k(x^*) + \sum_{i=1}^m \mu_i D^2f_i(x^*)$$

is positive definite on  $\{y \mid Dc_E(x^*)y = 0, Df_i(x^*)y = 0 \text{ for all } i \text{ with } \mu_i > 0\}$ .

proof: Assume  $x^*$  is not a strict optimum, but the conditions hold;

let  $y_k = x^* + \delta_k s_k$  with  $\delta_k > 0, |s_k| = 1, y_k \xrightarrow{k \rightarrow \infty} x^*$  such that  $f_0(y_k) \leq f_0(x^*)$

• wlog,  $\delta_k \rightarrow 0, s_k \rightarrow s^*, 0 \geq Df_0(x^*)s^*, 0 = Dc_E(x^*)s^*$  ← as in case with equality constraints

• for each (active) constraint  $f_j$  with  $\mu_j > 0, f_j(y_k) - f_j(x^*) \leq 0 \Rightarrow Df_j(x^*)s^* \leq 0$

→ if  $Df_j(x^*)s^* < 0$  for a  $j$ , then  $0 \geq Df_0(x^*)s^* = -\lambda^T Dc_E(x^*)s^* - \mu^T Dc_I(x^*)s^* > 0$   $\perp$

→ if  $Df_j(x^*)s^* = 0$  for all active constraints, use proof for equality constraints

□

# Optimality criteria: Inequality constraints

## Example

$$\text{Solve } \min_{x \in \mathbb{R}^3} |x - (0, 2, 0)|^2 \quad \text{s. t. } f_1(x) = |x|^2 - 1 \leq 0, \quad f_2(x) = x_1 - \frac{1}{2} \leq 0$$

1<sup>st</sup> order necessary condition:

$$0 = \nabla f(x) + \mu_1 \nabla f_1(x) + \mu_2 \nabla f_2(x) = 2\left(x - \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}\right) + 2\mu_1 x + \mu_2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$
$$0 \geq f_1(x), \quad 0 \leq \mu_1, \quad \mu_1 f_1(x) = 0$$
$$0 \geq f_2(x), \quad 0 \leq \mu_2, \quad \mu_2 f_2(x) = 0$$
$$\Rightarrow x_1 = 0, \quad x_3 = 0, \quad x_2 = 1, \quad \mu_2 = 0, \quad \mu_1 = 1$$

2<sup>nd</sup> order sufficient condition:

$$D^2 f \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \mu_1 D^2 f_1 \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \mu_2 D^2 f_2 \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} = 2I + 2\mu_1 I \gg 0$$
$$\Rightarrow \text{local minimum}$$

# Lagrange duality: dual function

## Lagrange dual

Recall  $(*)$   $\min_{x \in \mathbb{R}^n} f_0(x)$  s.t.  $c_I(x) \equiv \begin{pmatrix} f_1(x) \\ \vdots \\ f_m(x) \end{pmatrix} \leq 0$ ,  $c_E(x) \equiv \begin{pmatrix} h_1(x) \\ \vdots \\ h_p(x) \end{pmatrix} = 0$   
with domain  $\mathcal{D} = \bigcap_{i=1}^m \text{dom} f_i \cap \bigcap_{j=1}^p \text{dom} h_j$

Def: The Lagrange function to  $(*)$  is  $L: \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ ,  $\text{dom} L = \mathcal{D} \times \mathbb{R}^m \times \mathbb{R}^p$ ,

$$L(x, \mu, \lambda) = f_0(x) + \mu^T c_I(x) + \lambda^T c_E(x).$$

- $\mu, \lambda$  are called dual variables or Lagrange multipliers of  $(*)$
- $x$  is called primal variable
- $g: \mathbb{R}^m \times \mathbb{R}^p \rightarrow \mathbb{R}$ ,  $g(\mu, \lambda) = \inf_{x \in \mathcal{D}} L(x, \mu, \lambda)$ , is called the Lagrange-dual or dual function

Thm: The dual function  $g$  is concave (i.e.  $-g$  is convex).

proof:  $-g$  is pointwise supremum of affine functions  $\implies$  convex  $\square$   
see next chapter!

# Lagrange duality: dual function

## Lower bound on optimal value

Thm: Let  $p^*$  be the optimal value of  $(X)$ . For every  $\mu \geq 0, \lambda$  we have  $g(\mu, \lambda) \leq p^*$ .


proof: Let  $x$  be feasible, i.e.  $c_I(x) \leq 0, c_E(x) = 0$ , and let  $\mu \geq 0$ .

$$\Rightarrow g(\mu, \lambda) = \inf_{x \in S} L(x, \mu, \lambda) \leq L(x, \mu, \lambda) = f_0(x) + \underbrace{\mu^T c_I(x) + \lambda^T c_E(x)}_{\leq 0} \leq f_0(x).$$


Since  $x$  was arbitrary,  $g(\mu, \lambda) \leq \inf_{x \text{ feasible}} f_0(x) \leq 0$  □

Interpretation of dual function as "approximation":

$$\text{Set } \chi_-(u) = \begin{cases} 0, & u \leq 0 \\ \infty, & u > 0 \end{cases}$$



$$\chi_0(u) = \begin{cases} 0, & u = 0 \\ \infty, & u \neq 0 \end{cases}$$



$$\text{We have } (X) \Leftrightarrow \min_{x \in \mathbb{R}^n} \tilde{L}(x) = \min_{x \in \mathbb{R}^n} f_0(x) + \sum_{i=1}^m \chi_-(f_i(x)) + \sum_{j=1}^p \chi_0(h_j(x))$$

Replace  $\chi_-(u_i) \rightsquigarrow \mu_i u_i$  with  $\mu_i \geq 0$ ,  $\chi_0(u_i) \rightsquigarrow \lambda_i u_i$

(replace hard by soft constraints)

$$\Rightarrow p^* = \inf_{x \in \mathbb{R}^n} \tilde{L}(x) \geq \inf_{x \in \mathbb{R}^n} L(x, \mu, \lambda) = g(\mu, \lambda).$$

# Lagrange duality: dual function

## Example: least squares solution & LP

Ex:  $\min_x \|x\|^2 \quad \text{s.t.} \quad Ax = b$

Lagrange fcn:  $L(x, \lambda) = x^T x + \lambda^T (Ax - b)$  (convex, quadratic in  $x$ )

dual fcn:  $g(\lambda) = \inf_{x \in \mathcal{D}} L(x, \lambda) = -\frac{1}{4} \lambda^T A^T A \lambda - \lambda^T b$  (concave, quadratic in  $\lambda$ )

$0 = D_x L = 2x^T + \lambda^T A \Rightarrow x = -\frac{1}{2} A^T \lambda$

if  $AA^T$  invertible, property of dual fcn implies  $g(\lambda) = -\frac{1}{4} \lambda^T A A^T \lambda - \lambda^T b \leq \inf_{x \in \mathcal{D}} \{x^T x \mid Ax = b\} \forall \lambda$

$\Rightarrow b^T (A A^T)^{-1} b = g(-2(A A^T)^{-1} b) = \max_{\lambda} g(\lambda) \stackrel{\text{unconstrained!}}{\leq} \inf_{x \in \mathcal{D}} \{x^T x \mid Ax = b\}$

Ex:  $\min_x c^T x \quad \text{s.t.} \quad Ax = b, -x \leq 0 \quad (**)$

$L(x, \mu, \lambda) = c^T x - \mu^T x + \lambda^T (Ax - b) = -\lambda^T b + (c + A^T \lambda - \mu)^T x$

$g(\mu, \lambda) = \inf_{x \in \mathcal{D}} L(x, \mu, \lambda) = \begin{cases} -\infty & \text{if } \mu \neq c + A^T \lambda \\ -\lambda^T b & \text{else} \end{cases}$

$\Rightarrow (-\lambda^T b)$  for any  $\lambda$  with  $\mu := c + A^T \lambda \geq 0$  is lower bound for (\*\*).

# Lagrange duality: dual function

## Dual problem

Recall: Every  $(\mu, \lambda)$  with  $\mu \geq 0$  satisfies  $g(\mu, \lambda) \leq p^*$ .

Def: The dual problem or Lagrange-dual problem is

$$\max_{\mu \in \mathbb{R}^m, \lambda \in \mathbb{R}^p, \mu \geq 0} g(\mu, \lambda) \quad (\text{DP})$$

best lower bound that  $g$  gives on  $p^*$

- The original problem  $(*)$  is called primal problem.
- $(\mu, \lambda)$  is called dual-feasible if  $\mu \geq 0$  (pointwise) &  $g(\mu, \lambda) > -\infty$ .
- $(\mu^*, \lambda^*)$  is called dual-optimal if it solves (DP).

Thm: (DP) is a convex optimisation problem. (no matter how  $(*)$  looks)

proof: (DP)  $\Leftrightarrow \min_{(\tilde{\mu}, \lambda) \in \mathbb{R}^m \times \mathbb{R}^p} -g(-\tilde{\mu}, \lambda) \quad \text{s.t.} \quad \tilde{\mu} \leq 0$

and  $-g$  is convex.

□

Lagrange duality: dual function

Example: dual problem to LP

The dual problem to a linear program is again an LP:

$$\min_x c^T x \quad \text{s.t.} \quad Ax - b = 0, \quad Bx - v \leq 0$$

$$g(\mu, \lambda) = \inf_x c^T x + \mu^T (Bx - v) + \lambda^T (Ax - b) = \begin{cases} -\mu^T v - \lambda^T b & \text{if } A^T \lambda + B^T \mu + c = 0 \\ -\infty & \text{else} \end{cases}$$

$$\text{dual problem: } \max_{\mu \geq 0, \lambda} g(\mu, \lambda) = \max_{\substack{\mu \geq 0 \\ A^T \lambda + B^T \mu + c = 0}} -\mu^T v - \lambda^T b$$

$$\Leftrightarrow \min_{(\tilde{\mu}, \lambda)} -v^T \tilde{\mu} + b^T \lambda \quad \text{s.t.} \quad \tilde{\mu} \leq 0, \quad A^T \lambda - B^T \tilde{\mu} + c = 0$$

# Lagrange duality: strong duality

## Strong versus weak duality

Let  $p^*$  be the optimum value of the primal pb.  $(*)$ ,  $d^*$  of the dual pb. (DP).

We know  $d^* \leq p^*$ , in particular,

- if  $(*)$  is unbounded from below ( $p^* = -\infty$ ), then (DP) is infeasible ( $d^* = -\infty$ )
- if (DP) is unbounded from above ( $d^* = \infty$ ), then  $(*)$  is infeasible ( $p^* = \infty$ )

Def:

- The property  $d^* \leq p^*$  is called weak duality.
- $p^* - d^* \geq 0$  is called duality gap.
- If  $p^* = d^*$  we say that strong duality holds.

Typically, one does not have strong duality, however, for many convex optimisation problems (and some non-convex ones) one does!

We will later consider criteria that imply strong duality.

In case of strong duality,  $(*)$  and (DP) are equivalent!

# Lagrange duality: strong duality

## Examples

Ex:  $\min_{x \in \mathbb{R}} -x^2 \quad \text{s.t.} \quad x-1 \leq 0, -x-1 \leq 0$

(f0 concave, unbounded below)

$$L(x, \mu_1, \mu_2) = -x^2 + \mu_1(x-1) + \mu_2(-x-1)$$

$$g(\mu_1, \mu_2) = \inf_x L(x, \mu_1, \mu_2) = -\infty$$

$$\Rightarrow d^* = -\infty < -1 = p^*$$

(no strong duality)

Ex:  $\min_{x, y \in \mathbb{R}} e^{-x} \quad \text{s.t.} \quad \frac{x^2}{y} \leq 0, 5-y \leq 0$  (convex problem, degenerate constraints)

$$L(x, y, \mu_1, \mu_2) = e^{-x} + \mu_1 \frac{x^2}{y} + \mu_2(5-y)$$

$$g(\mu_1, \mu_2) = \inf_{x, y} L(x, y, \mu_1, \mu_2) = \begin{cases} 0 & \text{if } \mu_1 = \mu_2 = 0 \\ -\infty & \text{else} \end{cases}$$

$$\Rightarrow d^* = 0 < 1 = p^*$$

(no strong duality)

Ex:  $\min_{x, y \in \mathbb{R}} e^{-x} \quad \text{s.t.} \quad x^2 \leq 0, 5-y \leq 0$

(convex problem)

$$L(x, y, \mu_1, \mu_2) = e^{-x} + \mu_1 x^2 + \mu_2(5-y)$$

$$g(\mu_1, \mu_2) = \begin{cases} -\infty & \text{if } \mu_2 \neq 0 \text{ or } \mu_1 < 0 \\ e^{-x} + e^{-2x}/4\mu_1 & \text{else, where } e^{-x} = 2\mu_1 x \end{cases}$$

$$\Rightarrow d^* = \sup_{\mu_1 \geq 0, \mu_2 \geq 0} g(\mu_1, \mu_2) = 1 = p^*$$


let  $\mu_2 = 0, \mu_1 \rightarrow \infty$

(strong duality)

# Lagrange duality: strong duality


## Geometric intuition and nonconvex problems

Ex:  $\min_{x \in \mathbb{R}} f_0(x) \text{ s.t. } f_1(x) \leq 0$




primal fcn:  $w(u) = \inf \{ f_0(x) \mid f_1(x) = u \}$

dual fcn:  $g(\mu) = \inf_x f_0(x) + \mu f_1(x) = \inf_{(t,u) \in G} t + \mu u$  with  $G = \{ (f_0(x), f_1(x)) \mid x \in \mathbb{R} \}$



if  $g(\mu) = t^* + \mu u^*$ , then  $g(\mu)$  is the y-intercept of the line with slope  $-\mu$  through  $(u^*, t^*)$



strong duality only, if  $G$  lies on one side of tangent in  $x^*$

# Lagrange duality: strong duality

## Strong and weak duality of value sets

• As in the example, set  $G = \{(f_0(x), f_1(x), \dots, f_m(x), h_1(x), \dots, h_p(x)) \mid x \in \mathcal{D}\}$

$$\Rightarrow p^* = \inf \{ t \mid (t, u, v) \in G, u \leq 0, v = 0 \}$$

• Consider  $(1, \mu, \lambda)^T (t, u, v) = t + \mu^T u + \lambda^T v$

$$\Rightarrow g(\mu, \lambda) = \inf \{ (1, \mu, \lambda)^T (t, u, v) \mid (t, u, v) \in G \}$$

If infimum is finite,  $(t, u, v) \mapsto (1, \mu, \lambda)^T (t, u, v) = g(\mu, \lambda)$  defines a hyperplane orthogonal to  $(1, \mu, \lambda)$  which is tangent to  $G$  and such that  $G$  lies on one side of it.

$$\begin{aligned} d^* &= \sup_{\lambda, \mu \geq 0} g(\mu, \lambda) = \sup_{\lambda, \mu \geq 0} \inf \{ (1, \mu, \lambda)^T (t, u, v) \mid (t, u, v) \in G \} \\ &\leq \inf \{ t \mid (t, u, v) \in G, u \leq 0, v = 0 \} = p^* \end{aligned}$$


if  $\mu \geq 0$  &  $u \leq 0, v = 0$ , then  $t \geq (1, \mu, \lambda)^T (t, u, v)$

$\Rightarrow$  weak duality, and strong duality iff  $\mu^T u = 0$  at optimum

# Lagrange duality: strong duality

## Saddle point interpretation

Def: Let  $f: W \times Z \rightarrow \mathbb{R}$ .  $(\bar{w}, \bar{z}) \in W \times Z$  is called a saddle point if

$$f(\bar{w}, z) \leq f(\bar{w}, \bar{z}) \leq f(w, \bar{z}) \quad \forall (w, z) \in W \times Z, \text{ i.e.}$$
$$f(\bar{w}, \bar{z}) = \inf_{w \in W} f(w, \bar{z}) = \sup_{z \in Z} f(\bar{w}, z).$$


Thm:  $\sup_{z \in Z} \inf_{w \in W} f(w, z) \leq \inf_{w \in W} \sup_{z \in Z} f(w, z)$

proof:  $\sup_{z \in Z} \inf_{w \in W} f(w, z) \leq \sup_{z \in Z} \inf_{w \in W} \sup_{\tilde{z} \in Z} f(w, \tilde{z}) = \inf_{w \in W} \sup_{\tilde{z} \in Z} f(w, \tilde{z}) \quad \square$

Def:  $f$  possesses the saddle point property if  $\sup_z \inf_w f(w, z) = \inf_w \sup_z f(w, z)$ .

Thm: Strong duality  $\Leftrightarrow L$  satisfies the saddle point property on  $\mathbb{R}^n \times \mathbb{R}_+^m \times \mathbb{R}^p$

proof:  $p^* = \inf_{x, c_I(x) \leq 0, c_E(x) = 0} f_0(x) = \inf_x \sup_{\mu \geq 0, \lambda} L(x, \mu, \lambda)$

$d^* = \sup_{\mu \geq 0, \lambda} g(\mu, \lambda) = \sup_{\mu \geq 0, \lambda} \inf_x L(x, \mu, \lambda) \quad \square$

$\Rightarrow$  optimum is a saddle of  $L$ !

# Lagrange duality: optimality conditions

## Certificates

Notice: if  $x$  feasible and  $(\mu, \lambda)$  dual-feasible, then  
 $f_0(x) - p^* \leq f_0(x) - g(\mu, \lambda) =: \varepsilon$ , i.e.  
 $x$  is  $\varepsilon$ -suboptimal.

In particular, if  $f_0(x) = g(\mu, \lambda)$ , then  $x$  is optimal.

In general,  $p^*, d^* \in [g(\mu, \lambda), f_0(x)]$

Def:  $(\mu, \lambda)$  is called a certificate that  $x$  is  $\varepsilon$ -suboptimal/optimal.

This can be exploited as stopping criterion for optimisation algorithms!

# Lagrange duality: optimality conditions

## Complementarity

Assume strong duality, and let  $x^*$  primal-optimal and  $(\mu^*, \lambda^*)$  dual-optimal

$$f_0(x^*) = p^* = d^* = g(\mu^*, \lambda^*) = \inf_x \{ f_0(x) + \mu^{*\top} c_I(x) + \lambda^{*\top} c_E(x) \}$$

(A)  $\leq$   $f_0(x^*) + \mu^{*\top} c_I(x^*) + \lambda^{*\top} c_E(x^*)$  (B)  $\leq f_0(x^*)$ ,  $\mu^* \geq 0$

thus, equality holds in (A) and (B)

Equality in (A)  $\Rightarrow x \mapsto L(x, \lambda^*, \mu^*)$  is minimised by  $x^*$  (saddle point property)

Equality in (B)  $\Rightarrow \mu^{*\top} c_I(x^*) = 0 \Rightarrow \mu_i^* f_i(x^*) = 0, i=1, \dots, m.$

$\Rightarrow \mu_i^* = 0$  or  $f_i(x^*) = 0$  (complementary slackness)

# Lagrange duality: optimality conditions

## KKT conditions from strong duality

Thm: Consider  $\min_{x \in \mathbb{R}^n} f_0(x)$  s.t.  $c_I(x) \leq 0, c_E(x) = 0$  with  
 $f_0, c_E, c_I$  differentiable (i.e.  $\mathcal{X} = \bigcap_{i=0}^m \text{dom } f_i \cap \bigcap_{j=1}^p \text{dom } h_j$  open).

If strong duality holds and  $x^*$  is primal-,  $(\mu^*, \lambda^*)$  dual-optimal, then

before: / regularity

$$\left. \begin{aligned} c_I(x^*) \leq 0, c_E(x^*) = 0, \mu^* \geq 0, \mu^{*\top} c_I(x^*) = 0 \\ Df_0(x^*) + \mu^{*\top} Dc_I(x^*) + \lambda^{*\top} Dc_E(x^*) = 0 \end{aligned} \right\} \text{(KKT)}$$

proof: - saddle point property of  $L \Rightarrow x^*$  minimises  $x \mapsto L(x, \mu^*, \lambda^*)$   
 $\Rightarrow 0 = \frac{\partial}{\partial x} L(x, \mu^*, \lambda^*) = Df_0(x^*) + \mu^{*\top} Dc_I(x^*) + \lambda^{*\top} Dc_E(x^*)$   
 - complementary slackness  $\Rightarrow \mu^{*\top} c_I(x^*) = 0$  □

Thm: Let  $f_0, \dots, f_m$  convex,  $h_1, \dots, h_p$  affine.

Then (KKT)  $\Rightarrow x^*$  is primal-,  $(\mu^*, \lambda^*)$  dual-optimal.

proof: (KKT) implies feasibility of  $x^*, (\mu^*, \lambda^*)$ . Furthermore, due to  $\mu^* \geq 0$ ,  
 $x \mapsto \underbrace{f_0(x) + \mu^{*\top} c_I(x) + \lambda^{*\top} c_E(x)}_{L(x, \mu^*, \lambda^*)}$  is convex  $\stackrel{\text{(KKT)}}{\Rightarrow} x^*$  minimises  $L(x, \mu^*, \lambda^*)$   
 $\Rightarrow g(\mu^*, \lambda^*) = L(x^*, \mu^*, \lambda^*) = f_0(x^*)$ . □



## Motivation

- Several practical optimisation problems are convex
  - image processing / inverse problems ( $L^1$ -type optimisations)
  - production planning: linearised constraints + bounds  $\Rightarrow$  LP
- Sometimes the convex relaxation (convex envelope) of a nonconvex optimisation problem can be found
- Convex optimisation problems are simpler to solve (at each point clear what to go next)
- Convex optimisation problems satisfy many fine properties
  - local optimum = global optimum
  - often strong duality
- Convex optimisation problems occur naturally, e.g. as dual problem.

# Convex analysis: convex sets

## Definition & examples of convex sets

Def: A set  $C \subset \mathbb{R}^n$  is called convex if for all  $x, y \in C$  the line segment in between is also contained in  $C$ ,  $\theta x + (1-\theta)y \in C \quad \forall x, y \in C, \theta \in [0, 1]$



Ex: • Line: given  $x, y \in \mathbb{R}^n$ ,  $L = \{z \in \mathbb{R}^n \mid \exists \theta \in \mathbb{R} : z = \theta x + (1-\theta)y\}$  is convex

• Affine sets:  $C$  is called affine if for all  $x, y \in C$  also the line through  $x$  and  $y$  lies in  $C$ . If  $C$  is affine and  $x_0 \in C$ , then  $V = \{z \mid z + x_0 \in C\}$  is a linear subspace.

→ e.g. the solution set to a linear equation,  $C = \{x \mid Ax = b\}$ ,  $A \in \mathbb{R}^{k \times n}$ ,  $b \in \mathbb{R}^k$

→ affine hull of  $C \subset \mathbb{R}^n$ :  $\text{aff}(C) = \{y = \theta_1 x_1 + \dots + \theta_k x_k \mid k \in \mathbb{N}, \theta_1 + \dots + \theta_k = 1, x_1, \dots, x_k \in C\}$  is the smallest affine set containing  $C$  (homework)

→ affine dimension of  $C$  =  $\dim(\text{aff}(C))$  (note: for  $S^n$  this is 2!)


## Examples II

Ex: - Hyperplane:  $H = \{x \mid a^T x = b\}$  for  $a \in \mathbb{R}^n, b \in \mathbb{R}$ , is affine with  $\text{codim} H = 1$

- Halfspace:  $H = \{x \mid a^T x \leq b\}, a \in \mathbb{R}^n, b \in \mathbb{R}$ , is convex 

- Cone:  $C$  is called a cone if for all  $x \in C$  and all  $\theta \geq 0$  also  $\theta x \in C$ .

$C$  is called a convex cone if it is additionally convex, i.e.  $\forall x, y \in C$ :

$\forall \theta_1, \theta_2 \geq 0: \theta_1 x + \theta_2 y \in C$  


The positive orthant  $\mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid x_1, \dots, x_n \geq 0\}$  is a convex cone.

- (Norm-)ball: Let  $\|\cdot\|$  be a norm on  $\mathbb{R}^n$  (see a little later).

The (norm-)ball  $\{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$  is convex

- m-ellipse: Let  $x_1, \dots, x_m \in \mathbb{R}^n$ . The m-ellipse

$\{x \mid |x - x_1| + \dots + |x - x_m| \leq 1\}$  is convex.



# Convex analysis: convex sets

## Convex combinations

Def: Let  $x_1, \dots, x_k \in \mathbb{R}^n$ ,  $\theta_1, \dots, \theta_k \geq 0$  with  $\theta_1 + \dots + \theta_k = 1$ .  $y = \sum_{i=1}^k \theta_i x_i$  is called a convex combination of the  $x_i$ .

Thm:  $G \subset \mathbb{R}^n$  is convex  $\Leftrightarrow$  all convex combinations of any  $x_1, \dots, x_k \in G$  are contained in  $G$ .

proof: " $\Leftarrow$ " The line segment between  $x, y \in G$  are exactly all convex comb. of  $x, y$ .

" $\Rightarrow$ " induction over  $k$ :  $k=2$  is just definition of convexity

induction step: Consider  $x_1, \dots, x_{k+1} \in G$ ,  $\theta_1, \dots, \theta_{k+1} \geq 0$ ,  $\theta_1 + \dots + \theta_{k+1} = 1$ .

Wlog let  $\beta = \sum_{i=1}^k \theta_i > 0$  and set  $\theta'_i = \theta_i / \beta$ , then  $\sum_{i=1}^k \theta'_i = 1$ .

Thus,  $y' = \sum_{i=1}^k \theta'_i x_i \in G$ .

Furthermore,  $\beta + \theta_{k+1} = 1$ , i.e.  $\sum_{i=1}^{k+1} \theta_i x_i = \beta y' + \theta_{k+1} x_{k+1} \in G$ ,

since it can be written as convex combination of 2 elements.  $\square$

Short: convex combinations of convex combinations are again convex comb.

# Convex analysis: convex sets

## Convex hull

Thm: Let  $\{C_i\}_{i \in I}$  be a family of convex sets, then  $\bigcap_{i \in I} C_i$  is convex.

proof: Let  $x, y \in \bigcap_{i \in I} C_i$ , then  $\forall i \in I$  we have  $x, y \in C_i$ , thus  $\theta x + (1-\theta)y \in C_i \quad \forall \theta \in [0, 1]$   
 $\Rightarrow \forall \theta \in [0, 1], \quad \theta x + (1-\theta)y \in \bigcap_{i \in I} C_i. \quad \square$

Def: The convex hull  $\text{conv } C$  of a set  $C \subset \mathbb{R}^n$  is the intersection of all convex sets containing  $C$ .

Thm:  $\text{conv } C = \left\{ y = \sum_{i=1}^k \theta_i x_i \mid k \in \mathbb{N}, x_1, \dots, x_k \in C, \theta_1, \dots, \theta_k \geq 0, \sum_{i=1}^k \theta_i = 1 \right\} =: T$

proof: " $\subset$ ":  $T$  is convex: Any  $x, y \in T$  are convex combs. of points in  $C$ , thus any convex comb. of  $x$  and  $y$  is so, too, and thereby lies in  $T$ .

Also,  $C \subset T$ , thus  $\text{conv } C \subset T$ .

" $\supset$ ": Let  $S'$  be convex with  $S' \supset C$ .

By the previous slide,  $S'$  contains all its convex combinations, in particular also the convex combinations of  $C$ .

$\Rightarrow S' \supset T \xrightarrow[\text{previous thm}]{} \text{conv } C \supset T. \quad \square$

# Convex analysis: convex sets

## Caratheodory's theorem

Thm (Caratheodory): Every  $x \in \text{conv } G$ ,  $G \subset \mathbb{R}^n$ , can be written as convex combination of  $n+1$  elements from  $G$ .

proof: Consider an arbitrary convex combination  $x = \sum_{i=1}^k \theta_i x_i$  for  $k > n+1$ .

Claim: Without changing  $x$  one can change the  $\theta_i$  s.t. one  $\theta_i$  is 0.

Indeed,  $\{x_2 - x_1, \dots, x_k - x_1\} \subset \mathbb{R}^n$  are linearly dependent, since  $k-1 > n$ .

$\Rightarrow$  There are  $(\beta_2, \dots, \beta_k) \in \mathbb{R}^{k-1} \setminus \{0\}$  s.t.  $0 = \sum_{i=2}^k \beta_i (x_i - x_1) = \sum_{i=2}^k \beta_i x_i - \left(\sum_{i=2}^k \beta_i\right) x_1$ .

Define  $\theta'_i = \theta_i - t^* \beta_i$  for  $t^* = \theta_{i^*} / \beta_{i^*}$ ,  $i^* = \underset{i=1, \dots, k}{\text{argmin}} \theta_i / |\beta_i| =: -\beta_n$

$\Rightarrow \theta'_i \geq 0$  and  $\theta'_i = 0$  for at least one  $i$ .

Furthermore,  $\sum_{i=1}^k \theta'_i = \sum_{i=1}^k \theta_i - t^* \sum_{i=1}^k \beta_i = 1$

and  $\sum_{i=1}^k \theta'_i x_i = \sum_{i=1}^k \theta_i x_i - t^* \sum_{i=1}^k \beta_i x_i = x$ .

$$= \underbrace{\sum_{i=1}^k \theta_i x_i}_{=x} - \underbrace{t^* \sum_{i=1}^k \beta_i (x_i - x_1)}_{= \sum_{i=2}^k \beta_i (x_i - x_1) = 0} = x$$

$\leadsto$  similar technique in simplex method!

# Convex analysis: convex sets

## Examples III: norm balls


Def: A map  $\|\cdot\|: \mathbb{R}^n \rightarrow [0, \infty)$  is called a norm if for all  $x, y \in \mathbb{R}^n, \alpha \in \mathbb{R}$

- (a)  $\|x\| \geq 0$  (b)  $\|x\| = 0 \Leftrightarrow x = 0$  (c)  $\|\alpha x\| = |\alpha| \|x\|$  (d)  $\|x+y\| \leq \|x\| + \|y\|$

Ex: • Euclidean norm  $\|x\|_2 = |x| = \sqrt{x^T x} = \sqrt{\sum_{i=1}^n x_i^2}$

•  $l_1$ -norm  $\|x\|_1 = \sum_{i=1}^n |x_i|$  "Manhattan norm"

•  $l_\infty$ -norm  $\|x\|_\infty = \max_{i=1, \dots, n} |x_i|$



Def: • The norm-ball  $B(\tau, x) = \{y \in \mathbb{R}^n \mid \|x-y\| \leq \tau\}$  is convex. (homework)

• The ellipsoid  $B_p(\tau, x) = \{y \in \mathbb{R}^n \mid (x-y)^T P (x-y) \leq \tau^2\}$ ,  $P$  pos. def. is convex.

$\|x\|_p = \sqrt{x^T P x}$  defines a norm.


• The norm cone  $C = \{(x, t) \in \mathbb{R}^n \times \mathbb{R} \mid \|x\| \leq t\}$  is a convex cone. "2<sup>nd</sup> order cone" for  $\|\cdot\|_2$

# Convex analysis: convex sets


## Examples IV: simplices

Def: - The polyhedron  $P = \{x \in \mathbb{R}^n \mid a_j^T x \leq b_j, j=1, \dots, m, c_j^T x = d_j, j=1, \dots, p\}$ ,  $a_j, c_j \in \mathbb{R}^n, b_j, d_j \in \mathbb{R}$  is the intersection of  $m$  half spaces and  $p$  hyperplanes and thus is convex.

Compact notation:  $P = \{x \mid Ax \leq b, Cx = d\}$



- A polytope is a bounded polyhedron.
- $k+1$  points  $x_0, \dots, x_k \in \mathbb{R}^n$  are called affinely independent if  $x_1 - x_0, \dots, x_k - x_0$  are linearly independent.
- A simplex  $G$  is the convex hull of  $k+1$  affinely indep. points,  $G = \text{conv}\{x_0, \dots, x_k\}$ .
- The probability simplex is  $\text{conv}\{e_1, \dots, e_n\} \subset \mathbb{R}^n$



Thm: Every simplex is a polyhedron.

# Convex analysis: separation theorem

## Topological notions

Def: • The closed convex hull  $\overline{\text{conv}} S$  of a set  $S \subset \mathbb{R}^n$  is the intersection of all closed convex sets containing  $S$

• Let  $C \subset \mathbb{R}^n$  be convex and non-empty. If the interior of  $C$  is non-empty,  $\text{aff } C = \mathbb{R}^n$ .

The relative interior of  $C \subset \mathbb{R}^n$ ,  $\text{relint } C$ , is the interior of  $C$  w.r.t. the topology of  $\text{aff } C$

$$x \in \text{relint } C \iff x \in \text{aff } C \text{ and } \exists \delta > 0: \text{aff } C \cap B(x, \delta) \subset C$$


• The relative boundary  $\text{relbdy } C$  is defined analogously.

Ex: •  $C = \{x \in \mathbb{R}^3 \mid x_3 = 0, x_1^2 + x_2^2 \leq 1\}$

$$\text{aff } C = \{x \in \mathbb{R}^3 \mid x_3 = 0\}$$

$$\text{relint } C = \{x \in \mathbb{R}^3 \mid x_1^2 + x_2^2 < 1, x_3 = 0\}$$

$$\text{relbdy } C = \{x \in \mathbb{R}^3 \mid x_1^2 + x_2^2 = 1, x_3 = 0\}$$



# Convex analysis: separation theorem

## Operations preserving convexity

Thm: The intersection of convex sets is convex (see earlier)

Thm: The Cartesian product  $G = G_1 \times \dots \times G_n$  of convex sets  $G_i \subset \mathbb{R}^{n_i}$  is convex.

proof: homework

Thm: Let  $A: \mathbb{R}^n \rightarrow \mathbb{R}^m$  affine, i.e.  $A(x) = Bx + c$  for  $B \in \mathbb{R}^{m \times n}$ ,  $c \in \mathbb{R}^m$ . Let  $C \subset \mathbb{R}^n$ ,  $D \subset \mathbb{R}^m$  convex.

Then  $A(C)$  and  $A^{-1}(D) = \{x \in \mathbb{R}^n \mid A(x) \in D\}$  are convex.

proof: • Let  $x, y \in C$ . The image of  $[x, y] = \{\theta x + (1-\theta)y \mid \theta \in [0, 1]\}$  is  $[A(x), A(y)] \Rightarrow A(C)$  convex.

• Let  $x, y \in A^{-1}(D)$ , then  $A([x, y]) = [A(x), A(y)] \subset D \Rightarrow [x, y] \subset A^{-1}(D) \Rightarrow A^{-1}(D)$  convex.  $\square$

Thm:  $G \subset \mathbb{R}^n$  convex  $\Rightarrow$   $\text{int} G$ ,  $\text{relint} G$ ,  $\bar{G}$  convex.

proof: homework

# Convex analysis: separation theorem

## Extreme points

Def: Let  $G \subset \mathbb{R}^n$  convex,  $G \neq \emptyset$ .  $x \in G$  is called extreme point if there are

no  $x_1, x_2 \in G$ ,  $x_1 \neq x_2$ , with  $x = \theta x_1 + (1-\theta)x_2$  for a  $\theta \in (0,1)$ .

• Set of extreme points =  $\text{ext } G$



extreme points

Cor: • One may set  $\theta = \frac{1}{2}$  in above definition.

•  $x \in \text{ext } G \iff G \setminus \{x\}$  convex

Ex: •  $G = B(0,1) = \{x \in \mathbb{R}^n \mid |x| \leq 1\} \implies$  all  $x \in \mathbb{R}^n$  with  $|x|=1$  are extreme points:

Let  $x = \theta x_1 + (1-\theta)x_2$ ,  $\theta = \frac{1}{2}$ ,  $x_1 \neq x_2$ ,  $x_1, x_2 \in G$ .

$$1 = |x|^2 = |\theta x_1 + (1-\theta)x_2|^2 = 2 \left( \theta^2 |x_1|^2 + (1-\theta)^2 |x_2|^2 \right) - |\theta x_1 - (1-\theta)x_2|^2 < 1.$$

$$\leq 2\theta^2 + 2(1-\theta)^2 = 1 \quad = 0 \text{ iff } x_1 = \frac{1-\theta}{\theta} x_2 = x_2$$

• The  $B_n$ -ball has finitely many extreme points.

•  $\text{ext } G = \{0\}$  for a convex cone  $G$  which is not a halfhyperplane

•  $\text{ext } G = \emptyset$  for affine sets or halfspaces  $G$ .

# Convex analysis: separation theorem

## Properties of extreme points


Thm: If  $G \neq \emptyset$  compact and convex, then  $\text{ext } G \neq \emptyset$ .

proof: Due to compactness,  $x \mapsto |x|^2$  attains its maximum on  $G$  at some  $\bar{x}$ .

Claim:  $\bar{x} \in \text{ext } G$ . Indeed, let  $\bar{x} = \frac{x_1 + x_2}{2}$  for  $x_1, x_2 \in G, x_1 \neq x_2$ , then

$$|\bar{x}|^2 = \left| \frac{x_1 + x_2}{2} \right|^2 = \frac{1}{2} (|x_1|^2 + |x_2|^2) - \left| \frac{x_1 - x_2}{2} \right|^2 < |\bar{x}|^2 \quad \downarrow \quad \square$$

Thm:  $\text{ext } G \subset \text{relbdy } G$ .



Thm (Minkowski): If  $G \neq \emptyset$  compact and convex, then  $G = \text{conv}(\text{ext } G)$ .

proof: later...

Cor (see Carathéodory's Thm): If in addition  $\text{affdim } C = k$ , then every point in  $G$  is the convex combination of at most  $k+1$  extreme points.

# Convex analysis: separation theorem

## Faces

Def: A nonempty convex  $F \subset C$  is called a face of  $C$ , if for any  $x_1, x_2 \in C$ ,  
 $\text{relint}[x_1, x_2] \cap F \neq \emptyset \Rightarrow [x_1, x_2] \subset F$ .

Ex:  $x \in \text{ext}(C) \Leftrightarrow \{x\}$  is a (0-dimensional) face of  $C$

• 1-dimensional faces = edges



• face of polyhedra

•  $C$  itself is a face for  $C$  closed and convex

Thm: Let  $F$  be a face of the convex  $C \subset \mathbb{R}^n$ .  $\text{ext}(F) \subset \text{ext}(C)$

proof: Assume  $x \in \text{ext}F \setminus \text{ext}C$ , i.e. there are  $x_1, x_2 \in C$ ,  $x_1 \neq x_2$ ,  $x = \frac{x_1 + x_2}{2}$ .


By definition of a face,  $[x_1, x_2] \subset F$ , i.e.  $x_1, x_2 \in F \Rightarrow x \notin \text{ext}F \downarrow$ .  $\square$

# Convex analysis: separation theorem

## Exposed faces

Def: - A hyperplane  $H_{s,r} = \{x \in \mathbb{R}^n \mid s^T x = r\}$  supports  $G \subset \mathbb{R}^n$ , if  $G$  completely lies in one of the halfspaces  $\{x \in \mathbb{R}^n \mid s^T x \leq r\}$  or  $\{x \in \mathbb{R}^n \mid s^T x \geq r\}$ .

- $H_{s,r}$  supports  $G$  in  $x \in G$ , if in addition  $x \in H_{s,r}$
- $F \subset G \subset \mathbb{R}^n$  is an exposed face if there is a supporting hyperplane  $H$  with  $F = C \cap H$ .
- Exposed point = 0-dimensional exposed face (corner)



Thm: An exposed face is a face.

proof: Let  $F = G \cap H_{s,r}$  for a supporting hyperplane  $H_{s,r}$  (wlog,  $G \subset \{x \in \mathbb{R}^n \mid s^T x \leq r\}$ ).

Let  $x_1, x_2 \in G$ ,  $\alpha \in (0,1)$ , with  $x = \alpha x_1 + (1-\alpha)x_2 \in F \subset H_{s,r}$ .

$\Rightarrow r = s^T(\alpha x_1 + (1-\alpha)x_2)$ . Wlog assume  $x_1 \notin F \subset H_{s,r}$  i.e.  $s^T x_1 < r$ .

$\Rightarrow s^T(\alpha x_1 + (1-\alpha)x_2) < r \quad \Downarrow \quad \Rightarrow x_1, x_2 \in F \Rightarrow [x_1, x_2] \in F. \quad \square$

Cor: For  $C \subset \mathbb{R}^n$  convex,  $F$  an exposed face,  $\text{ext}(F) \subset \text{ext}(C)$  (by previous slide).

# Convex analysis: separation theorem

## Projections

Def: • A linear map  $P: \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called a projection, if  $P^2 = P$  (idempotence).  
• Let  $V \subset \mathbb{R}^n$  be a subspace.  $P_V: \mathbb{R}^n \rightarrow V$ ,  $x \mapsto v$  for  $x = v + v^\perp \in V \oplus V^\perp$  is called the orthogonal projection onto  $V$ .

Thm:  $P_V$  is linear, symmetric, positive semi-definite, idempotent, non-expansive ( $\|P_V\|_2 \leq 1$ ),  $x = P_V(x) + P_{V^\perp}(x)$ .

How to generalise the orthogonal projection onto closed sets?

Thm:  $P_V(x) = \underset{y \in V}{\operatorname{argmin}} \frac{1}{2} \|y - x\|_2^2$ .

proof: Let  $x = v + v^\perp \in V \oplus V^\perp$ , then  $\frac{1}{2} \|y - x\|_2^2 = \frac{1}{2} \|y - v\|_2^2 + \frac{1}{2} \|v^\perp\|_2^2 \Rightarrow y = v$  is minimiser  $\square$

Thm: Let  $C \subset \mathbb{R}^n$  be closed,  $x \in \mathbb{R}^n$ , then  $y \mapsto f_x(y) = \frac{1}{2} \|y - x\|_2^2$  attains its minimum on  $C$ .

proof: Let  $c \in C$  and  $S = \{y \in \mathbb{R}^n \mid f_x(y) \leq f_x(c)\}$ .  $f_x$  is continuous,  $C \cap S$  is compact  $\Rightarrow$  by Weierstrass' thm,  $f_x$  has a minimum on  $C \cap S$  and thus on  $C$ .  $\square$

# Convex analysis: separation theorem

## Orthogonal projections

Def: Let  $G \subset \mathbb{R}^n$  closed and convex. The (nonlinear) map  $P_G: \mathbb{R}^n \rightarrow G$ ,

$P_G(x) = \underset{y \in G}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$ , is called the orthogonal projection onto  $G$ .


Thm:  $P_G$  is well-defined.

proof: Only uniqueness remains to be shown: Let  $y_1 \neq y_2$  be minimisers,  $y_0 = \frac{y_1 + y_2}{2} \in G$ .

$$\begin{aligned} f_x(y_0) - \frac{1}{2}(f_x(y_1) + f_x(y_2)) &= \frac{\|x\|^2 - x \cdot y_1 - x \cdot y_2 + \|y_0\|^2}{2} - \frac{\|x\|^2 - 2x \cdot y_1 + \|y_1\|^2 + \|x\|^2 - 2x \cdot y_2 + \|y_2\|^2}{4} \\ &= \frac{1}{8} (-\|y_1\|^2 - \|y_2\|^2 + 2y_1 \cdot y_2) = -\frac{1}{8} \|y_1 - y_2\|^2 < 0 \quad \square \end{aligned}$$

Thm:  $P_G \circ P_G = P_G$      $P_G$  linear  $\Leftrightarrow G$  is a subspace of  $\mathbb{R}^n$ .

Thm:  $y_x = P_G(x) \Leftrightarrow (x - y_x) \cdot (y - y_x) \leq 0 \quad \forall y \in G$ .



proof: " $\Rightarrow$ " for  $y \in G$ ,  $\alpha \in [0, 1]$  arbitrary, consider  $y_\alpha = y_x + \alpha(y - y_x) \in G$ . We have

$$\frac{1}{2} \|y_x - x\|^2 = f_x(y_x) \leq f_x(y_\alpha) = \frac{1}{2} \|y_x - x + \alpha(y - y_x)\|^2$$

$$\Rightarrow 0 \leq \alpha(y_x - x) \cdot (y - y_x) + \frac{\alpha^2}{2} \|y - y_x\|^2; \text{ now divide by } \alpha \text{ and let } \alpha \rightarrow 0.$$

" $\Leftarrow$ "  $\forall y \in G$ ;  $0 \geq (x - y_x) \cdot (y - y_x) = \|x - y_x\|^2 + (x - y_x) \cdot (y - x) \geq \|x - y_x\|^2 - \|x - y_x\| \|y - x\|$

$\Rightarrow$  either  $y_x = x$  or  $\|x - y_x\| \leq \|y - x\|$ . □

# Convex analysis: separation theorem

## Separation of convex sets

Thm: Let  $C \subset \mathbb{R}^n$  convex, closed,  $x \notin C$ . Then exists  $s \in \mathbb{R}^n$  s.t.  $s \cdot x > \sup_{y \in C} s \cdot y$ .



halfspace H


proof: Set  $s := x - P_C(x)$ .

$$\text{For all } y \in C, 0 \geq (x - P_C(x)) \cdot (y - P_C(x)) = s \cdot (y - x + s) = s \cdot y - s \cdot x + \|s\|^2 \Rightarrow s \cdot y < s \cdot x \quad \square$$

Cor: Let  $C_1, C_2 \subset \mathbb{R}^n$  closed, convex, nonempty, disjoint. If  $C_2$  is bounded,

there exists  $s \in \mathbb{R}^n$  s.t.

$$\sup_{y \in C_1} s \cdot y < \min_{y \in C_2} s \cdot y$$



proof:  $C_1 - C_2 = \{y_1 - y_2 \mid y_1 \in C_1, y_2 \in C_2\}$  is convex:

indeed,  $C_1 \times C_2$  is convex, and  $C_1 - C_2 = i(C_1 \times C_2)$  for the linear  $i: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n, (a, b) \mapsto a - b$

$C_1 - C_2$  is closed, since  $C_1$  is closed and  $C_2$  compact:

indeed, let  $y_k = y_k^1 - y_k^2 \in C_1 - C_2$  with  $y_k \rightarrow y$ .  $C_2$  compact  $\Rightarrow y_k^2 \rightarrow y^2 \in C_2$   
up to subsequence

$\Rightarrow y_k^1 = y_k + y_k^2 \rightarrow y + y^2 \in C_1$  (since  $C_1$  closed)  $\Rightarrow y \in C_1 - C_2$

previous thm for  $x=0, C=C_1 - C_2 \Rightarrow \exists s \in \mathbb{R}^n: 0 = s \cdot 0 > \sup_{y \in C_1 - C_2} s \cdot y = \sup_{y_1 \in C_1} s \cdot y_1 - \min_{y_2 \in C_2} s \cdot y_2 \quad \square$

Note: statement can be wrong for  $C_1, C_2$  unbounded



if  $C_1, C_2$  only convex, nonempty, disjoint, one still gets  $\sup_{C_1} s \cdot y \leq \inf_{C_2} s \cdot y$  (homework)


# Convex analysis: separation theorem

## Consequences of separation property: Supporting hyperplanes

Thm (existence of supporting hyperplane): Let  $G \subset \mathbb{R}^n$  convex,  $x \in \partial G \cap G$ . There exists a supporting hyperplane to  $G$  at  $x$ ,

proof: Consider a sequence  $x_k \notin \bar{G}$  with  $x_k \rightarrow x$ . For each  $x_k$  there is  $s_k \in \mathbb{R}^n$  with  $s_k \cdot (x_k - y) > 0 \quad \forall y \in G$ . Wlog,  $\|s_k\|_2 = 1 \Rightarrow$  for a subsequence,  $s_k \rightarrow s \in S^{n-1}$ .  
By continuity  $s \cdot (x - y) \geq 0 \quad \forall y \in G \Rightarrow H_{s,r} = \{y \mid s \cdot y = r\}$  is the sought hyperplane for  $r = s \cdot x$ . □

Thm: For  $S \subset \mathbb{R}^n$  and halfspaces  $H_{s,r}^- = \{y \in \mathbb{R}^n \mid s \cdot y \leq r\}$ , set  $\Sigma_S = \{(s,r) \in \mathbb{R}^n \times \mathbb{R} \mid S \subset H_{s,r}^-\}$  and  $C_S = \bigcap_{(s,r) \in \Sigma_S} H_{s,r}^-$ . Then either  $C_S = \overline{\text{conv } S}$  or  $\overline{\text{conv } S} = \mathbb{R}^n$ .



proof: • Assume  $\overline{\text{conv } S} \neq \mathbb{R}^n$ , then  $C_S \supset \overline{\text{conv } S}$ .

• Now let  $x \notin \overline{\text{conv } S}$ . Separate  $\{x\}$  and  $\overline{\text{conv } S}$  by a hyperplane  $H_{s_0, r_0}$ , i.e.

$s_0 \cdot x > \sup_{y \in \overline{\text{conv } S}} s_0 \cdot y = r_0 \Rightarrow (s_0, r_0) \in \Sigma_S$ , but  $x \notin H_{s_0, r_0}^- \Rightarrow x \notin C_S$ . □

# Convex analysis: separation theorem

## Consequences of separation property: halfspaces & extr. points

Cor: Let  $C \subseteq \mathbb{R}^n$  closed convex, then  $C = C_G$ , i.e.  $C$  is intersection of halfspaces.

In particular, a polyhedron is the intersection of finitely many halfspaces.

Thm (Minkowski): If  $C \subset \mathbb{R}^n$  compact, convex, nonempty, then  $C = \text{conv}(\text{ext } C)$ .

proof: Induction in  $k = \text{affdim } C$ ; case  $k=0$  is trivial. Induction step  $k-1 \rightarrow k$ :

Let  $x \in C$ . • case  $x \in \text{relbdy } C$ : There is a hyperplane  $H \not\subset C$  supporting  $C$  in  $x$ .

$\Rightarrow C \cap H$  has affine dimension  $\leq k-1$

why?

$\Rightarrow x$  is convex combination of extreme points  $x_i$  in  $C \cap H$

$x_i$  are also extreme points of  $C$ , since  $C \cap H$  is an exposed face.

• case  $x \in \text{relint } C$ : Choose  $x' \neq x$ ,  $x' \in C$

$\Rightarrow$  line through  $x, x'$  intersects  $\text{relbdy } C$  in two points  $y, z$ .

$\Rightarrow$  by first case,  $y, z \in \text{conv}(\text{ext } C)$

$\Rightarrow x \in \text{conv}(\text{ext } C)$

□

# Convex analysis: convex functions

## Basic notions

Def: Let  $C \subset \mathbb{R}^n$  convex.  $f: C \rightarrow \mathbb{R} \cup \{\infty\}$  is called convex ( $f \in \text{Conv } C$ ), if

$$\forall x, y \in C, \theta \in [0, 1]: f(\theta x + (1-\theta)y) \leq \theta f(x) + (1-\theta)f(y).$$



•  $f$  is strictly convex, if  $\forall x \neq y \in C, \theta \in (0, 1): f(\theta x + (1-\theta)y) < \theta f(x) + (1-\theta)f(y)$

•  $f$  is concave if  $-f$  is convex.

• The graph of  $f$  is the set  $\{(x, f(x)) \mid x \in \text{dom } f\}$ .

• The epigraph of  $f$  is  $\text{epi } f = \{(x, \tau) \in \mathbb{R}^n \times \mathbb{R} \mid f(x) \leq \tau\}$ .



• The sublevel set of  $f$  wrt  $\tau$  is  $S_\tau(f) = \{x \in \mathbb{R}^n \mid f(x) \leq \tau\}$ .



Thm: •  $f$  convex  $\Rightarrow \text{dom } f$  convex

•  $f$  convex  $\Leftrightarrow \text{epi } f \subset \mathbb{R}^n \times \mathbb{R}$  convex

•  $(x, \tau) \in \text{epi } f \Leftrightarrow x \in S_\tau(f)$

# Convex analysis: convex functions

## Examples & properties

Ex: • characteristic or indicator function of a convex  $C \subset \mathbb{R}^n$

$$I_C(x) = \begin{cases} 0 & \text{if } x \in C \\ \infty & \text{else} \end{cases}$$

different from  $\chi_C = \begin{cases} 1 & \text{if } x \in C \\ 0 & \text{else} \end{cases} !$

- Any norm on  $\mathbb{R}^n$  is convex:  $\forall x, y \in \mathbb{R}^n, \theta \in (0, 1): \|\theta x + (1-\theta)y\| \leq \|\theta x\| + \|(1-\theta)y\| = \theta\|x\| + (1-\theta)\|y\|$
- The maximum function  $f(x) = \max_i x_i$  is convex.
- Linear and affine functions are convex; their epigraph is a halfspace.

Thm (Jensen's inequality):  $\forall f \in \text{Conv } \mathbb{R}^n, x_1, \dots, x_k \in \text{dom } f, \theta_1, \dots, \theta_k \geq 0, \sum_{i=1}^k \theta_i = 1: f\left(\sum_{i=1}^k \theta_i x_i\right) \leq \sum_{i=1}^k \theta_i f(x_i)$

proof:  $(x_i, f(x_i)) \in \text{epi } f \implies \sum_{i=1}^k \theta_i (x_i, f(x_i)) = \left(\sum_{i=1}^k \theta_i x_i, \sum_{i=1}^k \theta_i f(x_i)\right) \in \text{epi } f$ .  $\square$   
 $\uparrow$   
epi  $f$  convex

Rem: Jensen's inequality also holds for integrals: Let  $f \in \text{Conv } \Omega$  and

$(\Omega, \Sigma, P)$  be a probability space, then  $f\left(\int_{\Omega} x dP\right) \leq \int_{\Omega} f(x) dP$ .

expectation of random variable  $x$       exp. of  $f(x)$

# Convex analysis: convex functions

## Convexity and continuity

Thm: Let  $f \in \text{Conv } \mathbb{R}^n$ .  $f$  is continuous on  $\text{int}(\text{dom } f)$ . necessary?!

proof: • Let  $\bar{x} \in \text{int } \text{dom } f$ ;  $f$  is bounded in a neighbourhood of  $\bar{x}$ .

Indeed, there is a simplex  $Q = \text{conv} \{x_0, \dots, x_n\} \subset \text{dom } f$  with  $\bar{x} \in \text{int } Q$ .

Let  $x \in Q$ , i.e.  $x = \sum_{i=1}^n \theta_i x_i$ ,  $\theta_i \in [0, 1]$ ,  $\sum_{i=1}^n \theta_i = 1$ , then  $f(x) \leq \sum_{i=1}^n \theta_i f(x_i) \leq \max_i f(x_i) =: \tilde{M}$ .

Also, for  $\alpha = \frac{\text{dist}(\bar{x}, \partial Q)}{2 \text{diam}(Q)} \in (0, \frac{1}{2})$  set  $\hat{x} = \bar{x} - \frac{\alpha}{1-\alpha} (x - \bar{x}) \in Q$  (hence  $\bar{x} = \alpha x + (1-\alpha)\hat{x}$ )

$$\Rightarrow f(\bar{x}) \leq \alpha f(x) + (1-\alpha) f(\hat{x}) \Rightarrow f(x) \geq \frac{f(\bar{x}) - (1-\alpha) f(\hat{x})}{\alpha} \geq \min(-\tilde{M}, \frac{f(\bar{x}) - \tilde{M}}{\alpha}) =: -M$$

• Let  $v \in \mathbb{R}^n$  s.t.  $\bar{x} + v \in Q$ . For  $\beta \in [0, 1]$  we have

$$f(\bar{x} + \beta v) - f(\bar{x}) \leq \beta f(\bar{x} + v) + (1-\beta) f(\bar{x}) - f(\bar{x}) \leq \beta (M - f(\bar{x}))$$

• For  $v \in \mathbb{R}^n$  small enough s.t.  $\bar{x} \pm v \in Q$ ,  $\beta \in [0, 1]$  we have

$$f(\bar{x} + \beta v) - f(\bar{x}) \geq f(\bar{x} + \beta v) - \left[ \frac{\beta}{1+\beta} f(\bar{x} - v) + \frac{1}{1+\beta} f(\bar{x} + \beta v) \right] = \frac{-\beta}{1+\beta} (f(\bar{x} + \beta v) - f(\bar{x} - v)) \geq \frac{-2M\beta}{1+\beta}$$

• Summarising,  $|f(\bar{x} + \beta v) - f(\bar{x})| \leq 2\beta M$  for all  $v$  small enough and  $\beta \in [0, 1]$

$\Rightarrow f$  is locally Lipschitz continuous. □


Note: Not true in general  $\alpha$ -dimensional spaces (e.g.  $\|\cdot\|_{H^1} \in \text{Conv}(H^1)$  with norm  $\|\cdot\|_{L^2}$ ); also there exist discontinuous linear functionals on  $\alpha$ -dimensional Banach spaces)

# Convex analysis: convex functions

## Convex and affine functions

Thm: Let  $f \in \text{Conv } \mathbb{R}^n$ .  $\forall x_0 \in \text{relint dom } f$   $\exists s \in \mathbb{R}^n$ :  $f(x) \geq f(x_0) + s \cdot (x - x_0) \forall x \in \mathbb{R}^n$  necessary ?!

proof: • Note  $\text{aff epi } f = (\text{aff dom } f) \times \mathbb{R}$   
and  $\text{aff dom } f = x_0 + V$  for a subspace  $V \subset \mathbb{R}^n$ .



- $(x_0, f(x_0)) \in \text{relbdy}(\text{epi } f) \Rightarrow \exists$  hyperplane supporting  $\text{epi } f$  in  $(x_0, f(x_0))$ , i.e.  $\exists \tilde{s} \in V, \alpha \in \mathbb{R}, (\alpha \neq 0)$  s.t.  $\begin{pmatrix} \tilde{s} \\ \alpha \end{pmatrix} \cdot \begin{pmatrix} x \\ r \end{pmatrix} \leq \begin{pmatrix} \tilde{s} \\ \alpha \end{pmatrix} \cdot \begin{pmatrix} x_0 \\ f(x_0) \end{pmatrix} \quad \forall (x, r) \in \text{epi } f$ ,  
in particular  $-\alpha r \geq -\alpha f(x_0) + \tilde{s} \cdot (x - x_0)$  for all  $x \in \text{dom } f, r = f(x)$
- $\alpha < 0$ . Indeed, letting  $r \rightarrow \infty$  shows  $\alpha \leq 0$ ,  
and  $\alpha = 0$  implies  $0 \geq \tilde{s} \cdot (x - x_0) \forall x \in \text{dom } f$ , i.e.  $\tilde{s} = 0$ , a contradiction.
- Dividing by  $-\alpha$ ,  $f(x) \geq f(x_0) + s \cdot (x - x_0)$  for  $s = \frac{\tilde{s}}{-\alpha}$ .  $\square$

What is  $s$  if  $f$  is differentiable in  $x_0$ ?

# Convex analysis: convex functions

## Differentiable convex functions

Thm: Let  $C \subset \mathbb{R}^n$  open, convex,  $f: C \rightarrow \mathbb{R}$  differentiable.

$f$  is (strictly) convex on  $C \iff f(x) \geq (>) f(x_0) + \nabla f(x_0) \cdot (x - x_0) \forall x, x_0 \in C$

proof: " $\implies$ ": Let  $f$  be (strictly) convex,  $\alpha \in (0, 1)$ , then

$$f(x_0 + \alpha(x - x_0)) - f(x_0) = f(\alpha x + (1 - \alpha)x_0) - f(x_0) \leq \alpha f(x) + (1 - \alpha)f(x_0) - f(x_0) = \alpha(f(x) - f(x_0))$$

Dividing by  $\alpha$  and letting  $\alpha \rightarrow 0$  yields  $\nabla f(x_0) \cdot (x - x_0) \leq f(x) - f(x_0)$ .

If equality holds and  $f$  is strictly convex, let  $z = \theta x_0 + (1 - \theta)x$ , then

$$f(z) < \theta f(x_0) + (1 - \theta)f(x) = f(x_0) + \nabla f(x_0) \cdot (x - x_0)(1 - \theta) = f(x_0) + \nabla f(x_0) \cdot (z - x_0) \downarrow$$

" $\impliedby$ ": Let  $x_1 \neq x_2 \in C$ ,  $\alpha \in (0, 1)$ ,  $x_0 = \alpha x_1 + (1 - \alpha)x_2 \in C$ .

$$(E_i) \quad f(x_i) \geq f(x_0) + \nabla f(x_0) \cdot (x_i - x_0)$$

$$\alpha(E_1) + (1 - \alpha)(E_2) : \quad \alpha f(x_1) + (1 - \alpha)f(x_2) \geq f(x_0) + \nabla f(x_0) \cdot \underbrace{(\alpha x_1 + (1 - \alpha)x_2 - x_0)}_{x_0}$$

$$\implies f(\alpha x_1 + (1 - \alpha)x_2) \leq \alpha f(x_1) + (1 - \alpha)f(x_2) \quad \square$$

# Convex analysis: convex functions

## Convexity and monotonicity of first derivative

Thm: Let  $C \subset \mathbb{R}^n$  convex,  $f: C \rightarrow \mathbb{R}$  differentiable. If  $Df$  is (strictly) monotone, i.e.  $(Df(x) - Df(y)) \cdot (x - y) \geq (>) 0 \quad \forall x \neq y \in C$ , then  $f$  is (strictly) convex.

proof: Let  $x_1, x_2 \in C$ ,  $\alpha \in (0, 1)$ ,  $x = (1 - \alpha)x_1 + \alpha x_2 = x_1 + \alpha(x_2 - x_1) \in C$ ,  $x_s = x_1 + s(x_2 - x_1)$   
 $f(x) = f(x_1) + \int_0^\alpha Df(x_s)(x_2 - x_1) ds \quad (E_1)$       $f(x_2) = f(x_1) + \int_0^1 Df(x_s)(x_2 - x_1) ds \quad (E_2)$   
 $(1 - \alpha)(E_1) - \alpha(E_2) \Rightarrow f(x) = (1 - \alpha)f(x_1) + \alpha f(x_2) + \underbrace{[(1 - \alpha) \int_0^\alpha Df(x_s) ds - \alpha \int_\alpha^1 Df(x_s) ds]}_A (x_2 - x_1)$   
 $A = \alpha(1 - \alpha) \int_0^\alpha Df(x_{\alpha s}) - Df(x_{\alpha + (1 - \alpha)s}) ds (x_2 - x_1) \stackrel{A}{\leq} (<) 0. \quad \square$

Thm: Let  $C \subset \mathbb{R}^n$  convex,  $f: C \rightarrow \mathbb{R}$  differentiable.  $f$  (strictly) convex  $\Rightarrow Df$  (strictly) monotone.

proof:  $f(x_1) \stackrel{>}{\leq} f(x_2) + Df(x_2)(x_1 - x_2)$      &      $f(x_2) \stackrel{>}{\leq} f(x_1) + Df(x_1)(x_2 - x_1)$ .  
Now sum both inequalities. □

Cor: Let  $C \subset \mathbb{R}^n$  convex,  $f: C \rightarrow \mathbb{R}$  twice differentiable. Then  $\Leftarrow$  is false in general!

(a)  $D^2f$  (strictly) positive definite  $\Rightarrow f$  (strictly) convex     (b)  $f$  convex  $\Rightarrow D^2f$  ps. semi-def

proof: (strict) positive definiteness of  $D^2f \Leftrightarrow \Rightarrow$  (strict) monotonicity of  $Df. \quad \square$

# Convex analysis: convex functions

## Operations preserving convexity

Thm: i) Let  $f_1, \dots, f_m \in \text{Conv } \mathbb{R}^n$ ,  $t_1, \dots, t_m \geq 0$ ,  $x_0 \in \mathbb{R}^n$  with  $f_1(x_0), \dots, f_m(x_0) < \infty$ .

Then  $\sum_{i=1}^m t_i f_i \in \text{Conv } \mathbb{R}^n$ .

ii) Let  $f \in \text{Conv } \mathbb{R}^n$ ,  $A: \mathbb{R}^n \rightarrow \mathbb{R}^n$  affine with  $\text{range } A \cap \text{dom } f \neq \emptyset$ .

Then  $f \circ A \in \text{Conv } \mathbb{R}^n$ .

iii) Let  $f_i \in \text{Conv } \mathbb{R}^n$ ,  $i \in J$ ,  $x_0 \in \mathbb{R}^n$  with  $f_i(x_0) < M < \infty \forall i \in J$ .

Then  $x \mapsto \sup_{i \in J} f_i(x) \in \text{Conv } \mathbb{R}^n$ .

iv) Let  $f \in \text{Conv } \mathbb{R}^n$ ,  $h \in \text{Conv } \mathbb{R}$  monotonically increasing,  $x_0 \in \mathbb{R}^n$  with  $f(x_0) \in \text{dom } h$ .  $h \circ f \in \text{Conv } \mathbb{R}^n$ .

v) Let  $f \in \text{Conv } \mathbb{R}^n \times \mathbb{R}^{n_2}$ ,  $C \subset \mathbb{R}^{n_2}$  convex,  $g(x) = \inf_{y \in C} f(x, y)$ . If  $g(x) > -\infty \forall x \in \mathbb{R}^n$ ,  $g \in \text{Conv } \mathbb{R}^n$ .

Example: Moreau-Yosida approximation for  $f \in \text{Conv } \mathbb{R}^n$ ,  $\lambda > 0$ :

$$f_\lambda(x) = \inf_{y \in \mathbb{R}^n} f(y) + \frac{1}{2\lambda} \|x - y\|_2^2.$$

proof: i), ii), iv) homework

iii) Let  $f(x) = \sup_{i \in J} f_i(x)$ .  $\text{epi } f = \bigcap_{i \in J} \text{epi } f_i$  is convex.

v)  $\text{epi } g = \{(x, \epsilon) \mid \exists y \in C: (x, y, \epsilon) \in \text{epi } f\} = \text{P}_{\mathbb{R}^n \times \mathbb{R}}(\text{epi } f)$  is convex.  $\square$

# Convex analysis: convex functions

## Semi-continuity

Def:  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$  is called lower semi-continuous if  $\liminf_{y \rightarrow x} f(y) \geq f(x) \forall x \in \mathbb{R}^n$ .

Ex:  $f(x) = \begin{cases} 1 & x < 0 \\ 0 & x \geq 0 \end{cases}$  is lower semi-continuous,  $f(x) = \begin{cases} 1 & x \leq 0 \\ 0 & x > 0 \end{cases}$  is not.



Thm: Let  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ . The following are equivalent:

(a)  $f$  is lower semi-continuous

(b)  $\text{epi } f$  is closed in  $\mathbb{R}^n \times \mathbb{R}$

(c) The sublevel sets  $S_r(f)$  are closed for all  $r \in \mathbb{R}$ .

proof: (a)  $\Rightarrow$  (b): Let  $(y_k, r_k) \in \text{epi } f, (y, r) = \lim_{k \rightarrow \infty} (y_k, r_k)$ .

$$r = \lim_{k \rightarrow \infty} r_k \geq \liminf_{k \rightarrow \infty} f(y_k) \geq \liminf_{x \rightarrow y} f(x) \geq f(y), \text{ hence } (y, r) \in \text{epi } f.$$

(b)  $\Rightarrow$  (c): Let  $A_r(x) = (x, r), Q_r = \text{epi } f \cap (\mathbb{R}^n \times \{r\})$ .  $Q_r$  is closed,  $A_r$  continuous.

$$S_r(f) = \{x \in \mathbb{R}^n \mid f(x) \leq r\} = A_r^{-1}(Q_r) \text{ is closed.}$$



(c)  $\Rightarrow$  (a): Assume (a) is false, i.e.  $\exists y_k \rightarrow x$  with  $\rho := \lim_{k \rightarrow \infty} f(y_k) < f(x)$ . Let  $r \in (\rho, f(x))$

For  $k > k_0$  large enough,  $f(y_k) \leq r < f(x)$ , i.e.  $y_k \in S_r(f) \forall k > k_0$ , but  $x \notin S_r(f) \nmid \square$

# Convex analysis: convex functions

## Closed convex functions

Def: - A function is called closed if it is lsc on  $\mathbb{R}^n$  (or if  $\text{epi } f$  is closed).

• The relaxation (or lower semi-continuous envelope) of a function  $f$  is defined via  $\bar{f}(x) := \text{cl } f(x) := \liminf_{y \rightarrow x} f(y)$  or  $\text{epi } \bar{f} = \overline{\text{epi } f}$

Thm:  $f \in \text{Conv } \mathbb{R}^n \Rightarrow \bar{f} \in \text{Conv } \mathbb{R}^n$ , and  $f = \bar{f}$  on  $\text{reliant dom } f$ .

proof: •  $f$  convex  $\Rightarrow \text{epi } f$  convex  $\Rightarrow \text{epi } \bar{f} = \overline{\text{epi } f}$  convex  $\Rightarrow \bar{f}$  convex.

Also,  $\bar{f} \leq f \not\equiv \infty$  and  $\bar{f} > -\infty \forall x$  (due to minorisation by affine fns.)

•  $f|_{\text{aff dom } f}$  is continuous on  $\text{reliant dom } f$ ,

thus  $\bar{f}|_{\text{aff dom } f} = \overline{f|_{\text{aff dom } f}} = f|_{\text{aff dom } f}$  on  $\text{reliant dom } f$

$\Rightarrow f = \bar{f}$  on  $\text{reliant dom } f$ . □

# Convex analysis: conjugate functions

## The conjugate function

Def: The conjugate function (Legendre-Fenchel dual) to a function  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$  (not necessarily convex) is defined as

$$f^*(s) = \sup_{x \in \mathbb{R}^n} s \cdot x - f(x), \quad s \in \mathbb{R}^n$$

• The map  $f \mapsto f^*$  is the Legendre-Fenchel-transform



Thm: (Fenchel-Young-inequality)  $f^*(y) + f(x) \geq x \cdot y \quad \forall x, y \in \mathbb{R}^n$

Ex:  $f(x) = \begin{cases} x \log x - x, & x \geq 0 \\ \infty & \text{else} \end{cases}, \quad f^*(y) = e^y, \quad \text{Young-inequality } xy \leq x \log x - x + e^y \quad \forall x \geq 0, y \in \mathbb{R}$

Thm: Let  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ ,  $f \neq \infty$ ,  $f \geq g$  for an affine function  $g$ .  $f^*$  is closed convex.

proof: •  $f^*$  is supremum of affine functions  $y \mapsto f_x(y) = y \cdot x - f(x) \Rightarrow f^*$  is convex  
•  $\text{epi } f^* = \bigcap_{x \in \mathbb{R}^n} \text{epi } f_x$  is closed as intersection of half spaces  $\Rightarrow f^*$  is closed  
•  $f^* \neq \infty$ : Indeed, let  $g(x) = s_0 \cdot x + r_0$ , then  $f^*(s_0) = \sup_x s_0 \cdot x - f(x) \leq \sup_x s_0 \cdot x - g(x) = -r_0 \quad \square$

# Convex analysis: conjugate functions

## The biconjugate

Thm: Let  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ ,  $f \not\equiv \infty$ ,  $f(x) \geq s_0 \cdot x + r_0$  for some  $(s_0, r_0) \in \mathbb{R}^n \times \mathbb{R}$ .

We have  $\text{epi } f^{**} = \overline{\text{conv epi } f}$ , i.e.  $f^{**}$  is largest closed convex function below  $f$  ("convex relaxation", "convex, lower semi-continuous envelope").

Cor:  $f^{**} = f \Leftrightarrow f$  is closed convex



proof: Let  $\Sigma = \{(s, r) \in \mathbb{R}^n \times \mathbb{R} \mid f(x) \geq s \cdot x - r \ \forall x\}$  = parameters of supp. hyperplanes of  $\text{epi } f$

$$\swarrow \begin{pmatrix} s \\ -1 \end{pmatrix} \cdot \begin{pmatrix} x \\ f(x) \end{pmatrix} \leq r$$

$f^{**}(x) = \sup_{(s,r) \in \Sigma} (s \cdot x - r) = \sup$  over all supp. hyperplanes of their vertical coordinate at  $x$

Indeed,  $(s, r) \in \Sigma \Leftrightarrow r \geq \sup_{x \in \mathbb{R}^n} s \cdot x - f(x) = f^*(s)$ .

Thus,  $\sup_{(s,r) \in \Sigma} (s \cdot x - r) = \sup_{s, r \text{ with } f^*(s) \leq r} s \cdot x - r = \sup_s s \cdot x - f^*(s) = f^{**}(x)$ .

$\text{epi } f^{**} = \{(x, t) \mid f^{**}(x) \leq t\} = \{(x, t) \mid \sup_{(s,r) \in \Sigma} (s \cdot x - r) \leq t\} = \bigcap_{(s,r) \in \Sigma} \{(x, t) \mid s \cdot x - r \leq t\}$   
 = intersection of all half spaces containing  $\text{epi } f = \overline{\text{conv epi } f}$   $\square$

# Convex analysis: conjugate functions

## Examples

Ex: if  $f$  convex & differentiable,  $\text{dom } f = \mathbb{R}^n$ :

$$x^* = \underset{x}{\text{argmax}} \ y \cdot x - f(x) \Leftrightarrow y = \nabla f(x^*) \quad \Rightarrow \quad f^*(y) = x^* \cdot \nabla f(x^*) - f(x^*)$$

Let  $Q \in \mathbb{R}^{n \times n}$  symmetric positive definite,  $b \in \mathbb{R}^n$ ,  $f(x) = \frac{1}{2} x^T Q x + b^T x$ .

$$f^*(y) = \max_x \ y^T x - \frac{1}{2} x^T Q x - b^T x$$

$$y = \nabla f(x^*) = Q x^* + b \Rightarrow x^* = Q^{-1}(y - b) \Rightarrow f^*(y) = \frac{1}{2} (y - b)^T Q^{-1} (y - b)$$

Special case  $Q = I$ ,  $b = 0$ , i.e.  $f(x) = \frac{1}{2} \|x\|_2^2 \Rightarrow f^*(y) = \frac{1}{2} \|y\|_2^2$

Fenchel's inequality:  $x^T Q x + y^T Q^{-1} y \geq 2x \cdot y$  (case  $b = 0$ )

$$\|x\|^2 + \|y\|^2 \geq 2x \cdot y \quad (\text{case } b = 0, Q = I)$$

$f(x) = I_C(x) = \begin{cases} 0 & \text{if } x \in C \subset \mathbb{R}^n \\ \infty & \text{else} \end{cases} \Rightarrow I_C^*(y) = \sup_{x \in C} y \cdot x =: \zeta_C(y)$

"indicator function"

"support function"

special case  $C = \text{subspace of } \mathbb{R}^n$ :  $\zeta_C(y) = I_{C^\perp}(y)$

$C = \mathcal{B}(0, 1) = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\} \Rightarrow \zeta_C(y) = \|y\|_*$  for the dual norm  $\|y\|_* = \sup_{\|x\| \leq 1} |x \cdot y|$

$\|\cdot\|_p$  &  $\|\cdot\|_q$  with  $\frac{1}{p} + \frac{1}{q} = 1$  are dual norms  $\Rightarrow \zeta_{\mathcal{B}(0, 1), \|\cdot\|_p} = \|\cdot\|_q$

# Convex analysis: conjugate functions

## Calculation rules

Thm: Let  $f, f_i : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ ,  $f, f_i \neq \infty$ ,  $f, f_i \geq g, g$ : for affine  $g, g_i$ :

$h(x)$	$h^*(y)$	$h(x)$	$h^*(y)$
$f(x) + r$	$f^*(y) - r$	$f(x) + y_0 \cdot x$	$f^*(y - y_0)$
$t f(x), t > 0$	$t f^*(\frac{y}{t})$	$\sum_{j=1}^m f_j(x_j), x = (x_1, \dots, x_m), x_j \in \mathbb{R}^{n_j}$	$\sum_{j=1}^m f_j^*(y_j), y = (y_1, \dots, y_m)$
$f(tx), t \neq 0$	$f^*(\frac{y}{t})$	$\leq f$	$\geq f^*$
$f(Ax), A \in \mathbb{R}^{n \times m}$ invertible	$f^*(A^{-T}y)$	$\alpha f_1 + (1-\alpha)f_2, \alpha \in [0, 1]$	$\leq \alpha f_1^* + (1-\alpha)f_2^*$
$f(x - x_0)$	$f^*(y) + y \cdot x_0$	<i>"convexity of conjugation"</i>	

Def: Given  $f, g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ , their infimal convolution is

$$(f \square g)(x) = \inf_{y \in \mathbb{R}^n} f(y) + g(x - y) = \inf_{y \in \mathbb{R}^n} g(y) + f(x - y)$$

Moreau - Yosida - approximation  $f_\lambda = f \square \frac{1}{2\lambda} \|\cdot\|_2^2$

Thm:  $(f \square g)^* = f^* + g^*$  careful: if  $f, g, f^* \square g^*$  closed convex, then  $(f + g)^* = f^* \square g^*$ , but not generally

proof:  $(f \square g)^*(y) = \sup_x x \cdot y - \inf_{x_1 + x_2 = x} (f(x_1) + g(x_2)) = \sup_{x_1, x_2} (x_1 + x_2) \cdot y - f(x_1) - g(x_2) = f^*(y) + g^*(y) \square$

# Convex analysis: conjugate functions

## Coercivity and the conjugate

Def: A function  $f$  is called coercive or 0-coercive if  $\lim_{\|x\| \rightarrow \infty} f(x) = \infty$

$f$  is called 1-coercive if  $\lim_{\|x\| \rightarrow \infty} \frac{f(x)}{\|x\|} = \infty$

Thm: Let  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ ,  $f \not\equiv \infty$ ,  $f(x) \geq y_0 \cdot x + r_0$ .  $f$  1-coercive  $\Rightarrow f^* < \infty$

proof: Let  $y \in \mathbb{R}^n$ . (a)  $\exists R > 0: f(x) \geq \|y\|_2 \|x\|_2 \quad \forall \|x\|_2 \geq R$

show  $f^*(y) < \infty!$

$$\Rightarrow \begin{array}{l} \text{Cauchy-Schwarz} \\ x \cdot y - f(x) \leq 0 \quad \forall \|x\|_2 \geq R \Rightarrow \sup_{\|x\|_2 \geq R} x \cdot y - f(x) \leq 0 \end{array}$$

$$(b) \sup_{\|x\|_2 < R} x \cdot y - f(x) \leq \sup_{\|x\|_2 < R} x \cdot y - y_0 \cdot x - r_0 \leq R \|y - y_0\|_2 - r_0$$

$$(a) \& (b) \Rightarrow f^*(y) \leq \max(0, R \|y - y_0\|_2 - r_0) \quad \square$$

Thm:  $F_1 := \{\text{support functions of convex sets}\} = \{\text{closed convex positively homogeneous fcn's}\} =: F_2$

proof:  $b \in F_1 \Leftrightarrow b = I_C^*$  for some convex  $C \Rightarrow b$  closed convex with  $b(\lambda x) = |\lambda| b(x) \quad \forall \lambda \geq 0 \Leftrightarrow b \in F_2$

$$b \in F_2 \Rightarrow b^*(y) = \sup_x x \cdot y - b(x) = \sup_{\lambda \geq 0, x} \lambda x \cdot y - b(\lambda x) = \sup_{\lambda \geq 0} \lambda \delta^*(y) = I_C(y)$$

for  $C = \{y \in \mathbb{R}^n \mid b^*(y) = 0\}$ ;  $C$  must be convex since  $b^*$  is  $\Rightarrow b \in F_1 \quad \square$

Cor:  $f \in \text{Conv } \mathbb{R}^n$  pos. homogeneous  $\Rightarrow \text{cl} f = b_C$  for the closed convex  $C = \{x \in \mathbb{R}^n \mid x \cdot y \leq f(y) \quad \forall y \in \mathbb{R}^n\}$

proof:  $\text{cl} f \in F_2 \Rightarrow \text{cl} f \in F_1 \Rightarrow \text{cl} f = b_C$  for some convex  $C \subset \mathbb{R}^n$


$$f^*(x) = b_C^*(x) = I_C(x) = \begin{cases} \infty & \text{if } \exists y \in \mathbb{R}^n : x \cdot y > f(y) \\ 0 & \text{else} \end{cases} \quad \square$$

# Convex analysis: the subdifferential

## The subdifferential

Def: The subdifferential of  $f \in \text{Conv} \mathbb{R}^n$  is  $\partial f(x) = \{s \in \mathbb{R}^n \mid f(y) \geq f(x) + s \cdot (y-x) \forall y \in \mathbb{R}^n\}$

Elements of  $\partial f$  are called subgradients.



Cor:  $\partial f(x) \neq \emptyset$  for all  $x \in \text{relint dom } f$  (see earlier proof!)


Thm:  $\partial f(x)$  is closed convex for all  $x \in \text{dom } f$  (for  $x \notin \text{dom } f$ ,  $\partial f(x) = \emptyset$ )

proof:  $\partial f(x) = \bigcap_{y \in \mathbb{R}^n} \{s \in \mathbb{R}^n \mid s \cdot (y-x) \leq f(y) - f(x)\}$  is intersection of halfspaces.  $\square$

Thm:  $\partial f(x)$  is bounded for all  $x \in \text{int dom } f$  *, necessary?*

proof: There is  $\varepsilon > 0$  s.t.  $f$  is Lipschitz continuous on  $B(x, \varepsilon)$  (recall:  $f \in \text{Conv} \mathbb{R}^n$  !)

Let  $y = x + \frac{\delta s}{\|s\|_2}$  for  $s \in \partial f(x)$ ,  $\delta \in (0, \varepsilon)$ , then



$$f(x) + \underset{\substack{\uparrow \\ \text{Lipschitz constant}}}{L} \delta \geq f(y) \geq \underset{\substack{\uparrow \\ s \in \partial f(x)}}{f(x) + s \cdot \frac{\delta s}{\|s\|_2}} = f(x) + \delta \|s\|_2 \Rightarrow \|s\|_2 \leq L \delta$$

$\partial f$  is: nonempty  
 • compact  
 • convex

# Convex analysis: the subdifferential

## Subdifferential and directional derivatives

Def: Let  $f \in \text{Conv } \mathbb{R}^n$ . The directional derivative in direction  $v \in \mathbb{R}^n$  at  $x \in \text{dom } f$  is defined as

$$\partial_v f(x) = \liminf_{\lambda > 0, \lambda \rightarrow 0} \frac{f(x + \lambda v) - f(x)}{\lambda} = \inf_{\lambda > 0} \frac{f(x + \lambda v) - f(x)}{\lambda}$$

Rem:  $\partial_{(\cdot)} f(x)$  for  $f \in \text{Conv } \mathbb{R}^n$  is convex by definition.

Thm: Let  $f \in \text{Conv } \mathbb{R}^n, f(x) < \infty$ .  $s \in \partial f(x) \Leftrightarrow \partial_v f(x) \geq s \cdot v \quad \forall v \in \mathbb{R}^n$

• In fact,  $\text{cl}(\partial_{(\cdot)} f(x)) : \mathbb{R}^n \rightarrow \mathbb{R}$  is support function of closed convex  $\partial f(x)$ ,  $v \mapsto \sup_{s \in \partial f(x)} s \cdot v$

proof:  $s \in \partial f(x) \Leftrightarrow \frac{f(x + \lambda v) - f(x)}{\lambda} \geq s \cdot v \quad \forall \lambda > 0, v \in \mathbb{R}^n \Leftrightarrow \partial_v f(x) \geq s \cdot v \quad \forall v \in \mathbb{R}^n$

•  $\partial_{(\cdot)} f(x)$  is pos. homogeneous

$\Rightarrow \text{cl}(\partial_{(\cdot)} f(x)) = \mathcal{B}_c$  for  $c = \{s \in \mathbb{R}^n \mid s \cdot v \leq \partial_v f(x) \quad \forall v \in \mathbb{R}^n\} = \partial f(x)$   $\square$

Thm: Let  $f \in \text{Conv } \mathbb{R}^n$ .  $\partial f(x)$  nonempty & bdd  $\Leftrightarrow x \in \text{int dom } f$

proof:  $\partial f(x) \neq \emptyset$  & bdd  $\Leftrightarrow |\mathcal{B}_{\partial f(x)}| < \infty$  everywhere  $\Leftrightarrow \text{cl}(\partial_{(\cdot)} f(x)) < \infty$

$\Leftrightarrow$   $\partial_{(\cdot)} f(x) < \infty$   $\Leftrightarrow x \in \text{int dom } f$   $\square$

convexity of  $\partial_{(\cdot)} f(x)$

# Convex analysis: the subdifferential

## Relation to differential

Thm: Let  $f \in \text{Conv} \mathbb{R}^n$ .  $f$  differentiable in  $x$  with  $\nabla f(x) = s \iff \partial f(x) = \{s\}$

proof: " $\Rightarrow$ ": Let  $\tilde{s} \in \partial f(x)$ ,  $v \in \mathbb{R}^n$ . Subdifferential:  $f(x+v) \geq f(x) + \tilde{s} \cdot v$

$$\begin{aligned} \text{--- Taylor} \quad &: f(x+v) = f(x) + \nabla f(x) \cdot v + o(\|v\|) \\ & \geq (\tilde{s} - \nabla f(x)) \cdot v + o(\|v\|) \quad \forall v \in \mathbb{R}^n \end{aligned}$$

" $\Leftarrow$ ": (cl  $\partial_{(y)} f(x)$ )  $(y) = \partial_{\nabla f(x)}(y) = s \cdot y$

$\Rightarrow \partial_y f(x) = s \cdot y \quad \forall y$  (since for  $g$  convex cl  $g \neq g$  at most on relbdy dom  $g$ )

$$\Rightarrow \nabla f(x) = s \quad \square$$

Thm: Let  $f \in \text{Conv} \mathbb{R}^n$ .  $x^* = \underset{x}{\text{argmin}} f(x) \iff 0 \in \partial f(x^*)$

proof:  $x^* = \underset{x}{\text{argmin}} f(x)$


$$\Leftrightarrow f(x) \geq f(x^*) \quad \forall x \in \mathbb{R}^n$$

$$\Leftrightarrow f(x) \geq f(x^*) + 0 \cdot (x - x^*) \quad \forall x \in \mathbb{R}^n$$

$$\Leftrightarrow 0 \in \partial f(x^*) \quad \square$$

# Convex analysis: the subdifferential

## Examples and properties



Ex:  $f(x) = |x|, x \in \mathbb{R} \Rightarrow \partial f(x) = \begin{cases} \{\text{sign } x\} & , x \neq 0 \\ [-1, 1] & , x = 0 \end{cases}$

$f(x) = \|x\|_2, x \in \mathbb{R}^n \Rightarrow \partial f(x) = \begin{cases} \{x / \|x\|_2\} & , x \neq 0 \\ \mathbb{B}_{\| \cdot \|_2}(0, 1) & , x = 0 \end{cases}$

$f(x) = \|x\|, x \in \mathbb{R}^n \Rightarrow \partial f(0) = \{s \in \mathbb{R}^n \mid \|y\| \geq s \cdot y \ \forall y \in \mathbb{R}^n\} = \{s \in \mathbb{R}^n \mid \max_{\|y\|=1} s \cdot y \leq 1\} = \mathbb{B}_{\| \cdot \|_*}(0, 1)$

$f(x) = \|x\|_1, x \in \mathbb{R}^n \Rightarrow \partial f(0) = \mathbb{B}_{\| \cdot \|_1}(0, 1) = \text{conv}(\pm e_1, \dots, \pm e_n), \quad e_i = (0, \dots, 1, 0, \dots)$

Thm: Let  $f_1, \dots, f_m \in \text{Conv } \mathbb{R}^n, t_1, t_2 > 0$ , then necessary! (cf.  $f_1 = I_{\{y \geq x^2\}}, f_2 = I_{\{y \leq 0\}}$ )

(a)  $\partial(t_1 f_1 + t_2 f_2)(x) = t_1 \partial f_1(x) + t_2 \partial f_2(x) \quad \forall x \in \text{int}(\text{dom } f_1 \cap \text{dom } f_2)$  ("Moreau-Rockafellar")

(b)  $\partial(\max_{i=1, \dots, m} f_i)(x) = \text{conv}(\bigcup_{j \in J_x} \partial f_j(x))$  with  $J_x = \{j \mid f_j(x) = f(x)\}$

proof: (b) without proof (see e.g. Bauschke & Combettes)

(a) " $\supseteq$ ": Let  $s = t_1 s_1 + t_2 s_2$  with  $s_i \in \partial f_i(x), i=1, 2$ .

$$\begin{aligned} t_1 f_1(y) + t_2 f_2(y) &\geq t_1 (f_1(x) + s_1 \cdot (y-x)) + t_2 (f_2(x) + s_2 \cdot (y-x)) \\ &= (t_1 f_1 + t_2 f_2)(x) + s \cdot (y-x) \end{aligned}$$


# Convex analysis: the subdifferential

## Examples and properties cont'd

"c"  $\therefore$  Let  $s \in \partial(t_1 f_1 + t_2 f_2)(x)$ , set  $g_1 = t_1 f_1 - t_1 f_1(x) - s \cdot (\cdot - x)$ ,  $g_2 = t_2 f_2 - t_2 f_2(x) - s \cdot (\cdot - x)$   
 $\Rightarrow 0 \in \partial(g_1 + g_2)(x)$ ,  $0 = g_1(x) = g_2(x)$ ,  $\begin{cases} \partial g_2(x) = \partial t_2 f_2(x) = t_2 \partial f_2(x) \\ \partial g_1(x) = \partial t_1 f_1(x) - s = t_1 \partial f_1(x) - s \end{cases}$   
 $\Rightarrow$  to show:  $0 \in \partial g_1 + \partial g_2$

- $\Lambda_1 = \text{epi } g_1$ ,  $\Lambda_2 = -\text{epi } g_2$  are convex & have nonempty interior (why?)
- $\text{int } \Lambda_1 \cap \Lambda_2 = \emptyset$  (since  $g_1 + g_2 \geq 0$  and thus  $g_1 \geq -g_2$ )

$$\Rightarrow \exists \alpha \neq 0 \begin{pmatrix} q \\ a \end{pmatrix} \in \mathbb{R}^n \times \mathbb{R} : \sup_{(y,r) \in \Lambda_1} q \cdot y + ar \leq \inf_{(y,r) \in \Lambda_2} q \cdot y + ar$$



- due to  $(x, 0) \in \Lambda_1 \cap \Lambda_2$  even  $\max_{(y,r) \in \Lambda_1} q \cdot y + ar = q \cdot x = \min_{(y,r) \in \Lambda_2} q \cdot y + ar$
- $a < 0$ :  $(x, r) \in \Lambda_1 \forall r > 0 \Rightarrow a \leq 0$ , and  $a = 0$  would imply  $\sup_{y \in \text{dom } f_1} q \cdot y \leq \inf_{y \in \text{dom } f_2} q \cdot y$   
 which contradicts  $\text{int}(\text{dom } f_1 \cap \text{dom } f_2) \neq \emptyset$

$$\text{set } p = -\frac{q}{a} \Rightarrow \underbrace{\max_{(y,r) \in \Lambda_1} p \cdot y - r = p \cdot x}_{\text{choose } r = g_1(y) \Rightarrow r \in \partial g_1(x)} = \underbrace{\min_{(y,r) \in \Lambda_2} p \cdot y - r}_{\text{choose } r = -g_2(y) \Rightarrow -p \in \partial g_2(x)}$$

choose  $r = g_1(y) \Rightarrow r \in \partial g_1(x)$

choose  $r = -g_2(y) \Rightarrow -p \in \partial g_2(x)$

□

# Convex analysis: the subdifferential

## Subdifferential and conjugate

Thm: Let  $f \in \text{Conv } \mathbb{R}^n$ .  $s \in \partial f(x) \Leftrightarrow f^*(s) + f(x) = s \cdot x$

proof:  $s \in \partial f(x) \Leftrightarrow s \cdot y - f(y) \leq s \cdot x - f(x) \quad \forall y \in \text{dom } f$

$$\Leftrightarrow f^*(s) \leq s \cdot x - f(x)$$

•  $f^*(s) \geq s \cdot x - f(x)$  by Fenchel's inequality

Cor: Let  $f \in \text{Conv } \mathbb{R}^n$  be closed.  $s \in \partial f(x) \Leftrightarrow x \in \partial f^*(s)$

proof:  $s \in \partial f(x) \Leftrightarrow f^*(s) + f(x) = s \cdot x \quad \Leftrightarrow f^*(s) + f^{**}(x) = s \cdot x \quad \Leftrightarrow x \in \partial f^*(s) \quad \square$

Cor: Let  $f \in \text{Conv } \mathbb{R}^n$  be closed, then

$$x^* = \underset{x \in \mathbb{R}^n}{\text{argmin}} f(x) \quad \Leftrightarrow 0 \in \partial f(x^*) \quad \Leftrightarrow x^* \in \partial f^*(0)$$

In particular, if  $f^*$  is differentiable in 0, the minimiser is unique.

# Convex analysis: (strong) duality

## Slater's constraint qualification

Thm: (Slater) Consider  $\min_{x \in \mathbb{R}^n} f_0(x)$  s.t.  $f_1(x), \dots, f_m(x) \leq 0, h_1(x) = \dots = h_p(x) = 0$  (P)

for  $f_0, \dots, f_m$  convex,  $h_1, \dots, h_p$  affine (i.e.  $h_1(x) = \dots = h_p(x) = 0 \Leftrightarrow Ax = b$ ),  $\mathcal{D} = \bigcap_{i=1}^m \text{dom } f_i$ .

If (P) is strictly feasible, i.e.  $\exists \tilde{x} \in \text{relint } \mathcal{D}$  with  $f_i(\tilde{x}) < 0, i=1, \dots, m, A\tilde{x} = b$ ,

then strong duality holds, i.e.  $p^* = d^*$ . for proof without see later!

proof: Simplifying assumptions:  $\text{relint } \mathcal{D} = \text{int } \mathcal{D}$ ,  $\text{rank } A = p$

(both can be achieved by transformation to equivalent problems:

cancel linearly dependent rows of  $Ax = b$  & restrict to aff  $\mathcal{D}$ .)

Note: dual problem might change by this!

•  $\tilde{x}$  is feasible  $\Rightarrow p^* < \infty$ . wlog  $p^* > -\infty$  (else  $d^* = -\infty$  by weak duality)

• define  $M = \{(u, v, b) \in \mathbb{R}^{m+p+n} \mid \exists x \in \mathcal{D} : f_i(x) \leq u_i, i=1, \dots, m, h_i(x) = v_i, i=1, \dots, p, f_0(x) \leq b\}$

$N = \{(0, 0, s) \in \mathbb{R}^{m+p+n} \mid s < p^*\}$

$\Rightarrow M, N$  convex &  $M \cap N = \emptyset$  (otherwise there were some feasible  $x$  with  $f_0(x) < p^*$ )

## Slater's constraint qualification cont'd

- there exists a separating hyperplane, i.e.  $(\mu, \lambda, \nu) \neq 0, \alpha \in \mathbb{R}$  s.t.
  - (a)  $\mu \cdot u + \lambda \cdot v + \nu t \geq \alpha \quad \forall (u, v, t) \in M$
  - (b)  $\mu \cdot u + \lambda \cdot v + \nu t \leq \alpha \quad \forall (u, v, t) \in N$
- (a)  $\Rightarrow \mu \geq 0$  (comp.-wise),  $\nu \geq 0$  (else  $\mu \cdot u + \nu t$  would be unbounded below in  $M$ )
- (b)  $\Rightarrow \nu p^* \leq \alpha$
- together,  $\sum_{i=1}^m \mu_i f_i(x) + \sum_{i=1}^p \lambda_i h_i(x) + \nu f_0(x) \geq \alpha \geq \nu p^* \quad \forall x \in \mathcal{D}$ 
  - case  $\nu > 0$ :  $\Rightarrow L(x, \mu/\nu, \lambda/\nu) \geq p^* \Rightarrow g(\mu/\nu, \lambda/\nu) \geq p^* \Rightarrow d^* \geq p^*$
  - case  $\nu = 0$ :  $\Rightarrow \sum_{i=1}^m \mu_i f_i(\bar{x}) \geq 0 \Rightarrow \mu_i \geq 0, f_i(\bar{x}) < 0 \quad \mu = 0$   
 $\Rightarrow \sum_{i=1}^p \lambda_i h_i(x) \geq 0 \quad \forall x \in \mathcal{D}$   
 however,  $\sum_{i=1}^p \lambda_i h_i(\bar{x}) = 0$  together with  $\text{rank } A = p$   
 implies existence of  $x \in \mathcal{D}$  with  $\sum_{i=1}^p \lambda_i h_i(x) < 0 \quad \nabla$   
 $\Rightarrow \nu = 0$  impossible □

Cor:  $(\mu^*, \lambda^*) = (\mu/\nu, \lambda/\nu)$  are dual optimal!


# Convex analysis: (strong) duality

## Slater's constraint qualification - geometric intuition

$$\mathcal{G} = \{(u, v, t) \in \mathbb{R}^{m+p+n} \mid \exists x \in \mathcal{X}, f_0(x) = u, h_i(x) = v_i, f_0(x) = t\}$$


recall:  $g(\mu, \lambda) = \inf_x f_0(x) + \sum_i \mu_i f_i(x) + \sum_j \lambda_j h_j(x) = \inf_{(u, v, t) \in \mathcal{G}} t + \mu^T u + \lambda^T v; \mu \geq 0$

Ex:  $\min_x f_0(x)$  s.t.  $f_n(x) \leq 0$




(let  $\ell_c(u, v) = -\mu^T u - \lambda^T v + c$   
 $\Rightarrow \inf \{ \ell_c(0) \mid \exists (u, v) : (u, v, \ell_c(u, v)) \in \mathcal{G} \}$   
 $= \inf \{ t + \mu^T u + \lambda^T v \mid (u, v, t) \in \mathcal{G} \}$ )

Slater's condition holds



Slater's condition holds



nonconvex problem

no strictly feasible point

MQ N cannot be separated by hyperplane with finite slope

# Convex analysis: (strong) duality

## Implications of constraint qualification

Thm: Strong duality even holds if there is a feasible  $\bar{x} \in \text{relint} \mathcal{X}$  with  $f_i(\bar{x}) < 0$  only for all nonaffine  $f_i$ .

proof: see Rockafellar: Convex Analysis, Thm 28.2 □

In particular, we always have strong duality if all constraints are affine and  $f_0$  is convex, thus for all LPs and QPs!

Thm: Under a constraint qualification (i.e. a condition implying strong duality such as Slater's), if the dual problem is feasible, the KKT conditions hold for a convex optimisation problem (P) at an optimal point  $x$ , i.e.  $\exists \mu, \lambda$  s.t.

$$f_1(x), \dots, f_m(x) \leq 0, \quad h_1(x), \dots, h_p(x) = 0, \quad \mu \geq 0, \quad \mu^\top \begin{pmatrix} f_1(x) \\ \vdots \\ f_m(x) \end{pmatrix} = 0, \\ 0 \in \partial \left[ f_0 + \sum_{i=1}^m \mu_i f_i + \sum_{i=1}^p \lambda_i h_i \right] (x)$$

proof: strong duality  $\Rightarrow$  saddle point property of  $\mathcal{L} \Rightarrow x$  minimizes  $\mathcal{L}(\cdot, \mu, \lambda)$  □

take  $\mu, \lambda$  dual optimal (possible, already shown)

# Convex analysis: (strong) duality

## Fenchel-Rockafellar-duality

Thm: (Fenchel-Rockafellar) Let  $f, g \in \text{Conv } \mathbb{R}^n$ . If either

(a)  $\text{relint dom } f \cap \text{relint dom } g \neq \emptyset$  or

(b)  $\text{relint dom } f^* \cap -\text{relint dom } g^* \neq \emptyset$  and  $f, g$  closed,

then  $\inf_{x \in \mathbb{R}^n} f(x) + g(x) = \sup_{y \in \mathbb{R}^n} -g^*(-y) - f^*(y)$ .

Under (a) the supremum is attained, under (b) the infimum.

proof:  $\forall x, y \in \mathbb{R}^n : f(x) + f^*(y) \geq x \cdot y \geq -g(x) - g^*(-y)$  (Fenchel ineq.)

$$\Rightarrow \inf_x (f(x) + g(x)) \geq \sup_y -g^*(-y) - f^*(y)$$

• if  $\inf = -\infty$ , also  $\sup = -\infty$ , thus assume wlog  $-\infty < \alpha = \inf_x f(x) + g(x)$

• Let (a) hold; it suffices to show existence of some  $y \in \mathbb{R}^n$  with  $-g^*(-y) - f^*(y) \geq \alpha$

• Set  $C = \{(x, v) \in \mathbb{R}^n \times \mathbb{R} \mid v \geq f(x)\}$



$$D = \{(x, v) \in \mathbb{R}^n \times \mathbb{R} \mid v \leq -g(x) + \alpha\}$$

$\Rightarrow C, D$  convex &  $\text{relint } C = \{(x, v) \mid x \in \text{relint dom } f, f(x) < v\}$  is disjoint from  $D$

# Convex analysis: (strong) duality

## Fenchel-Rockafellar-duality cont'd

- $\exists$  separating hyperplane, i.e.  $(s, t) \in \mathbb{R}^n \times \mathbb{R}$  with  $\sup_{(x, y) \in D} s \cdot x + t y \leq \inf_{(x, y) \in C} s \cdot x + t y$ .
- Note:  $t \neq 0$  (otherwise the projections of  $D$  &  $C$ ,  $\text{dom } f$  &  $\text{dom } g$ , would be separated)
- $\Rightarrow$  separating hyperplane is graph of an affine function  $h(x) = \tilde{s} \cdot x + \beta$
- $\Rightarrow f(x) \geq \tilde{s} \cdot x + \beta \geq -g(x) + \alpha \quad \forall x \in \mathbb{R}^n$
- left inequality implies  $-\beta \geq \sup_x \tilde{s} \cdot x - f(x) = f^*(\tilde{s})$
- right ineq. implies  $\alpha - \beta \leq \inf_x \tilde{s} \cdot x + g(x) = -g^*(-\tilde{s})$
- for (er), all follows from duality, noting  $f = f^{**}$ ,  $g = g^{**}$ . □



## Fenchel-Rockafellar-duality: extensions

The result can be sharpened & generalized in different ways, e.g.

- if  $g$  is piecewise affine, we may replace  $\text{reclint dom } g^{(*)}$  by  $\text{dom } g^{(*)}$   
(Rockafellar, Thm. 31.1)

- if  $A: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a linear operator,  $\inf_{x \in \mathbb{R}^n} f(x) + g(Ax) = \sup_{y \in \mathbb{R}^m} -g^*(-y) - f^*(A^T y)$  if  
(a)  $\exists x \in \text{reclint dom } f$  with  $Ax \in \text{reclint dom } g$  or

(b)  $\exists y \in \text{reclint dom } g^*$  with  $-A^T y \in \text{reclint dom } f^*$

(Rockafellar, Cor. 31.2.1)

- under the same conditions,  $\inf_{x \in \mathbb{R}^n} (f(x) + g(Ax)) = \inf_{x \in \mathbb{R}^n} \sup_{y \in \mathbb{R}^m} x^T A^T y + f(x) - g^*(y)$   
and  $\inf$  &  $\sup$  may be swapped.

proof:  $\inf_{x \in \mathbb{R}^n} (f(x) + g(Ax)) \geq \inf_{x \in \mathbb{R}^n} (f(x) + g^{*^*}(Ax)) = \inf_{x \in \mathbb{R}^n} \sup_{y \in \mathbb{R}^m} x^T A^T y + f(x) - g^*(y)$

$$\geq \sup_{y \in \mathbb{R}^m} \inf_{x \in \mathbb{R}^n} x^T A^T y + f(x) - g^*(y) = \sup_{y \in \mathbb{R}^m} -f^*(-A^T y) - g^*(y) = \sup_{y \in \mathbb{R}^m} -f^*(A^T y) - g^*(-y)$$

$\Rightarrow$  strong duality implies equality □

# Convex analysis: (strong) duality

## Fenchel-Rockafellar-duality & Lagrange-duality

Rem: For convex  $f_0, c_I, c_E$ , Lagrange duality is a special case of Fenchel-Rockafellar duality:

choose  $f((x, u, v)) = f_0(x) + \mathbb{I}_{\{c_I(x) \leq u, c_E(x) = v\}}((x, u, v)), h((x, u, v)) = \mathbb{I}_{\{x \in \mathbb{R}^n, u \geq 0, v \geq 0\}}((x, u, v))$

$\Rightarrow f^*((y, \mu, \lambda)) = \sup_{x, u, v, u \geq c_I(x), v = c_E(x)} y \cdot x + \mu \cdot u + \lambda \cdot v - f_0(x), h^*((y, \mu, \lambda)) = \mathbb{I}_{\{0 \leq x \leq \infty\}}^m(y, \mu)$

Now Slater  $\Rightarrow \exists x \in \text{relint } \mathcal{D}$  with  $f((x, -\varepsilon, 0)) < \infty$  for  $\varepsilon > 0$  small enough (comp.-wise)

$\Rightarrow \underbrace{\text{relint dom } h}_{\mathbb{R}^n \times (-\infty, 0]^m \times \{0\}} \cap \text{relint dom } f \neq \emptyset$


Fenchel-Rockafellar

$\Rightarrow p^* = \inf_{x, c_I(x) \leq 0, c_E(x) = 0} f_0(x) = \inf_{(x, u, v)} f((x, u, v)) + h((x, u, v)) \stackrel{\downarrow}{=} \max_{(y, \tilde{\mu}, \tilde{\lambda})} -f^*((y, \tilde{\mu}, \tilde{\lambda})) - h^*((y, \tilde{\mu}, \tilde{\lambda}))$

$= \max_{\tilde{\mu} \leq 0, \tilde{\lambda}} \inf_{\substack{x, u, v \\ u \geq c_I(x), v = c_E(x)}} -\tilde{\mu} \cdot u - \tilde{\lambda} \cdot v + f_0(x) = \max_{\tilde{\mu}, \tilde{\lambda}} \left\{ \begin{array}{l} \inf_x -\tilde{\mu} \cdot c_I(x) - \tilde{\lambda} \cdot c_E(x) + f_0(x) \text{ if } \tilde{\mu} \leq 0 \\ \infty \text{ else} \end{array} \right\} = \max_{\mu \geq 0, \lambda} g(\mu, \lambda)$

$= d^*$

This also implies existence of a dual optimal  $(\mu, \lambda)$ . □



# Optimization algorithms: Simplex method (linear optimization)

## Linear program basics

how to obtain standard form? hint:  $\cdot x = x_0 - x_n$   
 $\cdot$  new variables for ineq. constraints

Def: - A linear program (LP) in standard form is given by

$$\min_{x \in \mathbb{R}^n} cx \quad \text{s.t.} \quad Ax = b, \quad x \geq 0 \text{ coordinate-wise,} \quad (\text{LP})$$

where  $c \in \mathbb{R}^{1 \times n}$ ,  $A \in \mathbb{R}^{p \times n}$  has full rank,  $p < n$ .

It can be viewed as minimisation of  $cx$  over the convex polyhedron  $K = \{x \geq 0 \mid Ax = b\}$ .

• Let  $B \in \mathbb{R}^{p \times p}$  be a matrix composed of  $m$  linearly independent columns  $a_i$ ,  $i \in J \subset \{1, \dots, n\}$  of  $A$ . A point  $x \in \mathbb{R}^n$  with  $x_i = 0 \forall i \notin J$  and  $Ax = b$  is called a basic point wrt. the basis  $B$ .

• A feasible point  $x$  satisfies  $Ax = b, x \geq 0$ ; a basic feasible point is basic & feasible.


We shall assume that any basic feasible point has  $x_i \neq 0 \forall i \in J$ .

Thm: If (LP) admits optimal points, then at least one of them is an extreme point of  $K$ .

proof: Let  $\text{ext } K = \{x_1, \dots, x_k\}$ , then  $\forall x \in K \exists \alpha_1, \dots, \alpha_k \geq 0: \alpha_1 + \dots + \alpha_k = 1, x = \alpha_1 x_1 + \dots + \alpha_k x_k$ .

Let  $z_0 = \min_{i=1, \dots, k} cx_i$ , then  $cx = \alpha_1 (cx_1) + \dots + \alpha_k (cx_k) \geq \alpha_1 z_0 + \dots + \alpha_k z_0 = z_0 \Rightarrow z_0$  is min!  $\square$


note: This does not depend on normal form!




# Optimization algorithms: Simplex method (linear optimization)

## Examples


Ex:  $\min_{x \in \mathbb{R}^2} \underbrace{(1 \ 2)}_c x$   
 s.t.  $\underbrace{(1 \ 1)}_A x = \underbrace{2}_b, x \geq 0$



$\min_{x \in \mathbb{R}^3} (1 \ 2 \ 3)x$   
 s.t.  $(1 \ 1 \ 1)x = 3, x \geq 0$



$\min_{x \in \mathbb{R}^3} (1 \ 2 \ 3)x$   
 s.t.  $\begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & 1 \end{pmatrix} x = \begin{pmatrix} 3 \\ 4 \end{pmatrix}, x \geq 0$



note: extreme points of  $K$  have  $n$ -rank  $A$  zeros!  
 $\Rightarrow$  basic points?  
 optimum?

# Optimization algorithms: Simplex method (linear optimization)

## Basic solutions & extreme points

Then:  $x \in \text{ext}K \Leftrightarrow x$  is a basic feasible point

proof: " $\Rightarrow$ " • wlog assume the nonzero entries of  $x$  to be  $x_1, \dots, x_k$

$$\Rightarrow b = x_1 a_1 + \dots + x_k a_k \quad \text{with } x_i > 0$$

columns of  $A$

• assume  $a_1, \dots, a_k$  are linearly dependent, i.e.  $0 = y_1 a_1 + \dots + y_k a_k$  for some  $y_i$

• letting  $y = (y_1, \dots, y_k, 0, \dots, 0)$ , there exists  $\varepsilon > 0$  with  $x + \varepsilon y \geq 0, x - \varepsilon y \geq 0$

$$\Rightarrow x = \frac{1}{2} \underbrace{(x + \varepsilon y)}_{\in K} + \frac{1}{2} \underbrace{(x - \varepsilon y)}_{\in K}, \text{ which contradicts } x \in \text{ext}K$$

" $\Leftarrow$ " • wlog assume  $x = (x_1, \dots, x_p, 0, \dots, 0)$  (i.e.  $a_1, \dots, a_p$  form a basis)

• assume there are  $y, z \in K, \alpha \in (0, 1)$  with  $x = \alpha y + (1 - \alpha)z$

$$\Rightarrow y_{p+1} = \dots = y_n = 0 \quad \& \quad z_{p+1} = \dots = z_n = 0 \quad (\text{due to } y, z \geq 0)$$

$$\Rightarrow y_1 a_1 + \dots + y_p a_p = b \quad \& \quad z_1 a_1 + \dots + z_p a_p = b$$

Since  $a_1, \dots, a_p$  are linearly independent, this implies  $y = z$

$$\Rightarrow x \in \text{ext}K$$

□

# Optimization algorithms: Simplex method (linear optimization)

## Simplex method: the idea

$$\begin{aligned} \# \text{ ext } K = \# \text{ basic feasible points} &\leq \# \text{ possibilities to choose } p \text{ lin. indep. columns from } A \\ &\leq \binom{n}{p} = \frac{n!}{p!(n-p)!} \end{aligned}$$

*↑ number of...*

Hence, to find an optimal point one only has to test at most  $\binom{n}{p}$  extreme points.

The simplex algorithm does better by generating a sequence  $x^k$  of extreme points, where the function value improves in each step.

### Algorithm (Simplex method):

0) (Initialisation) pick a basis / an extreme point

- choose columns  $a_{i_1}, \dots, a_{i_p}$ ,  $i_j \in J_1$  of  $A$  as a basis
- set  $x_i^0 = 0$  for  $i \notin J^0$  and solve  $Ax^0 = b \Leftrightarrow a_{i_1} x_{i_1}^0 + \dots + a_{i_p} x_{i_p}^0 = b$  for  $x_{i_1}^0, \dots, x_{i_p}^0$
- if the basic point is infeasible, choose a different basis
- note: we may transform  $Ax = b$  via Gaussian elimination s.t.  $a_{i_j} = e_j$
- note: superscript = iteration ; subscript = column / row index

# Optimization algorithms: Simplex method (linear optimization)

## Simplex method: the algorithm

iterate:

### 1) choose function-decreasing direction

- let  $x^{k+1}$  be the next iterate to be found
- $x_{i_1}^{k+1}, \dots, x_{i_p}^{k+1}$  can be solved for in terms of  $x_e^{k+1}$ ,  $e \notin J^k$ :  $x_{i_j}^{k+1} = \alpha_{j0} + \sum_{e \in J^k} \alpha_{je} x_e^{k+1}$  easy if  $\alpha_{ij} = e_j$
- $Cx^{k+1} = z_0 + \sum_{e \in J^k} (c_e + z_e) x_e^{k+1}$  for  $z_e = \sum_{j=1}^p c_{i_j} \alpha_{je}$
- pick  $e \in J^k$  with  $c_e + z_e < 0$ , then increasing  $x_e$  from 0 will decrease  $Cx$   
↖ if there is no such  $e$ ,  $x^k$  is optimal

### 2) move to an adjacent basis / extreme point

- want to add  $a_e$  to basis; which of  $a_{i_1}, \dots, a_{i_p}$  should it replace?
- $a_e = \gamma_1 a_{i_1} + \dots + \gamma_p a_{i_p} \Rightarrow (x_{i_1}^k - \alpha \gamma_1) a_{i_1} + \dots + (x_{i_p}^k - \alpha \gamma_p) a_{i_p} + \alpha a_e = b \quad \forall \alpha \in \mathbb{R}$
- choose  $x_{i_j}^{k+1} = x_{i_j}^k - \alpha \gamma_j$ ,  $j=1, \dots, p$ ,  $x_e^{k+1} = \alpha$  for some  $\alpha$

if  $\gamma_1, \dots, \gamma_p \leq 0$ :  $x^{k+1}$  cannot become a basic feasible point &  $p^* = -\infty$

else: let  $q = \operatorname{argmin}_{S=\{1, \dots, p\}, \gamma_s > 0} \frac{x_{i_s}^k}{\gamma_s}$ ; set  $\alpha = \frac{x_{i_q}^k}{\gamma_q}$ ,  $i_q = l$ ,  $J^{k+1} = \{i_1, \dots, i_p\}$

may again transform  $Ax = b$  s.t.  $a_{i_j} = e_j$ ! the only choice to make  $x^{k+1}$  basic feasible

# Optimization algorithms: Simplex method (linear optimization)

## Simplex method examples

Ex:  $\min_{x \in \mathbb{R}^3} (1 \ 2 \ 3)x \quad \text{s.t.} \quad (1 \ 1 \ 1)x = 3, \quad x \geq 0$

0)  $f^0 = \{3\} \Rightarrow x^0 = \begin{pmatrix} 0 \\ 0 \\ 3 \end{pmatrix} \geq 0$

1)  $(1 \ 1 \ 1)x = 3 \Rightarrow x_3^1 = 3 - x_1^1 - x_2^1 \Rightarrow (1 \ 2 \ 3)x^1 = x_1^1 + 2x_2^1 + 3(3 - x_1^1 - x_2^1)$   
 $= \underset{c_1 - 2a_1}{g^1} - 2x_1^1 - \underset{c_2 - 2a_2}{1}x_2^1 \Rightarrow \text{pick } \ell = 2!$

2) trivial:  $f^1 = \{2\} \Rightarrow x^1 = \begin{pmatrix} 0 \\ 3 \\ 0 \end{pmatrix}$

1)  $x_2^2 = 3 - x_1^2 - x_3^2 \Rightarrow cx^2 = 6 - x_1^2 + x_3^2 \Rightarrow \text{pick } \ell = 1!$

2) trivial:  $f^2 = \{1\} \Rightarrow x^2 = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}$

1)  $x_1^3 = 3 - x_2^3 - x_3^3 \Rightarrow cx^3 = 3 + x_2^3 + 2x_3^3 \Rightarrow x^3 \text{ is optimal!}$

Ex:  $\min_{x \in \mathbb{R}^3} (1 \ 2 \ 3)x \quad \text{s.t.} \quad \begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & 1 \end{pmatrix} x = \begin{pmatrix} 3 \\ 4 \end{pmatrix}, \quad x \geq 0$

0)  $f^0 = \{1, 3\} \Rightarrow x^0 = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$

1)  $Ax = b \Rightarrow \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x_1^1 \\ x_2^1 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \end{pmatrix} x_3^1 \Rightarrow \begin{pmatrix} x_1^1 \\ x_2^1 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 - x_2^1 \end{pmatrix}$

$\Rightarrow cx = 1 + 2x_2^1 + 3(2 - x_2^1) \Rightarrow \text{pick } \ell = 2!$

2)  $a_2 = 0 \cdot a_1 + 1 \cdot a_3 \Rightarrow (x_1^0 - \alpha \cdot 0)a_1 + (x_3^0 - \alpha \cdot 1)a_3 + \alpha a_2 = b$   
 $\Rightarrow q = 3, \alpha = x_3^0 = 2 \Rightarrow f = \{1, 2\}, x^2 = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$

1)  $\Rightarrow x^2 \text{ is optimal!}$

## Structure of Linesearch methods

In the following we shall assume  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  to be twice differentiable and bounded from below.

Def: A descent direction for  $f$  at  $x$  is a  $p \in \mathbb{R}^n$  with  $\nabla f(x)p < 0$   
(direction in which  $f$  decreases)

Alg. (Linesearch method):

given:  $f: \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $x_0 \in \mathbb{R}^n$

repeat 1) choose a descent direction  $p_k \in \mathbb{R}^n$

2) choose a step length  $\alpha_k > 0$

3)  $x_{k+1} = x_k + \alpha_k p_k$

4)  $k \leftarrow k + 1$

until  $x_{k+1}$  sufficiently minimises  $f$



## Stepsize control: Armijo condition

Def:  $\alpha_k > 0$  is said to satisfy Armijo's condition for  $0 < c_1 < 1$  if

$$f(x_k + \alpha_k p_k) \leq f(x_k) + c_1 \alpha_k Df(x_k) p_k \quad \text{expected descent!}$$

Cor: If  $f \in C^1(\mathbb{R}^n)$ ,  $0 < c_1 < 1$ , there exists a step length  $\alpha$  satisfying Armijo's condition

proof: Taylor's thm  $\Rightarrow f(x + \alpha p) = f(x) + \alpha Df(x)p + o(\alpha) < f(x) + c_1 \alpha Df(x)p$  for  $\alpha$  small enough  $\square$



Alg. (backtracking to find good step length):

$\alpha = 1$

if Armijo condition fulfilled

repeat  $\alpha \leftarrow 2\alpha$  until Armijo condition violated

while Armijo condition violated

$\alpha \leftarrow \alpha/2$


## Stepsize control: Wolfe conditions

Def:  $\alpha_k$  is said to satisfy the strong Wolfe conditions for  $0 < c_1 < c_2 < 1$  if it satisfies Armijo's cond. &

$$|Df(x_k + \alpha_k p_k) p_k| \leq -c_2 Df(x_k) p_k$$

Thm: If  $f \in C^1(\mathbb{R}^n)$  bounded from below,  $0 < c_1 < c_2 < 1$ , there exists  $\alpha_k$  satisfying the strong Wolfe cond.

proof: • Let  $g(\alpha) = f(x_k + \alpha p_k)$



- $\alpha_0 := \max \{ \alpha \in \mathbb{R} \mid f(x_k + \alpha p_k) \leq f(x_k) + c_1 \alpha Df(x_k) p_k \}$

- if  $g'(\alpha_0) \geq 0$  choose  $\alpha_k = \max \{ \alpha < \alpha_0 \mid g'(\alpha) = 0 \}$

*note: automatically satisfies Armijo's condition*

- else  $|Df(x_k + \alpha_0 p_k) p_k| = -g'(\alpha_0) < -\underbrace{c_1 g'(\alpha_0)}_{\text{slope of red line}} < -c_2 g'(\alpha_0) = -c_2 Df(x_k) p_k$

□

## Global convergence

Thm (Zoutendijk): If  $\exists L > 0: \|\nabla f(x) - \nabla f(y)\|_2 \leq L\|x - y\|_2$  (gradient is Lipschitz)

•  $\cos \theta_k = \frac{-\nabla f(x_k) p_k}{\|\nabla f(x_k)\|_2 \|p_k\|_2}$  (angle between  $-\nabla f(x_k)$  and search direction)

•  $\alpha_k$  satisfies strong Wolfe conditions,

then  $\sum_{k=1}^{\infty} \cos^2 \theta_k \|\nabla f(x_k)\|_2^2 < \infty$  (either  $\nabla f \rightarrow 0$  or angle degenerates)

the subsequent convergence rate proof is essentially start from here

proof: 1) use Wolfe condition & fact that  $\nabla f$  cannot change too fast to obtain lower bound on step length

$$\alpha_k L \|p_k\|_2^2 \geq (\nabla f(x_k + \alpha_k p_k) - \nabla f(x_k)) p_k \geq (c_2 - 1) \nabla f(x_k) p_k$$

$$\Rightarrow \alpha_k \geq \frac{c_2 - 1}{L \|p_k\|_2^2} \nabla f(x_k) p_k$$

2) use Armijo condition to estimate worst case function decrease

$$\Rightarrow f(x_{k+1}) \leq f(x_k) - c_1 \frac{1 - c_2}{L} \frac{(\nabla f(x_k) p_k)^2}{\|p_k\|_2^2} = f(x_k) - c_1 \frac{1 - c_2}{L} \cos^2 \theta_k \|\nabla f(x_k)\|_2^2$$

$$\Rightarrow -\infty < \inf f - f(x_0) \leq f(x_2) - f(x_0) = \sum_{k=0}^{2-1} f(x_{k+1}) - f(x_k) \leq -c_1 \frac{1 - c_2}{L} \sum_{k=0}^{2-1} \cos^2 \theta_k \|\nabla f(x_k)\|_2^2 \quad \square$$

Rem: If  $\nabla f$  is only Hölder with exponent  $\beta$ , one gets  $\sum_{k=1}^{\infty} |\cos \theta_k| \|\nabla f(x_k)\|_2^{1 + \frac{1}{\beta}}$  (homework)

$\Rightarrow \nabla f \rightarrow 0$  at a slower rate!



## Optimality bounds from gradient

Thm: If there exist  $m, M > 0$  s.t.  $m I \leq D^2 f(x) \leq M I \quad \forall x \in U(x^*)$ , then  $\forall x \in U(x^*)$

*in sense of quadratic forms, i.e. eigenvalues lie in  $[m, M]$*

- $\frac{1}{2M} \|Df(x)\|_2^2 \leq f(x) - p^* \leq \frac{1}{2m} \|Df(x)\|_2^2$
- $\frac{1}{M} \|Df(x)\|_2 \leq \|x - x^*\|_2 \leq \frac{1}{m} \|Df(x)\|_2$

↑  
neighbourhood of opt. point  $x^*$



proof: • By Taylor,  $f(y) = f(x) + Df(x)(y-x) + \frac{1}{2}(y-x)^T D^2 f(z)(y-x)$  for some  $z$  between  $x, y$

$$\Rightarrow f(y) \begin{cases} \leq \\ \geq \end{cases} f(x) + Df(x)(y-x) + \frac{M}{2} \|y-x\|_2^2$$

$\forall x, y \in U$  (if  $U$  convex)

• minimise both sides over  $y$ : on lhs,  $y = x^*$

on rhs,  $y = \tilde{y} = x - \frac{1}{m} \nabla f(x)$

$$\Rightarrow p^* = f(x^*) \leq f(x) - \frac{1}{2} \frac{1}{m} \|Df(x)\|_2^2$$

• As above,  $f(x) \geq f(x^*) + Df(x^*)(x-x^*) + \frac{M}{2} \|x-x^*\|_2^2 = p^* + \frac{M}{2} \|x-x^*\|_2^2$

$$\Rightarrow \|x-x^*\|_2^2 \leq \frac{2}{m} (f(x) - p^*) \leq \frac{1}{m^2} \|Df(x)\|_2^2 \quad \square$$

# Optimization algorithms: Linesearch methods for unconstrained optimization

## Gradient descent, steepest descent, Newton's method

Def: The gradient descent direction is  $p_k^g = -\nabla f(x_k)$

Let  $\|\cdot\|$  be a norm with dual norm  $\|\cdot\|_*$ ; the steepest descent direction is

$$p_k^s = \|\nabla f(x_k)\|_* \underset{\|v\|=1}{\operatorname{argmin}} Df(x_k)v \quad (\text{gradient descent} = \text{steepest descent with } \|\cdot\|_2)$$

(scaling s. t.  $-Df(x_k)p_k^s = \|\nabla f(x_k)\|_*^2$ )

The Newton step is  $p_k^N = -D^2f(x_k)^{-1} \nabla f(x_k)$

Rem: If  $D^2f$  is positive definite, all above are descent directions. Indeed,

$$Df(x_k)p_k^g = -\|Df(x_k)\|_2^2 < 0, \quad Df(x_k)p_k^s = -\|\nabla f(x_k)\|_*^2 < 0, \quad Df(x_k)p_k^N = -\|\nabla f(x_k)\|_{D^2f(x_k)}^2 < 0.$$

Rem: The Newton step finds minimum in a single step if  $f$  is quadratic and  $\alpha_k = 1$ :

$$f(x) = \frac{\alpha}{2} x^T A x + b^T x \text{ is minimised by } x^* = -A^{-1}b, \text{ and } x_k + \alpha_k p_k^N = x_k - A^{-1}(Ax_k + b) = x^*.$$

Cor: If  $mI \leq D^2f(x) \leq M I \quad \forall x \in \mathbb{R}^n$ , the  $\alpha_k$  satisfy the strong Wolfe conditions, then

$$Df(x_k) \xrightarrow{k \rightarrow \infty} 0 \quad \text{for gradient/steepest/Newton descent.}$$

proof: Follows from Zoutendijk's theorem with  $\cos \theta_k = 1$  for gradient descent,

$$\cos \theta_k \geq \frac{m}{M} \text{ for Newton's method, } \cos \theta_k \geq \frac{c}{C} \text{ for steepest descent with } c\|\cdot\| \leq \|\cdot\|_2 \leq C\|\cdot\|_* \quad \square$$

# Optimization algorithms: Linesearch methods for unconstrained optimization

## Experimental convergence rate

```
function experimentalConvergenceRate
```

```
% f(x) = x^4/4-5x^3/3+3x^2
```

```
% f'(x) = x*(x-2)*(x-3) = x^3-5x^2+6x; global minimum at 0
```

```
f = @(x) x.^4/4-5*x.^3/3+3*x.^2;
```

```
df = @(x) x*(x-2)*(x-3);
```

```
d2f = @(x) 3*x^2-10*x+6;
```

```
fun = @(x) combineFunctions(x,f,df,d2f);
```

```
x = -1:.01:4;
```

```
plot(x,f(x),'Linewidth',5);
```

```
pause;
```

```
x0 = 1.3;
```

```
maxIter = 100;
```

```
[x,iterSteepest] = descendSteepest( fun, x0, maxIter, true );
```

```
[x,iterNewton] = newtonMethod( fun, x0, maxIter, 1e-26, true );
```

```
semilogy(1:length(iterSteepest),abs(iterSteepest),'r','Linewidth',5,...
```

```
1:length(iterNewton),abs(iterNewton),'g','Linewidth',5);
```

```
end
```



```
function [A,B,C] = combineFunctions(x,a,b,c)
```

```
A = a(x);
```

```
B = b(x);
```

```
C = c(x);
```

```
end
```



# Optimization algorithms: Linesearch methods for unconstrained optimization

## Convergence rate gradient/steepest descent

Thm: Let  $S = \{x \in \mathbb{R}^n \mid f(x) \leq f(x_0)\}$ ,  $mI \leq D^2 f(x) \leq MI \quad \forall x \in S$ ,  $\left\{ \begin{array}{l} \frac{1}{\gamma} \| \cdot \|_1 \geq \| \cdot \|_2 \geq \gamma \| \cdot \|_x \\ \frac{\alpha}{\beta} \| \cdot \|_y \geq \| \cdot \|_z \geq \beta \| \cdot \|_1 \end{array} \right\}$ ,

$x_k$  the steepest descent iterates for backtracking with  $c_1 \in (0, \frac{1}{2})$ . Then

$$(f(x_k) - p^*) \leq \delta^k (f(x_0) - p^*) \quad \text{for } \delta = 1 - 2m c_1 \tilde{\gamma}^2 \min \left\{ 1, \frac{\gamma^2}{2M} \right\} < 1.$$

↑ linear convergence
↑ depends on the condition number  $\frac{m}{M}$  of  $D^2 f$ !



proof:  $f(x_k + \alpha \underbrace{p_k}_{\substack{\text{US} \\ p_k}}) \leq f(x_k) + \alpha Df(x_k) p_k + \frac{M}{2} \alpha^2 \|p_k\|_2^2 = f(x_k) + \alpha \left(1 - \frac{M}{2\gamma^2} \alpha\right) Df(x_k) p_k$

$$\leq \|p_k\|_2^2 / \gamma^2 = \|\nabla f(x_k)\|_x^2 / \gamma^2 = -Df(x_k) p_k / \gamma^2$$

$\Rightarrow$  any  $\alpha \in (0, \frac{2\gamma^2}{M}(1-c_1)) > (0, \gamma^2/M)$  satisfies Armijo's condition  $\Rightarrow \alpha_k \geq \min \left\{ 1, \frac{\gamma^2}{2M} \right\}$

$$\begin{aligned} \cdot f(x_{k+1}) &= f(x_k + \alpha_k p_k) \leq f(x_k) + c_1 \alpha_k Df(x_k) p_k \leq f(x_k) - c_1 \min \left\{ 1, \frac{\gamma^2}{2M} \right\} \|\nabla f(x_k)\|_x^2 \\ &\leq f(x_k) - c_1 \tilde{\gamma}^2 \min \left\{ 1, \frac{\gamma^2}{2M} \right\} \underbrace{\|\nabla f(x_k)\|_2^2}_{\geq 2m (f(x_k) - p^*)} \end{aligned}$$

$$\Rightarrow f(x_{k+1}) - p^* \leq f(x_k) - p^* - 2m c_1 \tilde{\gamma}^2 \min \left\{ 1, \frac{\gamma^2}{2M} \right\} (f(x_k) - p^*) = \delta (f(x_k) - p^*) \quad \square$$

- What does this imply about  $\|\nabla f(x_k)\|_2, \|x_k - x^*\|_2$ ?
- What rate follows from Zoutendijk's Thm?

# Optimization algorithms: Linesearch methods for unconstrained optimization

## Acceptance of Newton step

why important?

Thm: If  $D^2f$  continuous,  $mI \leq D^2f(x) \leq MI \quad \forall x$ ,  $c_0 < \frac{1}{2}$ , then  $\|p_k\| \rightarrow 0$ ,  $\lim_{k \rightarrow \infty} \frac{\|p_k^N - p_k\|}{\|p_k\|} = 0$ .

then for  $k$  sufficiently large,  $\alpha_k = 1$  satisfies the strong Wolfe conditions.

proof:  $0 \leq \frac{Df(x_k)p_k}{\|p_k\|_2} + m \leq \frac{Df(x_k)p_k}{\|p_k\|_2} + \frac{p_k^T D^2f(x_k)p_k}{\|p_k\|_2^2} = \frac{p_k^T D^2f(x_k)(p_k - p_k^N)}{\|p_k\|_2^2} \leq M \frac{\|p_k - p_k^N\|_2}{\|p_k\|_2} \rightarrow 0$

$\Rightarrow$  for  $k$  large enough,  $-Df(x_k)p_k \geq \frac{m}{2} \|p_k\|_2^2$

1) Armijo's condition satisfied for  $k$  large enough:

• Taylor's thm:  $f(x_k + p_k) = f(x_k) + Df(x_k)p_k + \frac{1}{2} p_k^T D^2f(x_k + q_k)p_k$  for some  $q_k \in [0, p_k]$

•  $f(x_k + p_k) - f(x_k) - \frac{1}{2} Df(x_k)p_k = \frac{1}{2} [Df(x_k)p_k + p_k^T D^2f(x_k + q_k)p_k]$

$= \frac{1}{2} [\underbrace{(Df(x_k)p_k + p_k^T D^2f(x_k)p_k)}_{=0} + \underbrace{p_k^T D^2f(x_k)(p_k - p_k^N)}_{\leq M o(\|p_k\|_2^2)} + \underbrace{p_k^T (D^2f(x_k + q_k) - D^2f(x_k))p_k}_{= o(\|p_k\|_2^2)}] = o(c_0 - \frac{1}{2}) Df(x_k)p_k$

2) strong Wolfe condition satisfied for  $k$  large enough:

$|Df(x_k + p_k)p_k| = \left| \underbrace{Df(x_k)p_k + p_k^T D^2f(x_k)p_k}_{= p_k^T D^2f(x_k)(p_k - p_k^N)} + \underbrace{\int_0^1 p_k^T (D^2f(x_k + t p_k) - D^2f(x_k))p_k dt}_{= o(\|p_k\|_2^2)} \right|$

$= p_k^T D^2f(x_k)(p_k - p_k^N) \leq M o(\|p_k\|_2^2) = o(\|p_k\|_2^2)$

while  $-c_2 Df(x_k)p_k \geq c_2 \frac{m}{2} \|p_k\|_2^2$

□

## Convergence rate Newton's method

Thm: Let  $D^2f$  Lipschitz with constant  $L$ ,  $mI \leq D^2f(x) \leq MI$ ,  $c_n < \frac{1}{2}$ ,

$p_k = p_k^N$ ,  $\alpha_k = 1$  where possible. Then  $\limsup_{k \rightarrow \infty} \frac{\|x_{k+1} - x^*\|_2}{\|x_k - x^*\|_2^2} \leq \frac{L}{m}$ . quadratic convergence

proof: • Zoutendijk  $\Rightarrow \|Df(x_k)\|_2 \rightarrow 0 \Rightarrow \|x_k - x^*\|_2 \rightarrow 0$  &  $p_k \rightarrow 0$

• By previous theorem,  $\alpha_k = 1$  for  $k$  large enough, i.e.  $x_{k+1} = x_k + p_k$

$$\begin{aligned} \bullet D^2f(x_k)(x_{k+1} - x^*) &= D^2f(x_k)(p_k - (x^* - x_k)) = Df(x^*) - Df(x_k) - D^2f(x_k)(x^* - x_k) \\ &= \int_0^1 (D^2f(x_k + t(x^* - x_k)) - D^2f(x_k))(x^* - x_k) dt \end{aligned}$$

• upon taking norms on both sides,

$$m \|x_{k+1} - x^*\|_2 \leq \|D^2f(x_k)(x_{k+1} - x^*)\|_2 \leq \left\| \int_0^1 (D^2f(x_k + t(x^* - x_k)) - D^2f(x_k))(x^* - x_k) dt \right\|_2 \leq L \|x_k - x^*\|_2^2 \quad \square$$

Rem: • If  $mI \leq D^2f(x^*) \leq MI$ , then by continuity, analogous bounds hold in a neighbourhood.

• If the algorithm at some point reaches such a neighbourhood, we have quadratic convergence

• If  $D^2f$  is only Hölder-continuous with exponent  $\alpha > 0$ ?

• Newton's method is invariant under coordinate transforms  $x \mapsto Ax$  (check!), but

analysis is not  $\Rightarrow$  actually, only  $\| (D^2f(x))^{-1} D^2f(x + t(x^* - x)) - Id \|_2 \leq C \|x^* - x\|_2^\alpha$  required,

i.e. no separate Lipschitz & strong convexity condition (related to "self-concordance")

## Complexity of Newton's method

*implies*  $\|x_k - x^*\|_2 < \min\left(\frac{1}{2} - c_1, c_2\right) \frac{m}{L}$

Thm: Under same conditions as before,  $\|Df(x_k)\|_2 < \min\left(\frac{1}{2} - c_1, \frac{c_2}{2}\right) \frac{m^2}{L}$  implies  $\|x_{k+1} - x^*\|_2 \leq \frac{L}{m} \|x_k - x^*\|_2^2$

proof: same proof holds, only need to show  $\alpha_k = 1$ ; for this repeat previous proof, using

$$p_k^{N^T} (D^2 f(x_k + \alpha p_k) - D^2 f(x_k)) p_k^N \leq L \|p_k^N\|_2^3 \leq L \frac{\|Df(x_k)\|_2}{m} \|p_k^N\|_2^2 \leq \begin{cases} \left(\frac{1}{2} - c_1\right) m \|p_k^N\|_2^2 \leq \left(c_1 - \frac{1}{2}\right) Df(x_k) p_k^N \\ \frac{c_2}{2} m \|p_k^N\|_2^2 \end{cases} \quad \square$$

Thm: Under same conditions as before and assuming  $\alpha_k$  to be chosen maximally up to a factor  $\beta \in (0, 1)$ ,

for all  $\eta > 0$  there is a  $\gamma > 0$  s.t.  $\|Df(x_k)\|_2 \geq \eta$  implies  $f(x_{k+1}) \leq f(x_k) - \gamma$ .

proof:  $-Df(x_k) p_k^N = p_k^{N^T} D^2 f(x_k) p_k^N \geq m \|p_k^N\|_2^2$

$f(x_k + \alpha p_k^N) \leq f(x_k) + \alpha Df(x_k) p_k^N + \frac{M}{2} \alpha^2 \|p_k^N\|_2^2 \leq f(x_k) + \alpha \left(1 - \frac{\alpha M}{2m}\right) Df(x_k) p_k^N$

$\Rightarrow \hat{\alpha} = 2 \frac{m}{M} (1 - c_1)$  satisfies Armijo's condition

case 1 ( $\hat{\alpha}$  satisfies Wolfe conditions):  $\alpha_k \geq \beta \hat{\alpha} \Rightarrow f(x_k + \alpha_k p_k^N) - f(x_k) \leq \beta \hat{\alpha} c_1 Df(x_k) p_k^N \leq -\gamma$

case 2 ( $Df(x_k + \hat{\alpha} p_k^N) p_k^N$  too negative):  $\alpha_k \geq \hat{\alpha} \geq \beta \hat{\alpha}$ ,  $\gamma = \frac{\beta \hat{\alpha} c_1 m^2}{m} \leq -\frac{1}{4} \eta^2$

case 3 ( $Df(x_k + \hat{\alpha} p_k^N) p_k^N$  too positive):  $f(x_k + \alpha_k p_k^N) - f(x_k) \leq f(x_k + \hat{\alpha} p_k^N) - f(x_k) \leq -\gamma$ . □

Cor: Newton's method reaches  $\|x_k - x^*\|_2 < \epsilon$  after  $\frac{f(x_0) - p^*}{\gamma} + \log_2 \left( \frac{\log \epsilon}{\log \min\left(\frac{1}{2} - c_1, \frac{c_2}{2}\right)} \right)$  iterations.

## Modified Newton's method

• If  $D^2f(x_k)$  is indefinite, one can choose  $p_k = -(D^2f(x_k) + M_k)^{-1} \nabla f(x_k)$  for  $M_k$  such that  $D^2f(x_k) + M_k$  is positive definite, e.g.

$$M_k = \max(0, \varepsilon - \lambda_{\min}) I \text{ for } \lambda_{\min} \text{ the smallest eigenvalue of } D^2f(x_k).$$

• In quasi-Newton methods we approximate the Hessian just from the past gradients.

Thm: Same conditions as before, only  $\lim_{k \rightarrow \infty} \frac{\|p_k - p_k^N\|_2}{\|p_k\|_2} = 0$ . Then  $\limsup_{k \rightarrow \infty} \frac{\|x_{k+1} - x^*\|_2}{\|x_k - x^*\|_2} = 0$ .

proof: As before,  $\|x_k - x^*\|_2 \rightarrow 0$  &  $p_k \rightarrow 0$  &  $x_{k+1} = x_k + p_k$

superlinear convergence

$$\underbrace{D^2f(x_k)(x_{k+1} - x^*)}_{\text{norm} \geq m \|x_{k+1} - x^*\|_2} = D^2f(x_k)(x_k - x^* + p_k^N - p_k^N + p_k) = \underbrace{Df(x^*) - Df(x_k)}_{\text{norm} \leq L \|x_k - x^*\|_2^2} - \underbrace{D^2f(x_k)(x^* - x_k)}_{\text{norm} \leq M O(\|p_k\|_2)} + D^2f(x_k)(p_k - p_k^N)$$

$$\Rightarrow m \|x_{k+1} - x^*\|_2 \leq L \|x_k - x^*\|_2^2 + M O(\|p_k\|_2)$$

$$= \|x_{k+1} - x_k\|_2 \leq \|x_{k+1} - x^*\|_2 + \|x_k - x^*\|_2$$

$$\Rightarrow \|x_{k+1} - x^*\|_2 \leq o(\|x_k - x^*\|_2)$$

□

# Optimization algorithms: Linesearch methods for unconstrained optimization

## Quasi-Newton methods

• Abbreviate:  $g_k = \nabla f(x_k)$ ,  $s_k = x_{k+1} - x_k$ ,  $y_k = g_{k+1} - g_k$ ,  $H_k = \text{approximation of } D^2 f(x_k)$

Rem: Taylor  $\Rightarrow \nabla f(x_k) = \nabla f(x_{k+n}) + D^2 f(x_{k+n})(x_k - x_{k+n}) + \text{h.o.t.} \Leftrightarrow D^2 f(x_{k+n})s_k = y_k + \text{h.o.t.}$

Def: The condition  $H_{k+n} s_k = y_k$  for  $H_{k+n} \in \mathbb{R}^{n \times n}$  is called the secant condition.



secant with slope  $H_{k+n}$

"secant method":

$$x_{k+n} = x_k - \alpha_k \left( \frac{f'(x_k) - f'(x_{k-1})}{x_k - x_{k-1}} \right)^{-1} f'(x_k)$$

Rem: If  $H_{k+1}$  is positive definite, the secant condition requires  $s_k^T y_k > 0$ .

Thm: A descent step satisfying the strong Wolfe conditions satisfies  $s_k^T y_k > 0$ .

proof: strong Wolfe  $\Rightarrow g_{k+1}^T s_k \geq c_2 g_k^T s_k \Rightarrow y_k^T s_k = \underbrace{(c_2 - 1)}_{< 0} \alpha_k \underbrace{g_k^T p_k}_{< 0} > 0$   $\square$

Quasi-Newton methods choose  $p_k = H_k^{-1} g_k$ , where  $H_0$  is an initial approximation of  $D^2 f(x_0)$  (typically  $H_0 = I$ ), and  $H_k$  is an improved approximation, which is updated in each step to satisfy the secant condition. Typically, the update is of low rank; different updates lead to different methods.

## Low-rank updates

Thm (Sherman-Morrison-Woodbury formula): Let  $A \in \mathbb{R}^{n \times n}$ ,  $U, V \in \mathbb{R}^{n \times r}$ .

$$B = A + UV^T \Rightarrow B^{-1} = A^{-1} - A^{-1}U(I + V^T A^{-1}U)^{-1}V^T A^{-1}$$

proof: By calculation (homework) □

Ex:  $(I + \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} (123))^{-1} v = \begin{pmatrix} 2 & 2 & 3 \\ 2 & 5 & 6 \\ 3 & 6 & 10 \end{pmatrix}^{-1} v = Iv - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} (1 + (123)\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix})^{-1} (123)v = v - \frac{\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot v}{15} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$

Def: The symmetric rank-1 (SR1) update is given by

$$H_{k+1} = H_k + \frac{(y_k - H_k s_k)(y_k - H_k s_k)^T}{(y_k - H_k s_k)^T s_k} \quad \text{rank-1 update}$$

• The Davidon-Fletcher-Powell (DFP) update is given by

$$H_{k+1} = \left( I - \frac{y_k s_k^T}{y_k^T s_k} \right) H_k \left( I - \frac{s_k y_k^T}{y_k^T s_k} \right) + \frac{y_k y_k^T}{y_k^T s_k} \quad \text{rank-2 update}$$

• The Broyden-Fletcher-Goldfarb-Shanno (BFGS) update is given by

$$H_{k+1} = H_k - \frac{H_k s_k s_k^T H_k}{s_k^T H_k s_k} + \frac{y_k y_k^T}{y_k^T s_k} \quad \text{rank-2 update}$$

Rem: By the S-M-W formula,  $B_{k+1} = H_{k+1}^{-1} = \begin{cases} B_k + \frac{(s_k - B_k y_k)(s_k - B_k y_k)^T}{(s_k - B_k y_k)^T y_k} & \text{(SR1)} \\ B_k - \frac{B_k y_k y_k^T B_k}{y_k^T B_k y_k} + \frac{s_k s_k^T}{y_k^T s_k} & \text{(DFP)} \\ \left( I - \frac{s_k y_k^T}{y_k^T s_k} \right) B_k \left( I - \frac{y_k s_k^T}{y_k^T s_k} \right) + \frac{s_k s_k^T}{y_k^T s_k} & \text{(BFGS)} \end{cases}$

# Optimization algorithms: Linesearch methods for unconstrained optimization

## SR1, Davidon-Fletcher-Powell, Broyden-Fletcher-Goldfarb-Shanno

Rem: • SR1 is the unique symmetric rank-1 update satisfying the secant condition (homework)

- The SR1 update can fail. (zero denominator)
- Even if  $H_k$  is pos. def.,  $H_{k+1}$  need not be. (homework)

• DFP is the rank-2 update satisfying the secant condition and being "closest" to  $H_k$ :

$$\text{DFP solves } H_{k+1} = \underset{H}{\operatorname{argmin}} \|G^{-\frac{1}{2}}(H - H_k)G^{-\frac{1}{2}}\|_F^2 \quad \text{s.t. } H = H^T, Hs_k = \gamma_k, \quad G = \int_0^1 D^2 f(x_k + \tau(x_{k+1} - x_k)) d\tau$$

- Under the strong Wolfe conditions, the DFP update is well-defined, since  $y_k^T s_k > 0$ .
- $H_k$  pos. def.  $\Rightarrow H_{k+1}$  pos. def.

(indeed, for  $z \in \mathbb{R}^n$  let  $w = (I - \frac{s_k y_k^T}{y_k^T s_k})z$ , then  $z^T H_{k+1} z = w^T H_k w + \frac{(y_k^T z)^2}{y_k^T s_k} \geq 0$ ,  
and  $z^T H_{k+1} z = 0 \Rightarrow y_k^T z = 0 \Rightarrow w = z$  &  $z^T H_{k+1} z = z^T H_k z > 0$   $\hookrightarrow$ )

• BFGS has the same properties as DFP, only for  $B_{k+1}$  instead of  $H_{k+1}$ .

# Optimization algorithms: Line search methods for unconstrained optimization

## Convergence of BFGS

*note: estimate of  $\cos \theta_k$  via eigenvalues of Hessian is here replaced by an estimate via  $\text{trace}(H_k)$ ,  $\det(H_k)$*

Thm: If  $f \in C^2(\mathbb{R}^n)$  with  $mI \leq D^2f \leq MI$ , then  $x_k \rightarrow x^*$  for the iterates  $x_k$  of the BFGS quasi-Newton method with  $H_0 = I$  and strong Wolfe stepsize control.

proof: Define  $m_k = \frac{y_k^T s_k}{s_k^T s_k}$ ,  $M_k = \frac{\|y_k\|_2^2}{y_k^T s_k}$ ,  $q_k = \frac{s_k^T H_k s_k}{s_k^T s_k}$ ,  $\cos \theta_k = \frac{s_k^T H_k s_k}{\|s_k\|_2 \|H_k s_k\|_2} = \frac{-g_k^T p_k}{\|g_k\|_2 \|p_k\|_2}$

• note:  $m_k \geq m$ ,  $M_k \leq M$

•  $\text{trace}(H_{k+1}) = \text{trace}(H_k) - \frac{\|H_k s_k\|_2^2}{s_k^T H_k s_k} + \frac{\|y_k\|_2^2}{y_k^T s_k} = \text{trace}(H_k) - \frac{q_k}{\cos^2 \theta_k} + M_k$  (homework)

•  $\det(H_{k+1}) = \det(H_k) \frac{y_k^T s_k}{s_k^T H_k s_k} = \det(H_k) \frac{m_k}{q_k}$  (homework)

• Let  $\Psi(H) = \text{trace} H - \ln(\det H)$ , then  $\Psi > 0$  (homework)

$$\text{and } \Psi(H_{k+1}) = \Psi(H_k) + \underbrace{(M_k - \ln m_k - 1)}_{\leq M - \ln m - 1 = c \geq 0} + \underbrace{\left[1 - \frac{q_k}{(\cos \theta_k)^2} + \ln \frac{q_k}{\cos^2 \theta_k}\right]}_{\leq 0} + \ln \cos^2 \theta_k$$

$$\Rightarrow 0 < \Psi(H_{k+1}) \leq \Psi(H_0) + c(k+1) + \sum_{j=0}^k \ln \cos^2 \theta_j$$

• assume  $\cos \theta_k \rightarrow 0$ , then  $\exists k_1 > 0 \forall k > k_1: \ln \cos^2 \theta_k \leq -2c$  (wlog,  $c > 0$ )

$$\Rightarrow 0 < \Psi(H_0) + c(k+1) + \sum_{j=0}^{k_1} \ln \cos^2 \theta_j - 2c(k - k_1) \xrightarrow{k \rightarrow \infty} -\infty \quad \downarrow$$

Hence,  $\cos \theta_k$  does not converge to 0  $\Rightarrow Df(x_k) \rightarrow 0$  for subsequence  $\Rightarrow x_k \rightarrow x^*$   
*20 kunden disk* *strong convexity*

□

## Linear convergence of BFGS

Thm: Under the same conditions,  $\exists \mu \in (0, 1), \rho > 0: \|x_k - x^*\|_2 \leq \rho \mu^k \|x_0 - x^*\|_2$ .

proof: • already know  $0 < \psi(H_0) + c(k+1) + \sum_{j=0}^k \ln \cos^2 \theta_j$

•  $\forall r \in (0, 1) \exists \kappa > 0: \cos^2 \theta_j \geq \kappa$  for at least  $\lfloor r(k+1) \rfloor$  indices  $j$  in  $\{0, \dots, k\} \forall k$ .

Indeed, let  $\eta_j = -\ln \cos^2 \theta_j$  and let  $\eta^k$  be the  $\lfloor r(k+1) \rfloor$  smallest of  $\{\eta_{01}, \dots, \eta_k\}$ .

Set  $I^k = \{i \in \{0, \dots, k\} \mid \eta_i > \eta^k\}$ . For  $i \in \{0, \dots, k\} \setminus I^k$ ,

$$\eta_i \leq \eta^k \leq \frac{1}{|I^k|} \sum_{j \in I^k} \eta_j \leq \frac{1}{1-r} \frac{1}{k+1} \sum_{j=0}^k \eta_j < \frac{1}{1-r} \left[ \frac{\psi(H_0)}{k+1} + c \right] \leq C$$

$$\Rightarrow \cos \theta_i \geq \exp(-C/2) =: \sqrt{\kappa}$$

• recall from proof of Zoutendijk's theorem:  $f(x_k) - f(x_{k+1}) \geq c_1 \frac{1-c_2}{M} \cos^2 \theta_k \underbrace{\|Df(x_k)\|_2^2}_{\geq 2m(f(x_k) - f(x^*))}$

$$\Rightarrow f(x_{k+1}) - f(x^*) \leq (1 - c_1 \frac{1-c_2}{M} 2m \cos^2 \theta_k) (f(x_k) - f(x^*))$$

$$\Rightarrow f(x_{k+1}) - f(x^*) \leq (1 - c_1 \frac{1-c_2}{M} 2m \kappa)^{\lfloor r(k+1) \rfloor} (f(x_0) - f(x^*))$$

monotonicity of  $f(x_k)$  and  $\cos^2 \theta_k \geq \kappa$  for fraction  $r$  of all indices

• result follows from  $\frac{1}{M} (f(x) - f(x^*)) \leq \|x - x^*\|_2^2 \leq \frac{1}{m} (f(x) - f(x^*))$

$$\text{for } \rho = \left(\frac{M}{m}\right)^{1/2}, \mu = (1 - c_1 \frac{1-c_2}{M} 2m \kappa)^{r/2}$$

□

# Superlinear convergence of BFGS *for simplicity, we rescale domain s.t. $D^2 f(x^*) = I$*

Thm: Under same conditions as before, if  $D^2 f$  is Lipschitz with constant  $L$ ,  $\lim_{k \rightarrow \infty} \frac{\|x_{k+1} - x^*\|_2}{\|x_k - x^*\|_2} = 0$ .

proof: Let  $G = D^2 f(x^*)$ ,  $\tilde{s}_k = G^{-\frac{1}{2}} s_k$ ,  $\tilde{y}_k = G^{-\frac{1}{2}} y_k$ ,  $\tilde{H}_k = G^{-\frac{1}{2}} H_k G^{-\frac{1}{2}}$

• as before, define  $\tilde{m}_k = \frac{\tilde{y}_k^T \tilde{s}_k}{\tilde{s}_k^T \tilde{s}_k}$ ,  $\tilde{M}_k = \frac{\|\tilde{y}_k\|_2^2}{\tilde{y}_k^T \tilde{s}_k}$ ,  $\tilde{q}_k = \frac{\tilde{s}_k^T \tilde{H}_k \tilde{s}_k}{\tilde{s}_k^T \tilde{s}_k}$ ,  $\cos \tilde{\theta}_k = \frac{\tilde{s}_k^T \tilde{H}_k \tilde{s}_k}{\|\tilde{s}_k\|_2 \|\tilde{H}_k \tilde{s}_k\|_2}$

• as before,  $\tilde{H}_{k+1} = \tilde{H}_k - \frac{\tilde{H}_k \tilde{s}_k \tilde{s}_k^T \tilde{H}_k}{\tilde{s}_k^T \tilde{H}_k \tilde{s}_k} + \frac{\tilde{y}_k \tilde{y}_k^T}{\tilde{y}_k^T \tilde{s}_k}$ ,  $\psi(\tilde{H}_{k+1}) = \psi(\tilde{H}_k) + (\tilde{M}_k - \ln \tilde{m}_k - 1) + \left[ 1 - \frac{\tilde{q}_k}{\cos^2 \tilde{\theta}_k} + \ln \frac{\tilde{q}_k}{\cos^2 \tilde{\theta}_k} \right] + \ln \cos^2 \tilde{\theta}_k$

•  $y_k - G s_k = \int_0^1 D^2 f(x_k + \tau s_k) - G d\tau s_k$

$$\Rightarrow \|\tilde{y}_k - \tilde{s}_k\|_2 \leq \underbrace{\|G^{-\frac{1}{2}}\|_2}_{\leq 1/\sqrt{m}} \|y_k - G s_k\|_2 \leq \sqrt{\frac{1}{m}} L \varepsilon_k \|s_k\|_2 \leq \frac{1}{m} L \varepsilon_k \|\tilde{s}_k\|_2 \quad \text{for } \varepsilon_k = \max(\|x_k - x^*\|_2, \|x_{k+1} - x^*\|_2)$$

• we have  $\tilde{y}_k^T \tilde{s}_k = (\tilde{s}_k + (\tilde{y}_k - \tilde{s}_k))^T \tilde{s}_k \geq \|\tilde{s}_k\|_2^2 (1 - \frac{L}{m} \varepsilon_k)$ ,  $\|\tilde{y}_k\|_2^2 \leq \|\tilde{s}_k\|_2^2 (1 + \frac{L}{m} \varepsilon_k)^2$

$$\Rightarrow \ln \tilde{m}_k \geq \ln(1 - \frac{L}{m} \varepsilon_k) \geq -C \varepsilon_k \quad ; \quad \tilde{M}_k \leq \frac{(1 + \frac{L}{m} \varepsilon_k)^2}{1 - \frac{L}{m} \varepsilon_k} \leq 1 + C \varepsilon_k \quad \text{for some } C > 0$$

$$\Rightarrow 0 < \psi(\tilde{H}_{k+1}) \leq \psi(\tilde{H}_k) + 3C \varepsilon_k + \ln \cos^2 \tilde{\theta}_k + \left[ 1 - \frac{\tilde{q}_k}{\cos^2 \tilde{\theta}_k} + \ln \frac{\tilde{q}_k}{\cos^2 \tilde{\theta}_k} \right] \quad \text{use previous thm \& geometric series}$$

$$\Rightarrow \sum_{j=0}^{\infty} \left( \ln \frac{1}{\cos^2 \tilde{\theta}_j} - \left[ 1 - \frac{\tilde{q}_j}{\cos^2 \tilde{\theta}_j} + \ln \frac{\tilde{q}_j}{\cos^2 \tilde{\theta}_j} \right] \right) \leq \psi(\tilde{H}_0) + 3C \sum_{j=0}^{\infty} \varepsilon_j < \infty, \text{ i.e. } \lim_{j \rightarrow \infty} \cos \tilde{\theta}_j = 1, \lim_{j \rightarrow \infty} \tilde{q}_j = 1$$

$$\cdot \frac{\|G^{-\frac{1}{2}}(H_k - G)s_k\|_2^2}{\|G^{1/2}s_k\|_2^2} = \frac{\|(\tilde{H}_k - I)\tilde{s}_k\|_2^2}{\|\tilde{s}_k\|_2^2} = \frac{\tilde{q}_k^2}{\cos^2 \tilde{\theta}_k} - 2\tilde{q}_k + 1 \xrightarrow{k \rightarrow \infty} 0$$

*now apply earlier convergence result!*

$$\Rightarrow \frac{\|p_k - p_k^*\|_2}{\|p_k\|_2} \leq \frac{\|D^2 f(x_k)(p_k - p_k^*)\|_2}{m \|p_k\|_2} = \frac{\|(D^2 f(x_k) - H_k)p_k\|_2}{m \|p_k\|_2} \leq \frac{\|(H_k - G)p_k\|_2 + \|(G - D^2 f(x_k))p_k\|_2}{m \|p_k\|_2} \xrightarrow{k \rightarrow \infty} 0 \quad \square$$

## Simple treatment of linear constraints

general :  $\min_{x \in \mathbb{R}^n} f(x)$  s.t.  $Ax = b$ ;  $f \in C^2(\mathbb{R}^n)$  convex,  $A \in \mathbb{R}^{p \times n}$ ,  $\text{rank } A = p < n$  (P)

• KKT condition :  $\exists \lambda \in \mathbb{R}^p$  s.t.  $\begin{cases} Ax^* = b \\ \nabla f(x^*) + A^T \lambda = 0 \end{cases}$

— example on next slide!

Elimination of constraints : • find  $F \in \mathbb{R}^{n \times (n-p)}$  with  $\text{range } F = \ker A$  &  $\hat{x}$  with  $A\hat{x} = b$

• (P)  $\Leftrightarrow \min_{z \in \mathbb{R}^{n-p}} f(Fz + \hat{x})$  (unconstrained optim.)

quadratic :  $\min_{x \in \mathbb{R}^n} \frac{1}{2} x^T P x + q^T x + r$  s.t.  $Ax = b$ ;  $P \in \mathbb{R}^{n \times n}$  sym. pos. semi-def.,  $q \in \mathbb{R}^n$ ,  $r \in \mathbb{R}$

• KKT condition :  $\exists \lambda \in \mathbb{R}^p$  s.t.  $\underbrace{\begin{pmatrix} P & A^T \\ A & 0 \end{pmatrix}}_{\text{KKT matrix}} \begin{pmatrix} x^* \\ \lambda \end{pmatrix} = \begin{pmatrix} -q \\ b \end{pmatrix}$

- if KKT matrix regular  $\Rightarrow \exists!$  primal-dual optimal pair  $(x^*, \lambda)$

- if KKT matrix singular &  $\exists$  solution  $\Rightarrow$  every solution is primal dual optimal pair

- if KKT matrix singular & there is no solution  $\Rightarrow p^* = -\infty$

# Optimization algorithms: Projection methods for equality-constrained optim.

## Projection methods for linear constraints

$$\min_{x \in \mathbb{R}^n} f_0(x) \quad \text{s.t.} \quad Ax = b, \quad A \in \mathbb{R}^{p \times n}, \text{rank } A = p < n$$

Thm: The orthogonal projection of  $d \in \mathbb{R}^n$  onto the tangent space  $T = \{s \in \mathbb{R}^n \mid As = 0\}$  to the set of feasible points is given by  $\text{proj}_T d = (I - A^T(AA^T)^{-1}A)d$ .

proof: write  $d = d_1 + d_2$ ,  $d_1 = \text{proj}_T d \in \ker A$ ,  $d_2 \in (\ker A)^\perp = \text{range } A^T$

$$\Rightarrow d = d_1 + A^T s \text{ for some } s \in \mathbb{R}^p \Rightarrow Ad = (AA^T)s \Rightarrow s = (AA^T)^{-1}Ad \Rightarrow d_1 = d - A^T(AA^T)^{-1}Ad \quad \square$$

Rem:  $-\text{proj}_T \nabla f_0(x)$  is a descent direction, since  $(-\text{proj}_T \nabla f_0(x)) \cdot \nabla f_0(x) = -\|\text{proj}_T \nabla f_0(x)\|_2^2$ .

Def: The linesearch methods with descent direction

$$p_k = \underset{p \in \mathbb{R}^n, Ap=0}{\text{argmin}} \quad f_0(x_k) + Df(x_k)p$$

$$p_k^N = \underset{p \in \mathbb{R}^n, Ap=0}{\text{argmin}} \quad f_0(x_k) + Df(x_k)p + \frac{1}{2} p^T D^2 f(x_k)p$$

} minimise first- / second order Taylor approximation under constraints

are called projected gradient / Newton descent, respectively. The directions are computed via


$$p_k = -\text{proj}_T \nabla f_0(x_k) \quad , \quad \begin{pmatrix} D^2 f_0(x_k) & A^T \\ A & 0 \end{pmatrix} \begin{pmatrix} p_k^N \\ w \end{pmatrix} = \begin{pmatrix} -\nabla f_0(x_k) \\ 0 \end{pmatrix}.$$

# Optimization algorithms: Projection methods for equality-constrained optim.

## Projection methods for nonlinear constraints

$$\min_{x \in \mathbb{R}^n} f_0(x) \quad \text{s.t.} \quad c_E(x) = 0$$

problem: At a feasible point  $x$  there is in general no feasible direction  $p$   
(a direction s.t.  $c_E(x + \alpha p) = 0$  for  $\alpha$  small).



Alg. (projection method):


given:  $f_0: \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $x_0 \in \mathbb{R}^n$ ,  $c_E(x_0) = 0$  tangent space to feasible set

repeat

- 1) choose a descent direction  $p_k \in T = \{p \in \mathbb{R}^n \mid Dc_E(x_k)p = 0\}$
- 2) choose a step length  $\alpha_k > 0$  e.g. projected gradient/Newton step

3)  $x_{k+1} = P(x_k + \alpha_k p_k)$   
↖ projection onto  $\{c_E(x) = 0\}$

4)  $k \leftarrow k + 1$



until  $x_{k+1}$  sufficiently minimises  $f_0$

Rem: For the projection  $P$  there are several possibilities.

• To choose a stepsize  $\alpha_k$  one can use the same criteria as earlier, applied to  $f_0 \circ P$ .


# Optimization algorithms: Projection methods for equality-constrained optim.

## Projection onto constraint set

From a point  $y_k = x_k + \alpha_k p_k$  close to  $x_k$ , need to find a point  $P(y_k) \in \{c_E(x) = 0\}$

• choose  $P(y_k)$  on the constraint set in a direction orthogonal to the tangent plane  $T$  at  $x_k$

$$\Rightarrow P(y_k) = y_k + \nabla c_E(x_k) s \text{ for } s \in \mathbb{R}^p \text{ s.t. } c_E(P(y_k)) = 0$$



• always possible for  $y_k$  close to  $x_k$

• perform Newton's method to find a zero  $s$  of  $c_E(y_k + \nabla c_E(x_k) s) =: h(s)$ , i.e.

$$s_0 = 0, \quad s_{i+1} = s_i - Dh(s_i)^{-1} h(s_i) = s_i - (Dc_E(y_{s_i}) \overset{=: y_{s_i}}{Dc_E(y_{s_i})}^T)^{-1} c_E(y_{s_i}) \quad \text{i.o.}$$

$$y_{s_0} = y_k, \quad y_{s_{i+1}} = y_{s_i} - Dc_E(x_k)^T [Dc_E(y_{s_i}) Dc_E(y_{s_i})^T]^{-1} c_E(y_{s_i})$$

• alternatively, use  $Dc_E(y_{s_i}) \approx Dc_E(x_k) \Rightarrow y_{s_{i+1}} = y_{s_i} - Dc_E(x_k) [Dc_E(x_k) Dc_E(x_k)^T]^{-1} c_E(y_{s_i})$   
same matrices as for gradient projection

Rem: The reduced gradient method introduced later can be seen as a projection method using a different projection  $P$ .

# Optimization algorithms: Projection methods for equality-constrained optim.

## Convergence rate of proj. grad. desc.: preliminaries

Idea: analyze simplified algorithm that asymptotically duplicates the considered method

• let  $S = \{x \in \mathbb{R}^n \mid c_E(x) = 0\}$ ,  $T_x S$  = tangent plane to  $S$  in  $x$


• imagine a bug living on  $S$  minimising  $f_0$  via gradient descent; at each iteration

- the bug chooses the (projected) gradient descent direction  $p_k$  and

- moves in that direction along a straight line with step length  $\alpha_k$

• "straight lines" in  $S$  are so-called geodesics; if iterates  $x_k$  converge against  $x^*$ ,

then the difference between geodesics and straight lines in  $\mathbb{R}^n$  converges to 0 even faster



$\Rightarrow$  the bug's iterates are asymptotically the same as the projected gradient descent ones

$\Rightarrow$  the bug's descent has the same convergence rate

# Optimization algorithms: Projection methods for equality-constrained optim.

## Convergence rate of proj. grad. desc.: geodesics & Lagrangians

Def: The arclength of a smooth curve  $x: [0, T] \rightarrow S$  is given by  $l[x] = \int_0^T |\dot{x}(t)| dt$ .

• A geodesic is a curve  $x(t)$  minimising  $l[x]$  for fixed  $x(0), x(T)$ . (Ex: great circles on sphere)

We shall assume  $|\dot{x}(t)| = 1 \forall t \in [0, T]$  (which can always be achieved).

Rem: Geodesics satisfy  $D C_E(x(t)) \dot{x}(t) = 0$  (from  $0 = \frac{d}{dt} C_E(x(t))$ )

and  $\ddot{x}(t) = \nabla C_E(x(t)) \omega(t)$  for some  $\omega: [0, T] \rightarrow \mathbb{R}^p$  (optimality condition for  $\min_x l[x]$ )

*acceleration only normal, not tangential to S*

The above two differential equations together with  $|\dot{x}(0)| = 1$  can be shown to uniquely

define a curve  $x(t)$ , a geodesic, with  $|\dot{x}(t)| = 1$  as long as  $x(t)$  is regular.

*$(g_1(x), \dots, g_p(x))^T$*

Thm: Let  $L_x(x, \lambda) = Df_0(x) + \lambda^T D C_E(x)$  and  $L_{xx}(x, \lambda) = D^2 f_0(x) + \sum_{i=1}^p \lambda_i D^2 g_i(x)$  denote derivative and

Hessian of the Lagrangian and introduce the Lagrange multiplier  $\lambda(x) = -[D C_E(x) D C_E(x)^T]^{-1} D C_E(x) \nabla f_0(x)$

Along a geodesic  $x(t)$  on  $S$ ,  $\frac{d}{dt} f_0(x(t)) = L_x(x, \lambda(x)) \dot{x}$ ,  $\frac{d^2}{dt^2} f_0(x(t)) = \dot{x}^T L_{xx}(x, \lambda(x)) \dot{x}$ .

proof:  $\frac{d}{dt} f_0(x(t)) = Df_0(x) \dot{x}(t) = \text{proj}_{T_x S} \nabla f_0(x) \cdot \dot{x}(t) = (I - D C_E^T [D C_E D C_E^T]^{-1} D C_E) \nabla f_0 \cdot \dot{x} = L_x(x(t), \lambda(x(t))) \dot{x}$

$\cdot \frac{d^2}{dt^2} f_0(x(t)) = \dot{x}^T D^2 f_0(x) \dot{x} + D^2 f_0(x) \ddot{x}$ ; also  $\dot{\lambda}^T D C_E(x(t)) = 0 \Rightarrow 0 = \dot{x}(t)^T \sum_{i=1}^p \dot{\lambda}_i D^2 g_i(x(t)) \dot{x}(t) + \dot{\lambda}^T D C_E(x(t)) \dot{x}(t)$

adding both equations,  $\frac{d^2}{dt^2} f_0(x(t)) = \dot{x}^T L_{xx}(x, \lambda) \dot{x} + (Df_0(x) + \dot{\lambda}^T D C_E(x)) \ddot{x}$   *$\perp \ddot{x}$  for  $\dot{\lambda} = \lambda(x)$*

□

# Optimization algorithms: Projection methods for equality constrained optim.

## Convergence rate of proj. grad. desc.

(assume  $f \in C^2$ )

Thm: Let the iterates  $x_k$  of geodesic gradient descent converge to  $x^*$  (where for simplicity we take  $\alpha_k$  as the minimising step length), and let  $mI \leq L_{xx}(x, \lambda(x)) \leq MI$  for  $x$  close to  $x^*$ , then asymptotically,  $\frac{f(x_{k+1}) - f(x^*)}{f(x_k) - f(x^*)} \leq \left(\frac{M-m}{M+m}\right)^2$ .

proof: abbreviate  $g(t) = L_x(x(t), \lambda(x(t)))^T$ ,  $H(t) = L_{xx}(x(t), \lambda(x(t)))$ ,  $g_k = L_x(x_k, \lambda(x_k))^T$ ,  $H_k = L_{xx}(x_k, \lambda(x_k))$

let  $x(t)$  be a geodesic with  $x(0) = x^*$ ,  $x(T) = x_k$ , then by Taylor expansion of  $f_0(x(t))$  &  $g(t)$ ,  $\rightarrow \text{dist}(x^*, x_k)$

$$f_0(x^*) - f_0(x_k) = -g_k^T \dot{x}(T)T + \frac{1}{2} T^2 \dot{x}^T(T) H_k \dot{x}(T) + o(T^2) \quad (*)$$

$$g_k = g(T) - g(0) = g'(T)T + o(T) = TH_k \dot{x}(T) + T \nabla_{C_E}(x_k) \frac{d}{dt}(\lambda \circ x)(T) + o(T)$$

let  $P_k$  the orthogonal projection onto  $T_{x_k} S$ , then  $g_k = P_k g_k = P_k H_k \dot{x}(T)T + o(T)$

$$\dot{x}_k = P_k \dot{x}_k$$

$$\tilde{H}_k \dot{x}(T)T = g_k + o(T)$$

$$\text{for } \tilde{H}_k = P_k H_k P_k$$

$$\left( \begin{array}{l} |\dot{x}_k| = 1 \\ \Rightarrow \end{array} \right) mT \leq |g_k| + o(T) \leq MT$$

in (\*)


$$f_0(x^*) - f_0(x_k) = -\frac{1}{2} g_k^T \tilde{H}_k^{-1} g_k + o(T^2) = -\frac{1}{2} g_k^T \tilde{H}_k^{-1} g_k (1 + o(1))$$

now let  $y(t)$  be the geodesic with  $y(0) = x_k$ ,  $\dot{y}(0) = \frac{P_k g_k}{\|P_k g_k\|} = -\frac{g_k}{\|g_k\|}$ , then

$$f_0(y(t)) = f_0(x_k) + t g_k^T \dot{y}(0) + \frac{t^2}{2} \dot{y}^T(0) H_k \dot{y}(0) + o(t^2) \quad \text{is minimised by } t_k = \frac{-g_k^T \dot{y}}{\dot{y}^T H_k \dot{y}} + o(t_k) \sim T$$

$$\Rightarrow f_0(x_k) - f_0(x_{k+1}) = \frac{1}{2} \frac{(g_k^T \dot{y})^2}{\dot{y}^T H_k \dot{y}} + o(T^2) = \frac{1}{2} \frac{(g_k^T \dot{y})^2}{\dot{y}^T H_k \dot{y}} (1 + o(1)) = \frac{1}{2} \frac{\|g_k\|_2^4}{g_k^T \tilde{H}_k g_k} (1 + o(1)) \quad \text{Kantorovich inequality}$$

$$f_0(x_{k+1}) - f_0(x^*) = f_0(x_k) - f_0(x^*) + f_0(x_{k+1}) - f_0(x_k) = [f_0(x_k) - f_0(x^*)] \left(1 - \frac{\|g_k\|_2^4 (1 + o(1))}{g_k^T \tilde{H}_k g_k g_k^T \tilde{H}_k^{-1} g_k}\right) \leq [f_0(x_k) - f_0(x^*)] \left(1 - \frac{c + o(1)}{(M+m) + (m+M)c}\right) \quad \square$$




# Optimization algorithms: Projection methods for equality-constrained optim.

## The Kantorovich inequality

Thm: Let  $p_1, \dots, p_n \geq 0, 0 < m \leq \lambda_1, \dots, \lambda_n \leq M$ , then  $\left(\sum_{i=1}^n p_i \lambda_i\right) \left(\sum_{i=1}^n p_i / \lambda_i\right) \leq \frac{(m+M)^2}{4mM} \left(\sum_{i=1}^n p_i\right)^2$

proof:  $\xi_1 = \frac{p_1}{\sum_{i=1}^n p_i}, \dots, \xi_n = \frac{p_n}{\sum_{i=1}^n p_i}$  are convex combination coefficients (i.e.  $\xi_i \in [0, 1], \sum_{i=1}^n \xi_i = 1$ )

$\Rightarrow (\lambda, r) := \sum_{i=1}^n \xi_i (\lambda_i, h(\lambda_i))$  lies in the convex hull of the  $(\lambda_i, h(\lambda_i))$



$$\Rightarrow r \leq \frac{\lambda - m}{M - m} h(M) + \frac{M - \lambda}{M - m} h(m) = \frac{m + M - \lambda}{mM}$$

$$\lambda_{\max} = \frac{m + M}{2}$$

$$\Rightarrow \left(\sum_{i=1}^n p_i \lambda_i\right) \left(\sum_{i=1}^n \frac{p_i}{\lambda_i}\right) / \left(\sum_{i=1}^n p_i\right)^2 = \left(\sum_{i=1}^n \xi_i \lambda_i\right) \left(\sum_{i=1}^n \frac{\xi_i}{\lambda_i}\right) = \lambda r \leq \lambda \frac{m + M - \lambda}{mM} \leq \max_{\lambda \in \mathbb{R}} \lambda \frac{m + M - \lambda}{mM} = \frac{(m + M)^2}{4mM} \quad \square$$

Cor: Let  $0 < m I \leq H \leq M I$  for a symmetric positive definite  $H \in \mathbb{R}^{n \times n}, x \in \mathbb{R}^n$ , then  $\frac{(x^T H x)(x^T H^{-1} x)}{\|x\|_2^4} \leq \frac{(m+M)^2}{4mM}$

proof: Let  $\lambda_1, \dots, \lambda_n \in [m, M]$  be the eigenvalues of  $H$  with orthonormal eigenvectors  $q_1, \dots, q_n \in \mathbb{R}^n$ .

Write  $x = \sum_{i=1}^n z_i q_i$ , then  $\|x\|_2^2 = \sum_{i=1}^n z_i^2, (x^T H x)(x^T H^{-1} x) = \left(\sum_{i=1}^n \lambda_i z_i^2\right) \left(\sum_{i=1}^n \frac{1}{\lambda_i} z_i^2\right)$ , set  $p_i = z_i^2$   $\square$

Rem: Note  $\frac{(m+M)^2}{4mM} = \frac{1}{4} \left(\sqrt{\frac{m}{M}} + \sqrt{\frac{M}{m}}\right)^2 = \frac{1}{1 - \left(\frac{M-m}{M+m}\right)^2}$ .   
 *condition number*

# Optimization algorithms: Reduced methods for equality-constrained optim.

## Linear constraints

$$\min_{x \in \mathbb{R}^n} f_0(x) \quad \text{s.t.} \quad Ax = b, \quad A \in \mathbb{R}^{p \times n}, \text{rank } A = p < n$$

choose a basis  $B$  of  $\mathbb{R}^p$  from the columns of  $A$  (for simplicity let  $B$  contain the first  $p$  columns)  
and subdivide  $x = (y, z) \in \mathbb{R}^p \times \mathbb{R}^{n-p}$ ,  $A = [B, C] \in \mathbb{R}^{p \times p} \times \mathbb{R}^{p \times (n-p)}$

Rem:  $z$  can be regarded as the independent and  $y = B^{-1}(b - Cz)$  as the dependent variable.

Def:  $\tilde{f}(z) = f_0(\underbrace{B^{-1}(b - Cz)}_{y(z)}, z)$  is called the reduced functional,  
 $\nabla \tilde{f}(z) = \left[ D_z f_0(y(z), z) - D_y f_0(y(z), z) B^{-1} C^T \right]^T$  the reduced gradient,  
and  $D^2 \tilde{f}(z)$  the reduced Hessian.

The reduced gradient/Newton method is a gradient/Newton descent for  $\tilde{f}$ .

## Nonlinear constraints

$$\min_{x \in \mathbb{R}^n} f_0(x) \text{ s.t. } c_E(x) = 0$$

• Let  $x = (y, z) \in \mathbb{R}^p \times \mathbb{R}^{n-p}$  and assume  $\nabla_y c_E(y, z)$  to be regular  $\xRightarrow{\text{implicit function}}$   $y = y(z)$  locally

• Define reduced functional  $\tilde{f}(z) = f_0(y(z), z)$ .

$$c_E(y(z), z) = 0 \Rightarrow D_y c_E(y(z), z) \frac{dy}{dz} + D_z c_E(y(z), z) = 0 \left. \vphantom{c_E} \right\} \Rightarrow \nabla \tilde{f}(z) = \nabla_z f_0(y(z), z) - \nabla_z c_E(y(z), z) \nabla_y c_E(y(z), z)^{-1} \nabla_y f_0(y(z), z)$$

$$\nabla \tilde{f}(z) = \left[ D_y f_0(y(z), z) \frac{dy}{dz} + D_z f_0(y(z), z) \right]^T$$

• analogously for reduced Hessian

• The reduced gradient/Newton method requires to find  $y(z_{k+1})$  for any given  $z_{k+1}$ .

As for projection methods, this can be done by a Newton iteration to find a zero of  $h(y) = c_E(y, z_{k+1})$

$$y_0 = y(z_k), \quad y_{i+1} = y_i - D_y c_E(y_i, z_{k+1})^{-1} c_E(y_i, z_{k+1})$$

or alternatively, using  $D_y c_E(y_i, z_{k+1}) \approx D_y c_E(y(z_k), z_k)$

$$y_{i+1} = y_i - D_y c_E(y(z_k), z_k)^{-1} c_E(y_i, z_{k+1})$$

# Optimization algorithms: Reduced methods for equality-constrained optim.

## Convergence rate of reduced gradient descent

Thm: Abbreviate  $K(z) = \frac{dy}{dz} = -D_y c_E(y(z), z)^{-1} D_z c_E(y(z), z)$  and  $C'(z) = \begin{pmatrix} K(z) \\ I \end{pmatrix} \in \mathbb{R}^{n \times (n-p)}$ , then for any  $\lambda \in \mathbb{R}^p$  satisfying  $D_y f_0(y(z), z) + \lambda^T D_y c_E(y(z), z) = 0$  we have  $D^2 \tilde{f}(z) = C'(z)^T L_{xx}(y(z), z, \lambda) C'(z)$ .

proof:  $\tilde{f}(z) = f_0(y(z), z) = f_0(y(z), z) + \lambda^T c_E(y(z), z)$

$$\cdot D \tilde{f}(z) = [D_y f_0(y(z), z) + \lambda^T D_y c_E(y(z), z)] K(z) + D_z f_0(y(z), z) + \lambda^T D_z c_E(y(z), z) = \underbrace{0}_{=0} L_y(y(z), z, \lambda) K(z) + L_z(y(z), z, \lambda)$$

$$\cdot D^2 \tilde{f}(z) = K(z)^T L_{yy}(y(z), z, \lambda) K(z) + L_{zy}(y(z), z, \lambda) K(z) + K(z)^T L_{yz}(y(z), z, \lambda) + L_{zz}(y(z), z, \lambda) \square$$

Thm: Let the iterates  $x_k$  of reduced gradient descent converge to  $x^*$  (where for simplicity we take  $\alpha_k$  as the minimising step length), and let  $mI \leq C'(z)^T L_{xx}(x, \lambda(x)) C'(z) \leq MI$  for  $x$  close to  $x^*$ , then asymptotically,  $\frac{f(x_{k+1}) - f(x^*)}{f(x_k) - f(x^*)} \leq \left(\frac{M-m}{M+m}\right)^2$ .

proof: homework  $\square$

Rem: The convergence rate of the projected gradient descent depends on the eigenvalues of  $\tilde{H} = P^T L_{xx}(x^*, \lambda(x^*)) P$  for  $P$  the orthogonal projection onto  $T_{x^*} S$ . We can write  $P = C'(z^*) (C'(z^*)^T C'(z^*))^{-\frac{1}{2}}$  (indeed, the columns of  $C$  span  $T_{x^*} S$  since  $(D_y c_E(x^*) D_z c_E(x^*)) C = 0$ , thus the columns of  $P$  form an orthonormal basis of  $T_{x^*} S$  due to  $P^T P = I$ ). Hence, the convergence rate of the reduced gradient descent depends on the eigenvalues of  $\sqrt{C'^T C'} \tilde{H} \sqrt{C' C'}$  <sup>depends on chosn 2</sup> and this is coordinate-dependent

# Optimization algorithms: Interior point methods for ineq.-constrained optim.

## Idea of interior point methods

$$\min f_0(x) \text{ s.t. } f_1(x), \dots, f_m(x) \leq 0, Ax = b, f_i \in C^2, \text{convex}, A \in \mathbb{R}^{p \times n}, \text{rank } A = p < n \quad (P)$$

assume: • optimal point  $x^*$  exists

• Slater's condition (strict feasibility) satisfied


⇒ strong duality, KKT conditions fulfilled

$$\left. \begin{aligned} Ax^* = b, f_i(x^*) \leq 0, \mu_i^* \geq 0, \mu_i^* f_i(x^*) = 0, i = 1, \dots, m \\ \nabla f_0(x^*) + \sum_{i=1}^m \mu_i^* \nabla f_i(x^*) + A^T \lambda^* = 0 \end{aligned} \right\} \quad (0)$$

idea: • Solve (P) or (0) by applying Newton's method to a sequence of modifications of (P) or (0) that only have equality constraints.

• E.g., replace  $f_i(x) \leq 0$  by adding a "barrier function" to  $f_0$

• interior point methods produce a sequence  $x_k$  with  $Ax_k = b, f_i(x_k) < 0, x_k \rightarrow x^*$



## Barrier methods: barrier functions

$$(P) \Leftrightarrow \min_x f_0(x) + \sum_{i=1}^m \chi_{-}(f_i(x)) \quad \text{s.t. } Ax = b \quad \text{for } \chi_{-}(s) = \begin{cases} 0 & \text{if } s \leq 0 \\ \infty & \text{else} \end{cases}$$

replace  $\chi_{-}(u)$  by  $I_{\epsilon}(u) = -\frac{1}{\epsilon} \ln(-u)$ ,  $\text{dom } I_{\epsilon} = \{u \in \mathbb{R} \mid u < 0\}$

Thm: For  $\epsilon > 0$ ,  $I_{\epsilon}$  is convex, differentiable, non-decreasing

$$\cdot I_{\epsilon}(u) \xrightarrow{u \uparrow 0} \infty, I_{\epsilon}(u) = 0 \text{ for } u > 0$$

□

Def: The logarithmic barrier function  $\phi$  to (P) is given by

$$\phi(x) = -\sum_{i=1}^m \ln(-f_i(x)), \quad \text{dom } \phi = \{x \in \mathbb{R}^n \mid f_1(x), \dots, f_m(x) < 0\}$$

$$\nabla \phi(x) = -\sum_{i=1}^m \frac{1}{f_i(x)} \nabla f_i(x), \quad \mathbb{D}^2 \phi(x) = \sum_{i=1}^m \frac{1}{f_i^2(x)} \nabla f_i(x) \nabla f_i(x)^T - \frac{1}{f_i(x)} \mathbb{D}^2 f_i(x)$$

$\phi$  is convex.

• The barrier problem to (P) is given by

$$\min_x t f_0(x) + \phi(x) \quad \text{s.t. } Ax = b \quad (P_{\epsilon})$$

• The central path is  $\{x^*(t) \mid t > 0\}$ , where  $x^*(t)$  is the optimal point of  $(P_{\epsilon})$ ; it

$$\text{satisfies } \exists \lambda \in \mathbb{R}^r: 0 = t \nabla f_0 + \nabla \phi + A^T \lambda$$

# Optimization algorithms: Interior point methods for ineq.-constrained optim.

## Barrier methods: algorithm

Thm: Let  $(x^*, \lambda^*)$  be optimal primal dual pair for  $(P_\varepsilon)$ , then  $(\mu^*, \lambda^*)$  is dually feasible for  $\mu_i^* = \frac{-1}{\varepsilon f_i(x^*)}$ , and  $f_0(x^*) - g(\mu^*, \lambda^*) = \frac{m}{\varepsilon}$ , i.e.  $x^*$  and  $(\mu^*, \lambda^*)$  are  $\frac{m}{\varepsilon}$ -suboptimal.

proof:  $\mu_i^* > 0$  due to  $f_i(x^*) < 0$

• optimality condition for  $(P_\varepsilon)$ :  $0 = \nabla f_0 + \frac{\nabla \phi}{\varepsilon} + A^T \lambda^* = \nabla f_0(x^*) + \sum_{i=1}^m \mu_i^* \nabla f_i(x^*) + A^T \lambda^*$

$\Rightarrow x^* = \underset{x}{\operatorname{argmin}} (f_0(x) + \sum_{i=1}^m \mu_i^* f_i(x) + \lambda^{*T} (Ax - b)) = \underset{x}{\operatorname{argmin}} L(x, (\mu^*, \lambda^*))$

•  $p^* \geq g(\mu^*, \lambda^*) = \underset{x}{\operatorname{min}} L(x, (\mu^*, \lambda^*)) = L(x^*, (\mu^*, \lambda^*)) = f_0(x^*) - \frac{1}{\varepsilon} \sum_{i=1}^m \frac{f_i(x^*)}{f_i(x^*)}$  □

Alg (barrier method): given: strictly feasible  $x$ ,  $t = t_0 > 0$ ,  $\beta > 1$ , tolerance  $\varepsilon > 0$

iterate until  $\frac{m}{t} < \varepsilon$ :

step 1: solve  $(P_t)$  via Newton's method with initial guess  $x$

step 2:  $x \leftarrow x^*(t)$ ,  $t \leftarrow \beta t$

Rem: The previous theorem implies convergence  $f_0(x^*(t)) \rightarrow p^*$ .

## Barrier methods: complexity

usually not fulfilled for barrier function  
but can be relaxed to a "self-concordance"  
assumption

Thm: Let  $D^2 f_0 + \frac{1}{\epsilon} D^2 \phi$  Lipschitz with constant  $L$ ,  $\tilde{m} I \leq D^2 f_0(x) + \frac{1}{\epsilon} D^2 \phi \leq M I$   
for some  $\tilde{m}, M > 0$  and all  $\epsilon > 0$ , then the barrier method reaches an  $x$  with  $\|x - x^*\|_2 < \epsilon$   
after  $N = \frac{\log \frac{\tilde{m}}{\epsilon_0 \epsilon}}{\log \beta} \left( \frac{m(\beta - 1 - \log \beta)}{\gamma} + C \log_2 \log \frac{1}{\epsilon} \right)$  steps for some  $C, \gamma > 0$ ,  
where  $\tilde{\epsilon}$  is the accuracy of solving  $(P_\epsilon)$ .

proof: • every iteration takes  $\frac{f(x) - p^*}{\gamma} + C \log_2 \log \frac{1}{\tilde{\epsilon}}$  Newton steps starting from  $x$   
where  $f = \epsilon f_0 + \phi$ ,  $x = x^*(\epsilon/\beta)$ ,  $x^+ = x^*(\epsilon)$ ,  $p^* = f(x^+)$

•  $\epsilon f_0(x) + \phi(x) - \epsilon f_0(x^+) - \phi(x^+) = \epsilon f_0(x) - \epsilon f_0(x^+) + \sum_{i=1}^m (-\log(-\mu_i(x)) + \log(-\mu_i(x^+)))$

$\mu_i = \frac{\beta}{-\epsilon f_i(x)}$  (multiplier from previous iteration)

$$= \epsilon f_0(x) - \epsilon f_0(x^+) + \sum_{i=1}^m \underbrace{\log(-\mu_i f_i(x^+))}_{\leq -\mu_i f_i(x^+) - 1} - m \log \beta$$

$$\leq \epsilon f_0(x) - \epsilon \left( f_0(x^+) + \sum_{i=1}^m \mu_i f_i(x^+) + \lambda^T (A x^+ - b) \right) - m - m \log \beta \leq m(\beta - 1 - \log \beta)$$

(similar to duality gap on previous slide!)

$\Rightarrow$  need less than  $\frac{m(\beta - 1 - \log \beta)}{\gamma} + C \log_2 \log \frac{1}{\tilde{\epsilon}}$  Newton steps

• in total  $\log \frac{\tilde{m}}{\epsilon_0 \epsilon} / \log \beta$  outer iterations

□

# Optimization algorithms: Interior point methods for ineq.-constrained optim.

## Barrier methods: finding a feasible starting point

Barrier methods require a strictly feasible starting point  $x_0$  with  $c_{\mathcal{I}}(x_0) = \begin{pmatrix} f_1 \\ \vdots \\ f_m \end{pmatrix}(x_0) < 0$ ,  $Ax_0 = b$ .

This is found by an auxiliary optimisation, a so-called phase I method.

method 1:  $\min_{x \in \mathbb{R}^n, s \in \mathbb{R}} s$  s.t.  $f_1(x), \dots, f_m(x) \leq s$ ,  $Ax = b$  - the actual optimisation problem is often called phase II

- strictly feasible optimisation problem, just choose some  $x$  with  $Ax = b$  and  $s > \max_i f_i(x)$

$\Rightarrow$  apply barrier method

- $p^* < 0 \Rightarrow$  feasible  $x_0$  exists (stop phase I method as soon as  $s < 0$  and take  $x$  as  $x_0$ )

$p^* > 0 \Rightarrow$  original problem infeasible

$p^* = 0 \Rightarrow$  original problem  $\begin{cases} \text{not feasible if minimum not attained} \\ \text{not strictly feasible else} \Rightarrow \text{barrier method not applicable} \end{cases}$

method 2:  $\min_{x \in \mathbb{R}^n, s \in \mathbb{R}^m} s_1 + \dots + s_m$  s.t.  $f_1(x) \leq s_1, \dots, f_m(x) \leq s_m$ ,  $Ax = b$ ,  $s_1, \dots, s_m \geq 0$

- also strictly feasible  $\Rightarrow$  apply barrier method

- if  $p^* > 0$ , at least some  $f_i(x)$  are  $> 0$ ; only those constraints  $f_i(x) \leq 0$  for which  $s_i > 0$

(and corresponding dual variables are 0) are violated

Optimization algorithms: Interior point methods for ineq.-constrained optim.

Barrier methods: phase I termination near phase II central path

Assume we know that a feasible point can be found with  $f_0(x) \leq M$ , then choose

$$\text{phase I: } \min_{x \in \mathbb{R}^n, s \in \mathbb{R}} s \quad \text{s.t. } f_1(x), \dots, f_m(x) \leq s, f_0(x) \leq M, Ax = 0$$

• The central path  $(s(t), x(t))$  for the logarithmic barrier method applied to this satisfies the optimality conditions  $t = \sum_{i=1}^m \frac{1}{s-f_i(x)}$ ,  $0 = \frac{1}{M-f_0(x)} \nabla f_0(x) + \sum_{i=1}^m \frac{1}{s-f_i(x)} \nabla f_i(x) + A^T \lambda$

• The central path  $x(t)$  for the barrier method applied to the original problem satisfies

$$t \nabla f_0(x) + \sum_{i=1}^m \frac{1}{-f_i(x)} \nabla f_i(x) + A^T \lambda = 0$$

•  $\Rightarrow$  If  $(s, x)$  lies on the central path of phase I with  $s=0$ , then  $x$  lies on the central path of phase II with parameter  $t = \frac{1}{M-f_0(x)}$  and duality gap  $m(M-f_0(x))$

# Optimization algorithms: Interior point methods for ineq.-constrained optim.

## Barrier methods: complexity of phase I

- Assume, we know the feasible set to lie in a ball of radius  $R$  (for simplicity no eq. constraints)
- Consider phase I problem  $\min_{x \in \mathbb{R}^n, s \in \mathbb{R}} s$  s.t.  $f_1(x), \dots, f_m(x) \leq s, a^T x \leq 1$   
 where  $a$  will be chosen with  $\|a\|_2 \leq \frac{1}{R}$  ( $\Rightarrow a^T x \leq 1$  is redundant); start with  $s = s_0, x = 0$
- Choose  $a$  &  $s_0$  such that  $(s = s_0, x = 0)$  lies on central path for phase I, i.e.

$$t_0 = \sum_{i=1}^m \frac{1}{s_0 - f_i(0)} \quad \text{and} \quad a = - \sum_{i=1}^m \frac{1}{s_0 - f_i(0)} \nabla f_i(0).$$

Also need  $\|a\|_2 \leq \frac{1}{R}$ :  $\|a\|_2 \leq \sum_{i=1}^m \frac{1}{s_0 - f_i(0)} \|\nabla f_i(0)\|_2 \leq \frac{mG}{s_0 - F}$  for  $G = \max_i \|\nabla f_i(0)\|_2, F = \max_i f_i(0)$

$\Rightarrow$  choose  $s_0 = mGR + F$  note: phase I has  $m+1$  constraints

- Initial duality gap  $\frac{m+1}{t_0} = (m+1)mGR / \sum_{i=1}^m \frac{1}{1 + (F - f_i(0)) / (mGR)} \leq (m+1)mGR$

needed accuracy  $\varepsilon = |p^*|$  (only need to determine whether  $p^* > 0$  or  $p^* < 0$ )  
(= measure of difficulty of determining feasibility)

$\Rightarrow$  number of Newton steps needed:

take  $\beta = 1 + \frac{1}{\sqrt{m+1}}$ ,  $\log(1+x) = x - \frac{1}{2}(\frac{x}{\beta})^2$  for some  $z \in [1, 1+x]$

$$N = \frac{\log \frac{m+1}{\log \beta} \varepsilon}{\log \beta} \left( \frac{(m+1)(\beta - 1 - \log \beta)}{\beta} + G \log_2 \log \frac{1}{\varepsilon} \right) \leq C_{m+1} \sqrt{m+1} \log \left( \frac{m(m+1)GR}{|p^*|} \right) \left( \frac{1}{2\beta} + c \right)$$

tends to  $\infty$  for  $\beta \rightarrow 1$  or  $\beta \rightarrow \infty \Rightarrow$  there is optimal  $\beta$

# Optimization algorithms: Interior point methods for ineq.-constrained optim.

## Primal dual interior point methods

Similar to barrier methods, but :

- no inner/outer iterations

- search directions from Newton's method applied to KKT conditions
- primal & dual iterates not necessarily feasible
- can exhibit superlinear convergence
- can be applied if problem not strictly feasible

Rem: In the logarithmic barrier method, the Newton step  $p_k^u$  of the inner iterations solves

$$\begin{pmatrix} D^2 f_0(x) + \frac{1}{\epsilon} D^2 \phi(x) & A^T \\ A & 0 \end{pmatrix} \begin{pmatrix} p_k^u \\ \lambda \end{pmatrix} = - \begin{pmatrix} \nabla f_0(x) + \frac{1}{\epsilon} \nabla \phi(x) \\ 0 \end{pmatrix}. \quad (*)$$

This can be interpreted as a Newton step for solving the modified KKT system

$$\begin{aligned} Df_0(x) + \sum_{i=1}^m \mu_i Df_i(x) + \lambda^T A &= 0 \\ -\mu_i f_i(x) &= \frac{1}{\epsilon}, \quad i=1, \dots, m \\ Ax - b &= 0 \end{aligned} \quad (m\text{KKT})$$

Indeed, eliminating  $\mu_i = -1/\epsilon f_i(x)$ , the above becomes  $Df_0(x) + \frac{1}{\epsilon} D\phi(x) + \lambda^T A = 0$ ,  $Ax - b = 0$  for which (\*) is a Newton step starting from an  $(x, \lambda)$  with  $Ax = b$ ,  $\lambda = 0$ .

The primal-dual search direction will be a Newton step for (mKKT) in  $(x, \mu, \lambda)$ .

## Primal dual interior point methods: algorithm

Def.: The primal dual step is a Newton step for (mKKT), i.e.,  $p_k^{pd} = (\Delta x, \Delta \mu, \Delta \lambda)$  with

$$\begin{pmatrix} D^2 f_0(x) + \sum_{i=1}^m \mu_i D^2 f_i(x) & \nabla c_I(x) & A^T \\ -\text{diag}(\mu_1, \dots, \mu_m) & DC_I(x) & 0 \\ A & 0 & 0 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta \mu \\ \Delta \lambda \end{pmatrix} = - \begin{pmatrix} r_{dual} \\ r_{central} \\ r_{primal} \end{pmatrix} := - \begin{pmatrix} \nabla f_0(x) + \nabla c_I(x) \mu + A^T \lambda \\ -\text{diag}(\mu_1, \dots, \mu_m) c_I(x) - \frac{1}{t} \\ Ax - b \end{pmatrix}$$

for the dual, central, and primal residual  $r_{dual}, r_{central}, r_{primal}$ .

The surrogate duality gap is  $\eta(x, \mu) = -\mu^T c_I(x)$ .

Rem: Since  $x$  need not be feasible (unlike in barrier methods), we cannot compute a duality gap, but

$$\left. \begin{array}{l} r_{dual} = 0, \mu \geq 0 \Rightarrow x = \text{argmin } L(x, (\mu, \lambda)) \\ r_{primal} = 0 \Rightarrow Ax = b \end{array} \right\} \Rightarrow p^* - d^* \leq f_0(x) - L(x, (\mu, \lambda)) = \eta(x, \mu)$$

then surrogate duality gap is true duality gap

Alg: given  $x_0$  with  $f_1(x_0), \dots, f_m(x_0) < 0, \mu_0 > 0, \lambda_0, \beta > 1, \varepsilon_{feas}, \varepsilon > 0$

repeat 1)  $t := \frac{\beta m}{\eta(x_k, \mu_k)}$  2) compute  $p_k^{pd}$  3) choose step size  $\alpha_k > 0$  4)  $\begin{pmatrix} x_{k+1} \\ \mu_{k+1} \\ \lambda_{k+1} \end{pmatrix} = \begin{pmatrix} x_k \\ \mu_k \\ \lambda_k \end{pmatrix} + \alpha_k p_k^{pd}$

until  $\|r_{primal}\|_2 < \varepsilon_{feas}, \|r_{dual}\|_2 < \varepsilon_{feas}, \eta(x_k, \mu_k) < \varepsilon$

Rem: If  $x_k, \mu_k, \lambda_k$  solved (mKKT),  $\eta$  would be the duality gap and  $\frac{m}{\eta}$  the barrier parameter  $t \Rightarrow$  step 1) means  $t \leftarrow \beta t$

# Optimization algorithms: Interior point methods for ineq.-constrained optim.

## Primal dual interior point methods: linesearch

Step 3) can be done via backtracking, using Armijo's condition, i.e., starting from  $\alpha_k = 1$ ,

$$\text{repeat } \alpha_k \leftarrow \frac{\alpha_k}{2} \text{ until } \begin{cases} c_I(x_k + \alpha_k \Delta x) < 0 \\ \mu_k + \alpha_k \Delta \mu_k > 0 \\ \|\tau((x_k, \mu_k, \lambda_k) + \alpha_k p_k^{pd})\|_2 \leq (1 - c_0 \alpha_k) \|\tau(x_k, \mu_k, \lambda_k)\|_2 \end{cases}$$

(  $\tau_{\text{primal}}$ ,  $\tau_{\text{central}}$ ,  $\tau_{\text{dual}}$  )

Thm: The backtracking terminates.

proof:  $c_I < 0$  and  $\mu > 0$  are satisfied for  $\alpha_k$  small enough by continuity.

• Let  $y = (x_k, \mu_k, \lambda_k)$ , then  $p_k^{pd} = -D\tau(y)^{-1} \tau(y)$

$$\Rightarrow \frac{d}{dt} \|\tau(y + t p_k^{pd})\|_2^2 \Big|_{t=0} = 2 \tau(y)^T D\tau(y) p_k^{pd} = -2 \tau(y)^T \tau(y)$$

$$\Rightarrow \frac{d}{dt} \sqrt{\|\tau(y + t p_k^{pd})\|_2^2} \Big|_{t=0} = -\|\tau(y)\|_2 \Rightarrow \|\tau(y + t p_k^{pd})\|_2 = (1 - t) \|\tau(y)\|_2 + O(t^2) \quad \square$$

Thm: Once a full Newton step is taken,  $\tau_{\text{primal}} = 0$  (i.e.,  $Ax_k = b$ ) for all following iterations.

proof: In that step,  $Ax_k = b$ ; but then, all following directions  $\Delta x$  are feasible.  $\square$

Thm: In each step,  $\tau_{\text{primal}}$  decreases by the factor  $(1 - \alpha_k)$ .

proof:  $Ax_{k+1} - b = Ax_k - b + \alpha_k A \Delta x = (1 - \alpha_k) (Ax_k - b)$   $\square$


# Optimization algorithms: Noninterior methods for ineq.-constrained optim.

## Penalty methods

$\min_{x \in \mathbb{R}^n} f_0(x)$  s.t.  $c_I(x) = (f_1(x), \dots, f_m(x)) \leq 0$  replace potential equality constraints  $g_i(x) = 0$  by  $g_i(x) \leq 0$  &  $-g_i(x) \leq 0$

Def: A penalty function for the constraints  $f_1(x), \dots, f_m(x) \leq 0$  is a function  $P(x) = \gamma(c_I^+(x))$  with  $c_I^+(x) = (\max\{0, f_1(x)\}, \dots, \max\{0, f_m(x)\})$  and  $\gamma: [0, \infty)^m \rightarrow [0, \infty)$  such that  $\gamma(0) = 0$ ,  $\gamma(p) > 0$  for  $p \neq 0$ ,  $\gamma$  continuous

Ex:  $\gamma(p) = \frac{1}{2} p^T \Gamma p$  for a positive definite  $\Gamma \in \mathbb{R}^{m \times m}$   
case  $\Gamma = I$  yields  $P(x) = \frac{1}{2} \sum_{i=1}^m (\max\{0, f_i(x)\})^2$



Alg (penalty method): given  $0 < c_1 < c_2 < \dots$ ,  $\lim_{k \rightarrow \infty} c_k = \infty$

for  $k = 1, 2, \dots$  do  $x_k = \underset{x}{\operatorname{argmin}} q(c_k, x) = \underset{x}{\operatorname{argmin}} f_0(x) + c_k P(x)$

We shall assume a solution  $x_k$  to exist for each  $k$ .

# Optimization algorithms: Noninterior methods for ineq.-constrained optim.

## Penalty methods: convergence

Thm: We have  $q(c_k, x_k) \leq q(c_{k+1}, x_{k+1})$ ,  $P(x_k) \geq P(x_{k+1})$ ,  $f_0(x_k) \leq f_0(x_{k+1})$ .

proof:  $q(c_{k+1}, x_{k+1}) = f_0(x_{k+1}) + c_{k+1} P(x_{k+1}) \geq f_0(x_{k+1}) + c_k P(x_{k+1}) \geq f_0(x_k) + c_k P(x_k) = q(c_k, x_k)$

$$\cdot f_0(x_k) + c_k P(x_k) \leq f_0(x_{k+1}) + c_k P(x_{k+1}) + f_0(x_{k+1}) + c_{k+1} P(x_{k+1}) \leq f_0(x_k) + c_{k+1} P(x_k) = (c_{k+1} - c_k) P(x_k) \leq (c_{k+1} - c_k) P(x_{k+1})$$

$$\cdot f_0(x_{k+1}) + c_k P(x_{k+1}) \geq f_0(x_k) + c_k P(x_k) \geq f_0(x_k) + c_k P(x_{k+1}) \Rightarrow f_0(x_{k+1}) \geq f_0(x_k) \quad \square$$

Thm: Let  $x^*$  solve (P), then  $f_0(x^*) \geq q(c_k, x_k) \geq f_0(x_k)$ .

proof:  $f_0(x^*) = f_0(x^*) + c_k P(x^*) \geq f_0(x_k) + c_k P(x_k) \geq f_0(x_k) \quad \square$

Thm: Let  $x_1, x_2, \dots$  be generated by the penalty method, then any limit point solves (P).

proof:  $\cdot$  Let  $K \subset \mathbb{N}$  indicate a subsequence with  $\lim_{k \in K} x_k = x$  (thus  $\lim_{k \in K} f_0(x_k) = f_0(x)$ )

$\cdot q(c_k, x_k)$  is monotone & bounded by  $f_0(x^*)$ , thus  $\lim_{k \in K} q(c_k, x_k) = q^* \leq f_0(x^*)$

$\cdot \lim_{k \in K} c_k P(x_k) = \lim_{k \in K} q(c_k, x_k) - f_0(x_k) = q^* - f_0(x) \Rightarrow P(x) = \lim_{k \in K} P(x_k) = 0$   <sup>$x$  feasible!</sup>

$\cdot f_0(x) = \lim_{k \in K} f_0(x_k) \leq f_0(x^*)$   <sup>$x$  optimal!</sup> by above thm □

# Optimization algorithms: Noninterior methods for ineq.-constrained optim.

## Active set methods

$$\min_{x \in \mathbb{R}^n} f_0(x) \quad \text{s.t.} \quad f_1(x), \dots, f_m(x) \leq 0, \quad g_1(x) = \dots = g_p(x) = 0 \quad (P)$$

Def: Given the optimal point  $x^* \in \mathbb{R}^n$ , the active set is defined as  $A = \{i \in \{1, \dots, m\} \mid f_i(x^*) = 0\}$ .

If  $A$  were known, (P) would reduce to an equality-constrained optimisation.

Active set methods guess and update  $A$  (the current guess is called "working set"  $W$ )

Alg (active set method): Given  $f_0, \dots, f_m, g_1, \dots, g_p, W \subset \{1, \dots, m\}$

repeat

1) minimise  $f_0(x)$  s.t.  $f_i(x) = g_j(x) = 0 \quad \forall j \in \{1, \dots, p\}, i \in W \quad (P_W)$   
if  $f_i(x) > 0$  for some  $i \notin W$ , add  $i$  to  $W$  and repeat step 1)  
*constraint was violated*

2) update working set:  $W = \{j \in \{1, \dots, m\} \mid f_j = 0 \text{ and } \mu_j \geq 0\}$   
*constraint is active*

until  $W$  does not change

*Lagrange multiplier for constraint  $f_j = 0$*

# Optimization algorithms: Noninterior methods for ineq.-constrained optim.

## Active set methods: convergence

Rem: Step 1) can be performed via any method for equality-constrained optimisation.

• When  $W$  does not change, then for some  $\mu \in \mathbb{R}^m$ ,  $\lambda \in \mathbb{R}^p$  we have

$$0 = Df_0(x) + \sum_{i=1}^m \mu_i Df_i(x) + \sum_{i=1}^p \lambda_i Dg_i(x) \quad , \quad g_1(x) = \dots = g_p(x) = 0 \quad ,$$

$$f_i(x) < 0 \quad , \quad \mu_i = 0 \quad , \quad i \notin W \quad , \quad f_i(x) = 0 \quad , \quad \mu_i \geq 0 \quad , \quad i \in W \quad ,$$

i.e.  $W=A$  and  $x$  satisfies the KKT conditions.

Thm: Suppose, for every  $W \subset \{1, \dots, m\}$  the problem  $(P_W)$  has a unique nondegenerate solution (i.e.  $\mu_i \neq 0 \forall i \in W$ ). Then the active set method converges to the optimal point of  $(P)$

proof: After the solution for one working set  $W$  is found, a decrease in the objective is made  $\Rightarrow$  it is impossible to return to that working set.

Process must terminate with  $W=A$  since there are only finitely many working sets.  $\square$

Rem: • Many solves with incorrect  $W$  needed.

- To update  $W$ , in principle,  $(P_W)$  needs to be solved exactly. *could have infinitely many changes of  $W$*
- In practice, one updates  $(P_W)$  prematurely (based on heuristics), but then convergence *unclear*.

# Optimization algorithms: Primal-dual methods

## Local convexity

$$f_0, \dots, f_m, g_1, \dots, g_r \in C^2$$

$$\min_{x \in \mathbb{R}^n} f_0(x) \quad \text{s.t.} \quad c_E(x) = (g_1, \dots, g_r)^T(x) = 0, \quad c_I(x) = (f_1, \dots, f_m)^T(x) \leq 0 \quad (P)$$

Let  $x^*$  be a local minimum of (P) & let it be a regular point wrt. the constraints.

$$\bullet \exists \lambda^* \in \mathbb{R}^r, \mu^* \in [0, \infty)^m : \quad Df_0(x^*) + \lambda^{*T} Dc_E(x^*) + \mu^{*T} Dc_I(x^*) = 0, \quad \mu_i^* f_i(x^*) = 0, \text{ and}$$

$$\bullet L_{xx}(x^*, \mu^*, \lambda^*) = D^2 f_0(x^*) + \sum_{i=1}^r \lambda_i^* D^2 g_i(x^*) + \sum_{i=1}^m \mu_i^* D^2 c_i(x^*) \text{ is positive semi-definite on the tangent space } T_{x^*} S \text{ to the active constraints (i.e. } c_E(x) = 0 \text{ and } f_i(x) \leq 0 \text{ for } i \in J = \{i \in \{1, \dots, m\} \mid f_i(x^*) = 0\})$$

$$\underline{\text{Def}}: \text{ We call } L_{xx}(x^*, \mu^*, \lambda^*) \text{ is positive definite on } \mathbb{R}^h \quad (C)$$

the local convexity assumption.

Thm: Under (C),  $x^*$  is a local minimum of the unconstrained problem  $\min_{x \in \mathbb{R}^n} f_0(x) + \lambda^{*T} c_E(x) + \mu^{*T} c_I(x)$ .

Also, for any  $(\mu, \lambda)$  in a neighbourhood of  $(\mu^*, \lambda^*)$  with  $\mu_i = 0$  for  $i \notin J$ ,  $\underbrace{f_0(x) + \lambda^T c_E(x) + \mu^T c_I(x)}_{L(x, \mu, \lambda)}$  has a local minimum  $x$  near  $x^*$ .

proof:  $x^*$  satisfies the second order sufficient conditions for optimality

$$\bullet L_{xx}(x^*, \mu^*, \lambda^*) \text{ pos. def. } \xrightarrow{\text{implicit function}} L_x(x, \mu, \lambda) = 0 \text{ has local solution } x(\mu, \lambda);$$

also, locally  $L_{xx}(x, \mu, \lambda)$  pos. def.  $\Rightarrow$  second order sufficient conditions for optimality fulfilled  $\square$

## Local duality

(becomes global duality for convex problems)

In the following we will assume all constraints to be active (else we have to ignore inactive constraints)

Def: The (local) dual function is defined for  $(\mu, \lambda)$  near  $(\mu^*, \lambda^*)$  as  $g(\mu, \lambda) = \min_{x \text{ near } x^*} L(x, \mu, \lambda)$ .

Under (C) this is well-defined. The minimiser is denoted  $x(\mu, \lambda)$ .

Thm:  $Dg(\mu, \lambda) = (c_I(x(\mu, \lambda))^T \quad c_E(x(\mu, \lambda))^T)$

$$D^2g(\mu, \lambda) = - \begin{pmatrix} D c_I \\ D c_E \end{pmatrix} L_{xx}(x(\mu, \lambda), \mu, \lambda)^{-1} (\nabla c_I \quad \nabla c_E) \quad \text{evaluated at } x = x(\mu, \lambda).$$

proof: Abbreviate  $\bar{x} = x(\mu, \lambda)$ , then  $g(\mu, \lambda) = f_0(\bar{x}) + \mu^T c_I(\bar{x}) + \lambda^T c_E(\bar{x}) = L(\bar{x}, \mu, \lambda)$

$$\Rightarrow \nabla g(\mu, \lambda) = \underbrace{L_x(\bar{x}, \mu, \lambda)}_{=0} D_{(\mu, \lambda)} \bar{x} + (c_I(\bar{x})^T \quad c_E(\bar{x})^T) = (c_I(\bar{x})^T \quad c_E(\bar{x})^T)$$

$$\Rightarrow D^2g(\mu, \lambda) = \begin{pmatrix} D c_I(\bar{x}) \\ D c_E(\bar{x}) \end{pmatrix} D_{(\mu, \lambda)} \bar{x}$$

← solve for  $D_{(\mu, \lambda)} \bar{x}$  and plug in

Now differentiating  $0 = L_x(\bar{x}, \mu, \lambda)$  w.r.t  $(\mu, \lambda)$ ,  $0 = L_{xx}(\bar{x}, \mu, \lambda) D_{(\mu, \lambda)} \bar{x} + (\nabla c_I(\bar{x}) \quad \nabla c_E(\bar{x}))$   $\square$

Thm (local duality): Let  $x^*$  be regular and locally optimal for (P) with Lagrange multipliers  $(\mu^*, \lambda^*)$ .

Under (C), the dual problem  $\max_{(\mu, \lambda) \in \mathbb{R}^{m+p}} g(\mu, \lambda)$  has the local solution  $(\mu^*, \lambda^*)$  with  $g(\mu^*, \lambda^*) = f_0(x^*)$ . strong duality

proof: By the above,  $x(\mu^*, \lambda^*) = x^*$  and  $\nabla g(\mu^*, \lambda^*) = (c_I(x^*)^T \quad c_E(x^*)^T) = 0$  &  $D^2g(\mu^*, \lambda^*)$  is neg. def.

$\Rightarrow (\mu^*, \lambda^*)$  is local maximum of  $g$  with  $g(\mu^*, \lambda^*) = \min_x f_0(x) + \mu^{*T} c_I(x) + \lambda^{*T} c_E(x) = f_0(x^*)$ .  $\square$

# Optimization algorithms: Primal-dual methods

## Dual problems and penalty functions

• Steepest descent convergence rate for (P) depends on condition number  $\kappa$  of  $L_{xx}(x^*, \mu^*, \lambda^*)$ , restricted to the tangent subspace  $T_{x^*} S$  to the active constraints.

• Steepest ascent convergence rate for dual problem depends on  $\kappa(D^2 g(\mu^*, \lambda^*))$  with  $D^2 g(\mu^*, \lambda^*) = - \begin{pmatrix} DC_E \\ DC_E \end{pmatrix}^{-1} L_{xx} \begin{pmatrix} DC_E \\ DC_E \end{pmatrix}$  at  $(x^*, \mu^*, \lambda^*)$ , a restriction of  $L_{xx}^{-1}$  to  $T_{x^*} S^\perp$ !

Problem:  $L_{xx}(x^*, \mu^*, \lambda^*)$  is in general not pos. def.  $\Rightarrow$  local duality not applicable!

• Steepest descent convergence rate of penalty method subproblem depends on condition number of  $D^2 q(c_k, x_k)$ , e.g. for  $P(x_k) = \frac{1}{2} \|C_E(x_k)\|_2^2 + \frac{1}{2} \|C_I^+(x_k)\|_2^2$

$$D^2 q(c_k, x_k) = D^2 f_0(x_k) + c_k \left[ \sum_{i=1}^p g_i(x_k) D^2 g_i(x_k) + \sum_{i=1}^m f_i^+(x_k) D^2 f_i^+(x_k) + (\nabla_{C_I^+}(x_k) \nabla_{C_E}(x_k)) \begin{pmatrix} DC_I^+(x_k) \\ DC_E(x_k) \end{pmatrix} \right]$$
$$= L_{xx}(x_k, \mu_k, \lambda_k) + c_k (\nabla_{C_I^+}(x_k) \nabla_{C_E}(x_k)) \begin{pmatrix} DC_I^+(x_k) \\ DC_E(x_k) \end{pmatrix} \quad \text{for } (\mu_k, \lambda_k) = c_k (C_I^+(x_k), C_E(x_k))$$

Problem: As  $c_k \rightarrow \infty$ , the condition number gets arbitrarily bad

(largest eigenvalue grows like  $c_k$ , smallest stays roughly same).

If Newton descent is used, then instead the efficiency of solving Newton's equations depends on condition number!

## Augmented Lagrangian method: Motivation by duality

$$\min_{x \in \mathbb{R}^n} f_0(x) \quad \text{s.t.} \quad c_E(x) = 0 \quad \Leftrightarrow \quad \min_{x \in \mathbb{R}^n} f_0(x) + \frac{c}{2} \|c_E(x)\|_2^2 \quad \text{s.t.} \quad c_E(x) = 0$$

*for any  $c \in \mathbb{R}^!$*

*Lagrangian of original formulation*

- Lagrangian of new formulation is  $L(x, \lambda) + \frac{c}{2} \|c_E(x)\|_2^2$ .
- Same first order optimality condition and Lagrange multiplier  $\lambda^*$ , but Hessian at the optimum is  $L_{xx}(x^*, \lambda^*) + c \nabla c_E(x^*) \nabla c_E(x^*)^T$ , pos. def. for  $c$  large enough!  
 $\Rightarrow$  dual method now applicable!
- Dual function  $g(\lambda) = \min_x f_0(x) + \lambda^T c_E(x) + \frac{c}{2} \|c_E(x)\|_2^2$  has at optimum the Hessian  $D^2 g(\lambda^*) = -D c_E(x^*) \left[ D^2 f_0(x^*) + \sum_{i=1}^r \lambda_i D^2 g_i(x^*) + c \nabla c_E(x^*) \nabla c_E(x^*)^T \right]^{-1} \nabla c_E(x^*)$   
 $\Rightarrow D^2 g(\lambda^*)$  approaches  $-\frac{1}{c} I$  for  $c \rightarrow \infty$ , which has condition number 1  
 $\Rightarrow$  extremely fast convergence of dual ascent / very efficient Newton method
- Apply modified Newton's method to dual problem, using approximate Hessian  $-\frac{1}{c} I$   
 $\Rightarrow \lambda_{k+1} = \lambda_k + c \nabla g(\lambda_k) = \lambda_k + c c_E(x_k)^T$  *can also be viewed as gradient ascent with stepsize  $c$*   
 where  $x_k$  minimises  $f_0(x) + \lambda_k^T c_E(x) + \frac{c}{2} \|c_E(x_k)\|_2^2$

# Optimization algorithms: Primal-dual methods

## Augmented Lagrangian method: Motivation by penalty

$$\min_{x \in \mathbb{R}^n} f_0(x) \text{ s.t. } c_E(x) = 0 \quad \Leftrightarrow \quad \min_{x \in \mathbb{R}^n} f_0(x) + \lambda^T c_E(x) \quad \text{s.t. } c_E(x) = 0$$

for any  $\lambda \in \mathbb{R}^p$

• penalty method objective  $q(c, x) = f_0(x) + \lambda^T c_E(x) + \frac{c}{2} \|c_E(x)\|_2^2$

• if  $\lambda = \lambda^*$  for the Lagrange multiplier of the original formulation, then the gradient of the penalty objective,  $\nabla f_0(x^*) + \lambda^{*T} \nabla c_E(x^*) + c c_E(x^*)^T \nabla c_E(x^*)$ , is zero

$\Rightarrow$  penalty method is exact even for finite  $c$ !  $\Rightarrow$  Hessian better conditioned!

• optimal point satisfies  $0 = \nabla f_0(x^*) + \lambda^{*T} \nabla c_E(x^*)$

penalty method iterate satisfies  $0 = \nabla f_0(x_k) + (\lambda_k + c c_E(x_k))^T \nabla c_E(x_k)$

$\Rightarrow$  choose  $\lambda_{k+1} \approx \lambda^* \approx \lambda_k + c c_E(x_k)$

## Augmented Lagrangian method: algorithm

Def:  $L_c(x, \lambda) = f_0(x) + \lambda^T c_E(x) + \frac{c}{2} \|c_E(x)\|_2^2$  is called the augmented Lagrangian to (P)

Alg: given  $f_0, c_E, x_0, \lambda_0$ , monotonically increasing sequence  $c_0, c_1, \dots, c_k = 0$

repeat  $x_{k+1} = \underset{x \in \mathbb{R}^n}{\operatorname{argmin}} L_{c_k}(x, \lambda_k)$

$$\lambda_{k+1} = \lambda_k + c_k c_E(x_{k+1})$$

$$c_k \leftarrow c_{k+1}$$

until  $c_E(x_{k+1}) = 0$

Thm: Let the 2<sup>nd</sup> order sufficient conditions for local optimality of (P) be satisfied at  $(x^*, \lambda^*)$ .

There is a  $c^* \in \mathbb{R}$  such that  $L_c(x, \lambda^*)$  has a local minimum at  $x^*$  for all  $c > c^*$ .

proof: • 1<sup>st</sup> order optimality satisfied at  $x^*$ , only need to check positive definiteness of  $D_x^2 L_c$

•  $D_x^2 L_c(x^*, \lambda^*) = L_{xx}(x^*, \lambda^*) + c D c_E(x^*)^T D c_E(x^*) = A + c B$  A pos. def. on  $T_{x^*} S = \ker B$   
B pos. def. on  $T_{x^*} S^\perp = \operatorname{span}(\nabla c_E(x^*))$

• suppose,  $\forall k \in \mathbb{N} \exists p_k: \|p_k\|_2 = 1, p_k^T (A + k B) p_k \leq 0$ , then for a subsequence,  $p_k \rightarrow p$

•  $0 \geq p_k^T B p_k \rightarrow p^T B p \Rightarrow p^T B p = 0 \Rightarrow p^T A p \leq 0$ , but this contradicts  $p \in \ker B$   $\square$

Rem: By continuity, the above also holds in a neighbourhood of  $(x^*, \lambda^*) \Rightarrow$  method well-defined

## Augmented Lagrangian method: example

Ex:  $\min_{(x,y) \in \mathbb{R}^2} 2x^2 + 2xy + y^2 - 2y \quad \text{s.t.} \quad x=0$

•  $L_c(x,y,\lambda) = 2x^2 + 2xy + y^2 - 2y + \lambda x + \frac{c}{2}x^2$

•  $(x_{k+1}, y_{k+1}) = \underset{(x,y)}{\operatorname{argmin}} L_{c_k}(x,y,\lambda_k) = \left( -\frac{2+\lambda_k}{2+c_k}, \frac{4+c_k+\lambda_k}{2+c_k} \right)$

$\lambda_{k+1} = \lambda_k + c_k x_{k+1} = \frac{2}{2+c_k} \lambda_k - \frac{2c_k}{2+c_k}$

•  $\lambda_k \xrightarrow{k \rightarrow \infty} -2$  with  $(\lambda_{k+1} + 2) = \frac{2}{2+c_k} (\lambda_k + 2) \Rightarrow$  linear convergence at rate  $\frac{2}{2+c}$

Ex:  $\min_{(x,y) \in \mathbb{R}^2} 2x^2 + 2xy + y^2 - 2y \quad \text{s.t.} \quad x=0, y \leq 0$

• replace by  $\min_{(x,y,z) \in \mathbb{R}^2} 2x^2 + 2xy + y^2 - 2y \quad \text{s.t.} \quad x=0, y+z^2=0$

•  $L_c(x,y,z,\lambda,\nu) = 2x^2 + 2xy + y^2 - 2y + \lambda x + \nu(y+z^2) + \frac{c}{2}x^2 + \frac{c}{2}(y+z^2)^2$

•  $(x,y,z)_{k+1} = \left( \frac{2\nu_k - 4 - 2\lambda_k - c_k \lambda_k}{(2+c_k)(4+c_k)} - \frac{2-\nu_k}{2+c_k} + \frac{2\lambda_k - (4\nu_k - 8)/(2+c_k)}{(2+c_k)(4+c_k)} - 4, 0 \right)$

$(\lambda, \nu)_{k+1} = (\lambda_k, \nu_k) + c_k (x_{k+1}, y_{k+1})$

•  $(\lambda_k, \nu_k) \xrightarrow{k \rightarrow \infty} (0, 2)$

(linear convergence for fixed  $c_k$ )

## Augmented Lagrangian method: convergence rate

Recall: update  $\lambda_{k+1} = \lambda_k + \overset{\text{assume fixed } c}{c} c_E(x_k) = \lambda_k + c \nabla g(\lambda_k)$  is gradient ascent

for dual function  $g$  to  $\min_{x \in \mathbb{R}^n} f_0(x) + \frac{c}{2} \|c_E(x)\|_2^2 \quad \text{s.t. } c_E(x) = 0$

$\Rightarrow$  convergence rate depends on eigenvalues of


$$D^2 g(\lambda^*) = -\underbrace{D c_E(x^*)}_{B} \left[ \underbrace{L_{xx}(x^*, \lambda^*)}_{A} + c \nabla c_E(x^*) \nabla c_E(x^*)^T \right]^{-1} \nabla c_E(x^*) = -B(A + cB^T B)^{-1} B^T$$

Thm: Let  $m$  be an eigenvalue of  $BA^{-1}B^T$ , then  $\frac{m}{1+cm}$  is an eigenvalue of  $B(A + cB^T B)^{-1} B^T$ .  
 $\leftarrow$  Hessian of dual problem to (P)

proof: By Sherman-Morrison-Woodbury,  $B(A + cB^T B)^{-1} B^T = F - c F(I + cF)^{-1} F$  for  $F = BA^{-1}B^T$

• let  $Fv = mv$ , then  $B(A + cB^T B)^{-1} B^T v = \left( m - c \frac{m^2}{1+cm} \right) v = \frac{m}{1+cm} v \quad \square$

Rem: The convergence rate of steepest ascent for the dual problem to (P) depends on the condition number  $\kappa(BA^{-1}B^T) = \frac{M}{m}$  for smallest / largest eigenvalue  $m/M$  of  $BA^{-1}B^T$ .



- Convergence rate of augmented Lagrangian method depends on  $\kappa(B(A + cB^T B)^{-1} B^T) = \frac{c + \frac{1}{m}}{c + \frac{1}{M}}$
- One can improve on the rate by using a quasi-Newton or Newton method for  $g$ .