

Table S1: Summary of studies comparing the treatment efficiency between conventional brackets and self-ligating brackets

Author	Study design	Number of Patients	Participants	Age (years)	Measurement	Treatment duration (Mean $\pm$ SD)			Result of statistical tests (p-values)
Pandis et al. 2007	CCT	n=54	• G1: CB, 0.022'' (Microarch) (n= 27) • G2: SB, 0.022'' (Damon2) (n= 27)	Mean age (SD): G1: 13.92 (1.43) G2: 13.48 (1.31)	• Rate of alignment	• G1: 114.51$\pm$46.44 days • G2: 91.03$\pm$31.94 days			No significant difference (p=0.06)
Scott et al. 2008	RCT	n=60	• G1: CB (Synthesis) (n= 28) (9 F, 19 M) • G2: SB (Damon3) (n= 32) (20 F, 12 M)	Mean age: G1: 16.38 G2: 16.19	• Rate of alignment	• G1: 243$\pm$82.5 days • G2: 253$\pm$63.6 days			No significant difference
Fleming et al. 2010	RCT	n=54	• G1: CB (Victory) (n= 26) (19 F, 7 M) • G2: SB (SmartClip) (n= 28) (17 F, 11 M)	Mean age (SD): G1: 15.48 (2.38) G2: 16.11 (2.74)	• Treatment duration • Total number of visits	• G1: 18.32$\pm$4.66 months • G2: 21.41$\pm$8.44 months			No significant difference for treatment duration (p=0.076), no sign. difference in number of visits
DiBiase et al. 2011	RCT	n=48	• G1: CB (Synthesis) (n= 21) (9 F, 20 M) • G2: SB (Damon3) (n= 27) (21 F, 12 M)	Mean age (SD): G1: 16.38 (5.28) G2: 16.19 (3.68)	• Treatment duration • Number of visits • Occlusal outcome	• G1: 23.00$\pm$4.86 months • G2: 24.48$\pm$6.72 months			No significant difference for treatment duration (p=0.992), No significant difference for no of visits
Johansson and Lundstrom 2012	RCT	n=90	• G1: CB (Gemini) (n= 46) (33 F, 13 M) • G2: SB (Time2) (n= 44) (31 F, 13 M)	Age range: 11.7 to 18.2	• Overall treatment time • Number of visits	• G1: 18.2$\pm$6.56 months • G2: 20.4$\pm$5.97 months			No significant difference for treatment duration and no. of visits
Songra et al. 2014	RCT	n=98	• G1: CB, 0.022'' (Omni) (n= 20) • G2: passive SB, 0.022'' (Damon 3MX) (n=41) • G3: active SB, 0.022'' (InOvation R) (n=37)	Age range:	• Labial segment alignment • Space closure	• G1: 251$\pm$107 days • G2: 422$\pm$124 days • G3: 399$\pm$107 days			No significant differences
Wahab et al. 2014	RCT	n=29	• G1: CB (Ormco) (n= 15) (10 F, 5 M) • G2: SB (Damon3) (n= 14) (11 F, 3 M)	Mean age: G1: 19.5 $\pm$ 3.9 G2: 21.9 $\pm$ 3.6	• Rate of alignment • Crowding alleviation		G1 (mm)	G2 (mm)	Differences in treatment time: T0-T1: p = 0.085 T1-T2: p < 0.05 T2-T3: p = 0.326 T3-T4: p = 0.793
						T0-T1	4.95 $\pm$ 2.58	1.94 $\pm$ 3.07	
						T1-T2	5.35 $\pm$ 2.27	2.13 $\pm$ 3.01	

Author	Study design	Number of Patients	Participants	Age (years)	Measurement	Treatment duration (Mean $\pm$ SD)			Result of statistical tests (p-values)
						T2-T3	1.57 $\pm$ 1.22	0.94 $\pm$ 1.31	Crowding alleviation showed a higher percentage with G2 (98%) compared with G1 (67%).
						T3-T4	1.14 $\pm$ 2.39	0.99 $\pm$ 0.65	
O'Dwyer et al. 2016	RCT	n=137	• G1: CB (Victory) (n=71) (42 F, 29 M) • G2: SB (SmartClip) (n=66) (43 F, 23 M)	Mean: G1: 14 years 6 months G2: 15 years 6 months	Treatment duration	• G1: 25.80 (months) • G2: 25.12 (months)			No sign. differences with regard to the number of visits and overall treatment time
Jung 2021	CCT	n=134	• G1: CB, 0.022'' (Clarity) (n=68) • G2: SB, 0.022'' (Clippy) (n=66)	Average age: 22.73	Treatment duration	• G1: 29.25$\pm$7.53 (months) • G2: 27.98$\pm$7.10 (months)			No significant difference

abbreviations: CB – Conventional bracket; CCT – Controlled clinical trial; F – Female; G1 – Group 1; G2 – Group 2; G3 – Group 3; M – Male; mm – millimeter; n – number of patients; NR – not reported, RCT – Randomized controlled trial; SB – Self-ligating bracket; SD – Standard Deviation; T0 – Pretreatment; T1 – first month review; T2 – Second month review; T3 – Third month review; T4 – Fourth month review

Table S2: Summary of studies comparing the rate of root resorption between conventional brackets and self-ligating brackets

Author	Study design	No. of Patients	Groups	Age (years) Mean $\pm$ SD	Evaluated Teeth	Assessment method	Duration of treatment	Root resorption (Mean $\pm$ SD)	Outcome of significance test
Scott et al. 2008	RCT	n=60	- G1: CB, 0.022'' (Synthesis) (n=28) (9 F, 19 M) - G2: SB, 0.022'' (Damon3) (n=32) (20 F, 12 M)	G1: 16.38 $\pm$ 5.28 G2: 16.19 $\pm$ 3.68	Mn right central incisor	PA	G1: 8.1 $\pm$ 2.7 months G2: 8.5 $\pm$ 2.1 months	EARR (mm) - G1: 1.21 $\pm$3.39 - G2: 2.26 $\pm$2.63	No significant difference
Pandis et al. 2008	CCT	n=96	- G1: CB, 0.022'' (Microarch) (n=48) (36 F, 12 M) - G2: SB, 0.022'' (Damon 2) (n=48) (31 F, 17 M)	G1: 13.14 $\pm$ 1.73 G2: 13.29 $\pm$ 1.57	- Mx incisors - Mn incisors	OPG	G1: 25.97 $\pm$ 6.65 months G2: 26.89 $\pm$ 5.94 months	EARR (mm) - Mx incisors: 1.23 $\pm$0.97 - Mn incisors: 1.36 $\pm$0.90	No significant difference
Leite et al. 2012	RCT	n=19	- G1: CB, 0.022'' (3M Unitek) (n = 8) (6 F, 2 M) - G2: SB, 0.022'' (Easyclip) (n = 11) (5 F, 6 M)	Mean: 20.6 Range: 11–30	- Mx central incisor - Mx lateral incisor - Mn central incisor - Mn lateral incisor	CBCT	6 months	EARR (mm) - Mx central incisor: G1: 0.33 $\pm$ 0.19 G2: 0.34 $\pm$ 0.24 - Mx lateral incisor: G1: 0.44 $\pm$ 0.33 G2: 0.43 $\pm$ 0.33 - Mn central incisor: G1: 0.31 $\pm$ 0.21 G2: 0.39 $\pm$ 0.52 - Mn lateral incisor: G1: 0.40 $\pm$ 0.24 G2: 0.23 $\pm$ 0.23	No significant difference
Jacobs et al. 2014	CCT	n=213	- G1: CB, 0.022'' (Victory) (n= 74) (51 F, 23 M) - G2: SB, 0.022'' (Smart clip)	Mean: 12.4 $\pm$ 2.2	- Mx incisors - Mn incisors	OPG	G1: 18.1 $\pm$ 5.3 months G2: 20.7 $\pm$ 4.9 months	rRR (%) G1: 4.5 $\pm$6.6; G2: 3.0 $\pm$5.6	No significant difference

Author	Study design	No. of Patients	Groups	Age (years) Mean $\pm$ SD	Evaluated Teeth	Assessment method	Duration of treatment	Root resorption (Mean $\pm$ SD)	Outcome of significance test
			(n=139) (83 F, 56 M)						
Chen et al. 2015	CCT	n=70	- G1: CB, 0.022'' (3M Unitek) (n= 35) (19 F, 16 M) - G2: SB, 0.022'' (Damon 3) (n= 35) (18 F, 17 M)	G1: 13.42 $\pm$ 2.50 G2: 13.52 $\pm$ 2.84	- Mx central incisor - Mx lateral incisor - Mn central incisor - Mn lateral incisor	PA	Mean $\pm$ SD: G1: 20.34 $\pm$ 3.40 months G2: 20.53 $\pm$ 3.62 months	EARR (mm) - Mx central incisor: G1: 0.5 $\pm$0.3; G2: 0.3 $\pm$0.4 - Mx lateral incisor: G1: 0.3 $\pm$0.5; G2: 0.2 $\pm$0.3 - Mn central incisor: G1: 0.4 $\pm$0.5; G2: 0.4 $\pm$0.4 - Mn lateral incisor: G1: 0.3 $\pm$0.5; G2: 0.3 $\pm$0.3	No significant differences
Handem et al. 2016	CCT	n=52	- G1: CB, 0.022'' (Roth) (n= 27) (13 F, 14 M) - G2: SB, 0.022'' (Damon) (n= 25) (12 F, 13 M)	Mean age range: G1: 16.77-18.47 G2: 16.04-18.06	- Mx central incisor - Mx lateral incisor - Mn central incisor - Mn lateral incisor	PA	Mean: G1: 1.70 years G2: 2.02 years	EARR (mm) - Mx right central incisor: G1: 0.59; G2: 0.72 - Mx right lateral incisor: G1: 0.70; G2: 0.72 - Mx left central incisor: G1: 0.66; G2: 0.88 - Mx left lateral incisor: G1: 0.74; G2: 0.80 - Mn right central incisor: G1: 0.66; G2: 0.64 - Mn right lateral incisor: G1: 0.40; G2: 0.48 - Mn left central incisor: G1: 0.62; G2: 0.60 - Mn left lateral incisor: G1: 0.55; G2: 0.56	No significant difference
Aras et al. 2018	RCT	n=32	G1: CB, 0.022'' (Titanium Orthos) (n= 16) (10 F, 6 M)	Average: G1: 14.94 $\pm$ 1.06	- Central incisors - Lateral incisors	CBCT	9 months	Volumetric changes (mm ³) - Central incisor: - G1: 28.29 $\pm$13.48 - G2: 27.08 $\pm$12.71 - Lateral incisor:	No significant difference

Author	Study design	No. of Patients	Groups	Age (years) Mean $\pm$ SD	Evaluated Teeth	Assessment method	Duration of treatment	Root resorption (Mean $\pm$ SD)	Outcome of significance test
			G2: SB, 0.022'' (Damon Q) (n= 16) (12 F, 4 M)	G2: 15.00 $\pm$ 1.03				- G1: 18.77 $\pm$11.05 - G2: 20.32 $\pm$11.67	
Qin and Zhou 2019	RCT	n=98	- G1: CB, 0.019'' (3M Unitek) (n= 49) (23 F, 26 M) - G2: SB, 0.019'' (Damon 3) (n= 49) (24 F, 25 M)	G1: 15.15 $\pm$ 4.52 G2: 15.21 $\pm$ 4.43	- Mx central incisors - Mx lateral incisors	OPG	M $\pm$ SD: G1: 20.25 $\pm$ 5.11 months G2: 22.10 $\pm$ 5.15 months	EARR (mm) - Mx right central incisor: G1: 0.40 $\pm$0.28 G2: 0.32 $\pm$0.24 - Mx right lateral incisor: G1: 0.30 $\pm$0.25 G2: 0.27 $\pm$0.28 - Mx left central incisor: G1: 0.39 $\pm$0.31 G2: 0.33 $\pm$0.27 - Mx left lateral incisor: G1: 0.31 $\pm$0.26 G2: 0.28 $\pm$0.25	No statistically significant difference

abbreviations: CB – Conventional bracket; CBCT – Cone Beam Computed Tomography; CCT – Controlled clinical trial; EARR – External apical root resorption; F – Female; G1 – Group 1; G2 – Group 2; M – Male; Mn – Mandibular; Mx – Maxillary; mm – millimeter; n – number of patients; OPG – Orthopantomography; PA – Periapical Radiograph; RCT – Randomized controlled trial; rRR – relative root resorption; SB – Self-ligating bracket; SD – Standard Deviation;

Table S3: Summary of studies comparing the level of pain between conventional brackets and self-ligating brackets

Author	Study design	No. of Patients	Participants	Age (years)	Assessment method	Pain level			Outcome of significance tests
Miles et al. 2006	CCT, split-mouth	n=58 (40 F, 18 M)	G1: - Mn right quadrant: SB, (Damon2) - Left quadrant: CB, (Victory) (n=29) G2: - Mn right quadrant: CB, (Victory) - Left quadrant: SB (Damon2) (n=29)	Average age: 16.3 (10.5 – 46.5)	VRS (Comparing left and right, which side is more painful)		SB n (%)	CB n (%)	1-2 d: p = .04 untying: p = .56 tying: p = .004
						days 1-2	2 (3.4)	9 (15.5)	
						untying	7 (12.1)	5 (8.6)	
						tying	21 (36.2)	6 (10.3)	
Scott et al. 2008	RCT	n=62	- G1: CB (Synthesis) (n=29) (9 F, 20 M) - G2: SB (Damon3) (n=33) (21 F, 12 M)	Mean age: 16 years 3 months	VAS (0-100mm) 4 hours 24 hours 3 days 1 week		G1	G2	No significant difference
						4 hours	60.11±23.3	63.67±23.18	
						24 hours	57.44±24.24	58.3±20.74	
						3 days	38.19±26.43	42.94±23.08	
						1 week	15.11±20.32	15.12±14.32	
Pringle et al. 2009	RCT	n=52	- G1: CB (Tru Straight) (n=28) (13 F, 15 M) - G2: SB (Damon3) (n=24) (15 F, 9 M)	Mean age (SD): G1: 16.1 (7.4) G2: 15.2 (6.8)	VAS (0-10cm) Twice a day, over 7 days	Mean MPI: - G1: 55.71 mm (95% CI, 45.93-65.50 mm) - G2: 40.92 mm (95% CI, 30.40-51.44 mm) - G2 reported lower mean MPI (difference, 11.77 mm; 95% CI, -0.15-23.68 mm) than did the G1			- Mean maximum pain intensity: P = 0.053 - Mean pain intensity: P = 0.012
Fleming et al. 2009	RCT	n=48	- G1: CB (Victory) (n=22) (16 F, 6 M) - G2: SB (SmartClip) (n=26) (16 F, 10 M)	Mean age: G1: 15.65±2.1 G2: 16.23±2.91	VAS (0-100 mm) 4 hours 24 hours 3 days 7 days - Archwire removal - Archwire insertion		G1	G2	at 4 hours (p = .958), 24 hours (p = .289), 72 hours (p = .569), and 7 days (p = .756) wire removal (p = .001), wire insertion (p = .013)
						4 hours	46.43±24.07	46.81±24.41	
						24 hours	67.7±17.83	61.46±25.35	

Author	Study design	No. of Patients	Participants	Age (years)	Assessment method	Pain level				Outcome of significance tests
						3 days	46.45±24.15	42.58±23.87		
						7 days	18.52±18.31	17.08±16.47		
						Archwire removal	15.52±14.3	33.65±21.8		
						Archwire insertion	19.74±17.53	38.02±27.47		
Tecco et al. 2009	CCT	n=30	• G1: CB (Victory) (n=15) (8 F, 7 M) • G2: SB (Damon SL II) (n=15) (10 F, 5 M)	Mean age: G1: 16.9 ± 3.8 G2: 16.4 ± 4.5	VAS (0-100mm) Each evening from placement of the first archwire for 3 months		G1 mean (SD)	G2 mean (SD)	Significant differences: • Day 1 (p<0.05) • Day 3 (p<0.05)	
						Day 1	52 (45.07)	42 (20.25)		
						Day 3	50.3 (34.93)	18.83 (16.50)		
Miles and Weyant 2010	RCT	n=60	• G1: CB (Clarity) (n=30) (19 F, 11 M) • G2: SB, (InOvation C) (n=30) (19 F, 11 M)	Mean age: 13.5+1.5	7-point Likert scale • 4 hours • 24 hours • 3 days • 1 week	Time	G1 Mean (SD)	G2 Mean (SD)	P-Value showed no significant differences • 4 hours: 0.42 • 1-day: 0.33 • 3-days: 0.87 • 7-days: 0.29 • Total: 0.90	
						4 hours	2.4 (1.1)	2.8 (1.5)		
						1-day	3.9 (1.7)	3.4 (1.8)		
						3-days	2.4 (0.9)	2.5 (1.5)		
						7-days	1.3 (0.6)	1.6 (1.1)		
						Total	10.1 (3.2)	10.2 (5.2)		
Bertl et al. 2013	CCT, split-mouth	n=18 (15 F, 3 M)	SB: (SmartClip) CB: (AO) • G1: SB on left side and CB on the right (n=11) • G2: SB on right side and CB on the left (n=7)	Mean age: 22.2±6.4	NRS (0 – 10) • Archwire engagement • Archwire disengagement	NRS Scores (SB minus CB)				• Treatment: not significant (p = 0.21) • archwire disengagements (p = .027), SB was more painful
						Archwire Engagement		Archwire Disengagement		
						Mean (SD)	95% CI	Mean (SD)	95% CI	
						0.97 (2.25)	-0.15 (2.08)	1.04 (1.87)	0.10 (1.97)	
						1.55 (2.63)	-0.33 (3.43)	1.60 (3.27)	-0.74 (3.94)	
Atik and Ciger 2014	RCT	n=33	• G1: CB (Forestadent) (n=17, all F)	Mean age: G1: 14.5±1.2	VAS (0-100 mm) • 24 hours • 3 days	G1 (mean (CI95%))		G2		Intergroup diff: 24 h: p = 0.465 3 d: p = 0.363
						•24 h: 1.7 (0.2-2.8) •3 d: -2.1 (-4.8 – 0.1)		•24 h: 1.3 (0.3 - 2.8) •3 d: -2.1 (-5.5 – -1.1)		

Author	Study design	No. of Patients	Participants	Age (years)	Assessment method	Pain level	Outcome of significance tests
			G2: SB, (Damon3): (n=16, all F)	G2: 14.8±1.0	1 week 1 month	7 d: -1.9 (-3.9 – -0.2) 30 d: -0.8 (-1.6 - -0.1) 7 d: -1.9 (-3.9 – -0.0) 30 d: 0 (-0.8 - 0.0)	7 d: p = 0.763 30 d: p = 0.709
Rahman et al. 2016	RCT	n=113	- G1: CB (Victory) (n=60) (38 F, 22 M) - G2: SB (SmartClip) (n=53) (34 F, 19 M)	Mean: G1: 14 years 5 months G2: 15 years 4 months	VRS (none, mild, moderate, severe) - Day 1 - Day 3 - Day 5	- Pain scores - teeth > soft tissues - most pain encountered on day 1 - G2 – more discomfort (engaging/disengaging archwires)	No significant difference
Lai et al. 2019	RCT	n=88	- G1: CB (OPA-K): (n=44) (25 F, 19 M) - G2: SB (Damon Q): (n=44) (27 F, 17 M)	Median age: G1: 19.5 G2: 19.0	VAS (0-100mm) - Primary outcome: 24 h (baseline) after bracket bonding - Secondary outcome: 4 h, 3 days, 1 week and 1 month after bracket bonding	Estimate = 3.09. SE = 3.22	No significant difference (p = 0.34)
González-Sáez et al. 2021	RCT	n=90	- G1: CB (Diamond+) (n=30) (19 F, 11 M) - G2: SB (Bio-smile): (n=30) (19 F, 11 M) - G3: low-friction CB (Synergy): (n=30) (18 F, 12 M)	Mean ±SD: G1: 20.4±5.9 G2: 22.7±7.2 G3: 22.1±9.2	VAS (0-10cm) 4 hours after BB (T1) 8 hours after BB (T2) 24 hours after BB (T3) 2 days after BB (T4) 3 days after BB (T5) 4 days after BB (T6) 5 days after BB (T7) 6 days after BB (T8) 7 days after BB (T9)	- G3 being the one showing the highest pain values at time points (T1: 3.2 ± 3, T2: 3.8 ± 3.1, T7: 1.9 ± 2.3, T8: 1.7 ± 2.4, T9: 1.2 ± 2.3) - maximum peak at 24 hours in G1 (4.5 ± 2) followed by G3 (4.0 ± 2.9) and, at 48 hours, by G2 (3.4 ± 1.7)	No significant difference between G1 and G2 Only G3 showed significant difference in pain: - p < 0.01 in T1, T7 and T9 vs. G1 & G2 - p < 0.05 in T2 and T8 vs. G1 & G2

abbreviations: 95% CI – 95% confidence interval; BB _ bracket bonding; CB – Conventional bracket; CCT – Controlled clinical trial; F – Female; G1 – Group 1; G2 – Group 2; M – Male; mm – millimeter; Mn – Mandibular; MPI – Maximum pain intensities; n – number of patients; NRS – Numeric rating scale; RCT – Randomized controlled trial; SB – Self-ligating bracket; SD – Standard Deviation; SE – Standard error; VAS – Visual analog scale; VRS – Verbal rating scale

Table S4: Summary of studies comparing the periodontal status between conventional brackets and self-ligating brackets

Author	Study design	Number of Patients	Participants	Age (years)	Measured Periodontal Index				Additional comments
					PI	GI	PD	BOP	
Baka et al. 2013	CCT, Split mouth	n=20 (M)	- CB (Dentaurum) with steel ligature - SB (DamonQ) - G1: CB in the Mx right and Mn left dentitions and SB in the Mx left and Mn right dentitions (n=10) - G2: SB in the Mx right and Mn left dentitions and CB in the Mx left and Mn right dentitions (n=10)	Mean age: 14.2 ± 1.5	Yes • G1: 1.37±0.34 • G2: 1.16±0.38 • P = 0.091	No	Yes • G1: 0.68±0.32 • G2: 0.72±0.31 • P = 0.882	Yes • G1: 38.00±14.53 • G2: 37.84±15.87 • P = 0.871	
Atik and Ciger 2014	RCT	n=33 (F)	- G1: CB (Forestadent) (n=17) - G2: SB (Damon3) (n=16)	Mean age: G1: 14.5 ± 1.2 G2: 14.8 ± 1.0	Yes • G1: 0.23±0.52 • G2: 0.17±0.45 • P value: 0.746	Yes • G1: -0.08±0.62 • G2: 0.16±0.39 • P value: 0.201	Yes • G1: 0.25±0.35 • G2: 0.12±0.27 • P value: 0.286	No	
Nalcaci et al. 2014	RCT	n=46	- G1: CB (Mini Taurus) with elastomeric ligatures (n=23) (13 F, 10 M) - G2: SB (DamonQ) (n=23) (11 F, 12 M)	Mean age: G1: 13.30 ± 1.61 G2: 14.48 ± 1.27	Yes • G1: 0.60±0.06 • G2: -0.60±0.07 P = 0.967	Yes • G1: -0.63±0.67 • G2: 0.48±0.07 • P = 0.141	No	Yes • G1: -0.11±0.008 • G2: 0.11±0.11 P = 0.092	G1 with elastomeric ligatures – increase in halitosis with the accretion of plaque accumulation
Uzuner et al. 2014	RCT	n=40	- G1: CB (OPAL) with stainless steel ligature (n=20) (15 F, 5 M) - G2: SB (Leone) (n=20) (16 F, 4 M)	Mean age (SD): G1: 15.05 ± 0.97 G2: 14.67 ± 1.06	Yes Median (Min-Max) • G1: 0.93 (0.00-3.20) • G2: 1.00 (0.09-3.20)	Yes Median (Min-Max) • G1: 0.61 (0.09-2.10) • G2: 1.02 (0.12-2.70)	Yes Median (Min-Max) • G1: 2.20 (1.10-3.50) • G2: 3.00 (1.23-3.50)	No	

Author	Study design	Number of Patients	Participants	Age (years)	Measured Periodontal Index				Additional comments
					PI	GI	PD	BOP	
					• P = 0.069	• P = 0.028	• P = 0.007		
Folco et al. 2014	CCT	n=22	- G1: CB (Synthesis) with elastomeric ligatures (n=23) (13 F, 10 M) - G2: SB (Damon3) (n=11) (11 F, 12 M)	16 to 30	Yes p > 0.05	Yes p > 0.05	Yes p > 0.05	No	
Cardoso et al. 2015	CCT, Split mouth	n=16	- CB (Kirium) with metallic ligature - SB (Portia) - G1: CB on the lower dental arch and SB on the upper arch (n=8) - G2: SB in the lower arch and CB in the upper arch (n=8)	12 to 16	Yes • G1: 1.38±0.68 • G2: 1.48±0.85 • P > 0.05	No	Yes • G1: 2.01±0.53 • G2: 2.05±0.52 • P > 0.05	Yes • G1: 0.93±1.03 • G2: 0.73±0.59 • P > 0.05	
Kaygisiz et al. 2015	RCT	n=60 (28 F, 32 M)	- G1: CB (Avex) ligated with steel ligatures (n=20) - G2: SB (Leone) (n=20) - G3: control group (treatment started at the end of the study) (n=20)	Mean age (SD): G1: 14 (1.01) G2: 14.7 (1.39) G3: 14.4 (1.46)	Yes • G1: 0.78±0.52 • G2: 0.81±0.43 • G3: 0.63±0.71 • P = 0.837	Yes • G1: 0.38±0.35 • G2: 0.58±0.52 • G3: 0.47±0.51 • P = 0.346	Yes • G1: 1.42±0.30 • G2: 1.46±0.23 • G3: 1.63±0.54 • P = 0.067	Yes • G1: 2.11±0.94 • G2: 2.34±0.56 • G3: 1.97±0.73 • P = 0.286	
Chhibber et al. 2017	RCT	n=61	- G1: Clear aligner (Invisalign) (n=24) - G2: SB (Carriere) (n=17) - G3: CB (Carlsbad) with elastomeric ligatures (n=20)	Mean age: G1: 16.56 ± 3.99 G2: 15.39 ± 3.54 G3: 14.56 ± 3.92	Yes • G1: 0.92±0.58 • G2: 1.07±0.59 • G3: 1.32±0.67 • P > 0.05	Yes • G1: 0.75±0.53 • G2: 1.00±0.65 • G3: 1.32±0.67 • P > 0.05	No	Yes • G1: 0.46±0.72 • G2: 0.73±1.1 • G3: 0.84±1.01 • P > 0.05	

abbreviations: BOP – Bleeding on probing; CB – Conventional bracket; CCT – Controlled clinical trial; F – Female; G1 – Group 1; G2 – Group 2; G3 – Group 3; GI – Gingival index; M – Male; Mn – Mandibular; Mx – Maxillary; n – number of patients; PD – Probing depth; PI – Plaque index; RCT – Randomized controlled trial; SB – Self-ligating bracket; SD – Standard Deviation