

Potential Theory on Gromov Hyperbolic Manifolds of Bounded Geometry

Matthias Kemper, Joachim Lohkamp

Abstract

In the 1980s Alano Ancona developed a profound potential theory on Gromov hyperbolic manifolds of bounded geometry. Since then, such hyperbolic spaces have become basic in geometry, topology and group theory. This also stimulated a broader interest into analytic aspects of these spaces. In this paper we make Ancona's original work, addressed to a rather advanced audience, approachable for a wider readership without firm background in potential theory. We streamline the arguments to make the results depend only on coarse geometro-analytic invariants and we derive useful details not found in the original sources. Finally, we explain remarkable relations of these results to recent developments in the potential theory and quasi-conformal geometry of domains in \mathbb{R}^n .

Contents

1	Introduction	2
1.1	Goals of this paper	2
1.2	Organization of the paper	3
2	Basic Concepts	3
2.1	Geometric Structures	3
2.2	Analytic Structures	6
2.3	Potentials and Balayage	7
3	Local Maximum Principles and Harnack Inequalities	9
3.1	Maximum Principles	9
3.2	Harnack Inequalities	11
4	Global Results from Resolvent Equations and Bounded Geometry	12
4.1	Resolvent Equation	12
4.2	Behavior of Green's functions	13
4.3	Relative Maximum Principles	14
5	Hyperbolicity and Boundary Harnack Inequalities	16
5.1	Global Behavior: Φ -Chains	16
5.2	Boundary Harnack Inequality	19
5.3	The Martin Boundary	21
6	Geometry and Analysis on Uniform Domains	25
6.1	Uniform Domains	25
6.2	Quasi-Hyperbolic Geometry	27
6.3	Martin Theory for (Non-)Uniform Domains	28
	References	30

1 Introduction

In this paper we are interested in the elliptic analysis on *non-compact* complete manifolds and its relation to the geometry at infinity. Non-compact spaces occur even if one is primarily interested in compact cases, for instance as limit spaces under blow-ups or by deforming bounded domains. An example are Riemann uniformizations of planar domains which also turn the boundary of the domain into a boundary at infinity.

It is reasonable to focus on non-compact manifolds of *bounded geometry*. In simple terms, bounded geometry excludes a variety of awkward analytic phenomena resulting from degenerations towards infinity. For instance, it makes sure that basic analytic controls from local estimates, most prominently Harnack inequalities, hold uniformly all over the manifold. Moreover, a complete manifold of bounded geometry is *stochastically complete*. This means that solutions of stochastic differential equations associated to typical elliptic problems on the manifold exist for an infinitely long time.

However, having bounded geometry alone is not yet enough to gain a control as strong as in the classical case of harmonic or holomorphic functions on smoothly bounded planar domains. Finer details in the style of Cauchy's integral formula over the boundary of the given domain are reserved to the subclass of Gromov hyperbolic manifolds. For reasonable elliptic operators there is a contour integral for (positive) solutions formally taken over the so-called Martin boundary which, and this is a deep insight of Ancona's potential theory on such hyperbolic spaces, simply *equals* the natural geometric boundary at infinity, namely the Gromov boundary.

1.1 Goals of this paper

Our main goals are to explain how bounded geometry paves the way to a controllable potential theory for suitable elliptic operators and how adding Gromov hyperbolicity refines these basic controls to a remarkable degree. We will see how even mild violations of Gromov hyperbolicity can cause a breakdown of these refined controls.

This is also a tour through a variety of beautiful concepts and fundamental results from geometric hyperbolicity notions over the potential theoretic balayage and the resolvent equation to (boundary) Harnack inequalities and their associated Martin theory.

The exposition is modelled on the original sources [Anc87] and [Anc90], but we streamline and replace some of the arguments and derive a priori estimates only depending on coarse geometric and analytic data such as the hyperbolicity and coercivity constants of the considered spaces and operators. Our presentation emphasizes the central role of the boundary Harnack inequality on Gromov hyperbolic spaces and we derive some useful versions of this inequality not present in earlier literature.

Our second theme is a strategy, also initiated by Ancona [Anc87, Anc90], to transfer the potential theory on hyperbolic spaces to Euclidean domains, provided they admit conformal deformations into such a hyperbolic space. The idea is to transform the operator in such a way that the notion of a solution remains unchanged under the deformation. The effect then is that the estimates and results derived on the hyperbolic space simply remain valid under this conformal deformation.

More recent work in quasi-conformal geometry due to Bonk, Heinonen and Koskela [BHK01] has shown that so-called *uniform domains* are precisely those domains admitting the type of hyperbolizing deformations needed here. Remarkably, the particular potential theory on uniform domains obtained this way also characterizes their uniformity condition. Aikawa [Aik01, Aik04] has developed an independent analytic argument showing that, under some weak regularity assumptions, uniform domains are exactly those Euclidean domains which admit boundary Harnack inequalities for the Laplace operator. In turn, this underlines the sharpness of Ancona's results for the Gromov hyperbolic case.

1.2 Organization of the paper

Section 2 We first recall the concepts of manifolds of bounded geometry and of Gromov hyperbolicity. We also introduce the ideal boundary of such space at infinity, the Gromov boundary. Then we discuss some analytic counterparts of these geometric concepts, uniform ellipticity and coercivity of operators, which are necessary for Ancona’s theory. Finally, we introduce the basic potential theoretic notion of balayage which we use to globalize local potential theoretic considerations.

Section 3 We use the bounded geometry and uniform ellipticity to derive local estimates for Green’s functions of elliptic operators on uniformly sized balls and their relation to Harnack inequalities. The original sources [Anc87, Anc90] employ advanced analytic tools due to Stampacchia [Sta65]. We reorganize the arguments and base them on more common and elementary techniques.

Section 4 Here we additionally use the coercivity properties to formulate resolvent equations for Green’s functions considered as operators. We apply resolvent equations to derive local and, by means of chains of Harnack inequalities, also global growth estimates for Green’s functions.

Section 5 Now we also use that our manifold is Gromov hyperbolic. We employ this hyperbolicity in two places, most notably in Proposition 5.2 “growth recovery along Φ -chains” to derive our main technical result, Theorem 5.1 “Green’s functions along Φ -chains”. Secondly, the hyperbolicity readily implies the existence of an abundance of Φ -chains which allows us to deduce the hyperbolic boundary Harnack inequality 5.8 and that the so-called Martin boundary is homeomorphic to the Gromov boundary. This gives us contour integrals for positive solutions of $Lw = 0$ similar to the classical case of harmonic functions on smoothly bounded planar domains.

Section 6 Manifolds of bounded geometry that are Gromov hyperbolic form a rather special class of non-compact manifolds. However, many other geometries admit canonical deformations into such hyperbolic manifolds. Basic examples are the so-called *uniform* domains in \mathbb{R}^n . This remarkable concept covers smoothly bounded domains but also some classes of fractals like snowflakes. We show how we can use the hyperbolic theory to gain control over the classical potential theory on such Euclidean domains from the insights of section 5 .

2 Basic Concepts

2.1 Geometric Structures

We start with a discussion of those properties we assume throughout for our spaces and operators. Our space (M^n, g) is a connected, complete C^∞ -smooth Riemannian manifold of dimension $n \geq 2$. We assume the following two conditions. The first one, of having bounded geometry, is basic for a fine control of the potential theory. It is a homogeneity condition saying that the space locally looks the same around any point, up to a uniformly controlled deviation:

Bounded Geometry *For each ball $B_\sigma(p) \subset M$ there is a smooth ℓ -bi-Lipschitz chart ϕ_p to an open subset of U_p of \mathbb{R}^n with its Euclidean metric, where $\sigma = \sigma_M > 0$, $\ell = \ell_M \geq 1$ are fixed constants for the given manifold M . We call this a space of bounded geometry, or more precisely, (σ, ℓ) -bounded geometry.*

On spaces of bounded geometry one can link any two points $p, q \in M$ by a sequence of such balls transferring information inductively from one end to the other. Such configurations are known as Harnack chains due to their frequent use in the context of Harnack inequalities in elliptic analysis.

Harnack Chains For some fixed $r \in (0, \sigma)$ we call a sequence of balls $B_r(x_1), \dots, B_r(x_k)$ with

$$x_1 = p, x_k = q \text{ and } d(x_i, x_{i+1}) < r/2 \text{ for } i = 1, \dots, k-1$$

a **Harnack chain** of length k .

Thus any point x_i is contained in the ball $B_{r/2}(x_{i+1})$. If we link p and q by a curve $\gamma_{p,q}$ with $\text{length}(\gamma_{p,q}) = d(p, q)$ we can cover $\gamma_{p,q}$ by balls forming a Harnack chain. From this we see that, for given r , we can choose a chain of length proportional to the distance of the endpoints.




Figure 1: Harnack Chains

The second condition, of Gromov hyperbolicity, that we impose on M is, in some sense, complementary to the bounded geometry assumption. It has no local impact but strong consequences for the geometry near infinity.

Gromov Hyperbolicity A Riemannian manifold (M^n, g) , now regarded as a metric space, is Gromov hyperbolic or, quantitatively, **δ -hyperbolic**, if all its geodesic triangles are **δ -thin**, for some $\delta = \delta_M > 0$. This means that each point on the edge of any geodesic triangle is within δ -distance of one of the other two edges.

The concept of Gromov hyperbolic spaces embraces a broad range of metric spaces, prototypes from either side are classical hyperbolic spaces \mathbb{H}^n and abstract trees. While the former can be generalized to simply connected manifolds of sectional curvature less than a negative constant, δ -hyperbolicity of a graph expresses that it is near to a tree on a large scale. Gromov hyperbolicity is designed to study the asymptotic behavior of spaces near infinity, cf. [BH99, ch. III.H] and [KB02] for detailed expositions.

Every compact manifold is Gromov hyperbolic since we can choose $\delta = \text{diam}(M)$. This shows that this type of hyperbolicity has no local impact. However, near infinity Gromov hyperbolicity can be more demanding even than having constant negative sectional curvature. An instructive example are \mathbb{Z}^2 -coverings of genus ≥ 2 Riemann surfaces equipped with some hyperbolic metric. They have bounded geometry but they are not Gromov hyperbolic since we get arbitrarily large geodesic triangles uniformly close to Euclidean triangles in Hausdorff metric.

The motivating example for Ancona in [Anc87] were Cartan-Hadamard manifolds of sectional curvature pinched between two negative constants, which also ensures bounded geometry.

A larger and arguably more basic class of further examples for Gromov hyperbolic spaces of bounded geometry is given by uniform domains in Euclidean space with their quasi-hyperbolic metric as explained in [section 6](#).

Gromov boundary For a complete Gromov hyperbolic space X we define the set $\partial_G X$ of equivalence classes $[\gamma]$ of geodesic rays, from a fixed basepoint $p \in X$, where two rays are equivalent if they have finite Hausdorff distance. $\partial_G X$ is called the **Gromov boundary** of X . ($\partial_G X$ does not depend on the choice of the basepoint p .)

To define a topology on $\overline{X}^G = X \cup \partial_G X$, we first recall that a *generalized geodesic ray* $\gamma : I \rightarrow X$ is an isometric embedding of an interval $I \subset \mathbb{R}$ into X , where either $I = [0, \infty)$, then γ is a proper geodesic ray, or $I = [0, R]$, for some $R \in (0, \infty)$. Then γ is a geodesic arc. When we fix a basepoint $p \in X$ we can assign to any $x \in X$ a (not necessarily unique) generalized ray γ_x with endpoint $\gamma_x(R) = x$. For the following discussion we extend the definition of such a ray to $I = [0, \infty]$ setting $\gamma_x(t) = x$ for $t \in [R, \infty]$.

Then the topology on \overline{X}^G can be characterized from the following notion of a converging sequence: $x_n \in \overline{X}^G$ converges to $x \in \overline{X}^G$ if there exist generalized rays γ_{x_n} with $\gamma_{x_n}(0) = p$ and $\gamma_{x_n}(\infty) = x_n$ subconverging (on compact sets) to a generalized ray γ_x with $\gamma_x(0) = p$ and $\gamma_x(\infty) = x$. The canonical map $X \hookrightarrow \overline{X}^G$ is a homeomorphism onto its image, $\partial_G X$ is closed and \overline{X}^G is compact, see [BH99, III.H.3.7]. \overline{X}^G is called the **Gromov compactification** of X . It is a metrizable space.

To exploit Gromov hyperbolicity analytically Ancona introduced the following concept of Φ -chains. As in the case of bounded geometry versus hyperbolicity, Harnack chains and Φ -chains act complementary. This is best seen in the discussion preceding [Proposition 5.2](#) “[growth recovery along \$\Phi\$ -chains](#)” where Harnack chains give basic estimates which, however, weaken the overall control, while Φ -chains can be used to recover the apparently lost details.

Φ -Chains For a monotonically increasing function $\Phi : [0, \infty) \rightarrow (0, \infty)$ with $\Phi_0 := \Phi(0) > 0$ and $\Phi(d) \xrightarrow{d \rightarrow \infty} \infty$, a **Φ -chain** on X is a finite or infinite sequence $U_1 \supset U_2 \supset \dots \supset U_m$ of open subsets of X together with a sequence of **track points** x_1, x_2, \dots, x_m such that

- (i) $\Phi_0 \leq d(x_i, x_{i+1}) \leq 3\Phi_0$,
 - (ii) $x_i \in \partial U_i$,
 - (iii) $d(x, \partial U_{i\pm 1}) \geq \Phi(d(x, x_i))$, for every $x \in \partial U_i$
- for every i where applicable.¹

Note that a Φ -chain traversed backwards, i.e., with sets $X \setminus U_m \supset X \setminus U_{m-1} \supset \dots \supset X \setminus U_1$, is again a Φ -chain with the same track points.

The existence of infinite Φ -chains can be considered as a partial hyperbolicity property of the underlying space. It is easy to see that neither the Euclidean nor asymptotically flat spaces admit any infinite Φ -chains.

A first non-trivial example can be created as follows: with the coordinates $(x, y) \in \mathbb{R} \times \mathbb{R}^{n-1}$ we consider the metric $(1 + |y|^2)^2 \cdot g_{\mathbb{R}} + g_{\text{Eucl}}$ on $\mathbb{R} \times \mathbb{R}^{n-1}$. Then the half-spaces $U_i := \mathbb{R}^{\geq i} \times \mathbb{R}^{n-1}$ form a Φ -chain with track points $x_i = (i, 0)$ for $\Phi(t) := 1 + t^2$. But, as in the Euclidean case, the half-spaces $U_i[k] := \mathbb{R}^k \times \mathbb{R}^{\geq i} \times \mathbb{R}^{n-k-1}$, for $1 \leq k \leq n-1$, do not make up a Φ -chain.

The hyperbolic n -space \mathbb{H}^n carries Φ -chains in all directions and for the same Φ . Namely, when we think of \mathbb{H}^n as the upper half-space and $B_{1/2^i}(0)$ is the Euclidean ball of radius $1/2^i$, then $U_i = B_{1/2^i}(0)$, $i \geq 1$, form a Φ -chain for $\Phi(t) = a + b \cdot t$ for suitable $a, b > 0$ independent

¹For notational convenience this is slightly different from Ancona’s version in [Anc90, définitions V.5.1], but essentially the same.




Figure 2: Φ -Chains

even of n . Due to the homogeneity of \mathbb{H}^n this also gives a Φ -chain along any hyperbolic geodesic γ , i.e., with track points on γ .

This ubiquity of Φ -chains which we see already from considering hyperbolic half-spaces relative to geodesics extends to arbitrary non-homogenous Gromov hyperbolic spaces.

Φ -Chains on Hyperbolic Spaces [BHK01, Section 8] *For a general δ -hyperbolic X we choose a geodesic $\gamma : \mathbb{R} \rightarrow X$, and define*

$$U_i := \{x \in X \mid \text{dist}(x, \gamma([22\delta i, +\infty))) < \text{dist}(x, \gamma((-\infty, 22\delta i]))\} .$$

Then the U_i form a Φ_δ -chain with track points $x_i = \gamma \cap \partial U_i$ for $\Phi_\delta(t) = a_\delta + b_\delta \cdot t$ for suitable $a_\delta, b_\delta > 0$. Moreover, the U_i and the $X \setminus U_i$ form neighborhood bases in the topology of \bar{X}^G around the two ideal endpoints of the geodesic γ in $\partial_G X$.

The constants a_δ and b_δ can be inferred from the proof of [BHK01, Lemma 8.9].

2.2 Analytic Structures

Now we turn to the analytic side. On a complete Riemannian manifold M with bounded geometry, we consider an operator L with the following properties:

Adaptedness of L *L is a linear second order elliptic operator on M such that relative to the charts ϕ_p ,*

$$L(u) = - \sum_{i,j} a_{ij} \cdot \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_i b_i \cdot \frac{\partial u}{\partial x_i} + c \cdot u$$

for $(2, \beta)$ -Hölder continuous a_{ij} , $(1, \beta)$ -Hölder continuous b_i and β -Hölder continuous c , $\beta \in (0, 1]$, with

$$k^{-1} \sum_i \xi_i^2 \leq \sum_{i,j} a_{ij} \xi_i \xi_j \leq k \sum_i \xi_i^2$$

for any $\xi \in \mathbb{R}^n$ and

$$|a_{ij}|_{C^2(U_p)}, |b_i|_{C^1(U_p)}, |c|_{L^\infty(U_p)} \leq k$$

for some $k = k_L \geq 1$ and any $p \in M$.

Solutions of the equation $Lw = 0$ are called **L-harmonic** functions. We also consider L -sub- and superharmonic functions satisfying, in the smooth case, $Lw \leq 0$ or $Lw \geq 0$, respectively. More general notions of L -sub/superharmonicity will be introduced later on.

The adaptedness assumption is a generalized form of uniform ellipticity. It is a compromise between generality and readability, trimmed to ensure that one can apply the usual maximum principles, all weak solutions are classical $C^{2,\beta}$ -regular solutions, the coefficients are bounded and that the adjoint operator L^* shares (even quantitatively) the same properties because it still has Hölder continuous coefficients bounded by $3k$, cf. [GT98, Chapter 8]. For more general conditions, see [Anc87].

Towards the potential theory of such operators, we need the existence of a **Green's function** G for L . This is a function $G(x, y) > 0$ on $M \times M$ which is singular on the diagonal $\{(x, x) \mid x \in M\}$ and satisfies the equation $LG(\cdot, y) = \delta_y$, in a distributional sense, where δ_y is the Dirac delta function with basepoint y . Put differently, for any $y \in M$, $G(\cdot, y) > 0$ is a (singular) function so that for a given function f , the function $u(x) = \int_M G(x, y) f(y) \, d\text{Vol}(y)$ is a solution of $Lu = f$, where $d\text{Vol}$ is the volume measure on M . Then a Green's function associated to L^* is given by $G^*(x, y) = G(y, x)$.

The existence of a Green's function is not always granted. As a quite typical example for our further discussion we consider the Laplacian $-\Delta$ on the flat \mathbb{R}^n . The operator $-\Delta - \lambda \text{Id}$ admits a Green's function if and only if $\lambda \leq 0$. Moreover, we will see later on that there is a fine distinction between the cases $\lambda = 0$ and $\lambda < 0$. The following assumption we make for our operator L means that we are focussing on the second case.

Weak Coercivity For some $t > 0$, $L_t := L - t \text{Id}$ admits a Green's function G^t .

The weak coercivity condition is generic as soon as a Green's function exists. More precisely there is a *generalized principal eigenvalue* $\tau = \tau_L$ such that for all $t < \tau$ the Green's function G^t exists, while for $t > \tau$ there is no globally defined positive L_t -harmonic function. In the borderline case $t = \tau$, the Green's function G^τ might or might not exist, but in any case there is a globally defined positive L_τ -harmonic function [Pin95, sec. 4.3]. Then weak coercivity amounts to $\tau > 0$.

Another way to express weak coercivity is to say that there is a positive supersolution u of the equation $Lu = 0$ with the quantitative estimate $Lu \geq \varepsilon \cdot u$, for some $\varepsilon > 0$.

Note that with L the operator L_t is again adapted and for $t < \tau$ even weakly coercive.

2.3 Potentials and Balayage

The sum of a Green's function and a positive solution of $Lu = 0$ is again a Green's function. Therefore one usually focusses on the uniquely determined *minimal* Green's function which does not admit a sum decomposition into another Green's function and a positive solution. More generally, any Green's function is an **L-superharmonic** function, that is a lower semi-continuous function with values in $(-\infty, \infty]$ that is larger than the Dirichlet solution on any ball with the same boundary conditions and finite on a dense set. On smooth parts, this is equivalent to $Lu \geq 0$. An L -superharmonic function $p \geq 0$ without a minorizing positive L -harmonic function h , that is with $p \geq h > 0$, is called a **potential**. Minimal Green's functions are not only examples for potentials, but also the basic building blocks:

Theorem 2.1 (Integral Representation of Potentials) [Her62, 22.], [Hel14, Thm. 6.18]
Every potential p on M can be represented by a unique (positive) Radon measure μ_p as

$$p(x) = G(\mu_p)(x) := \int_M G(x, y) d\mu_p(y) .$$

The support of p (i.e. the complement of the largest open set where p is L -harmonic) equals the support of μ_p .

The fact that every positive L -superharmonic function is uniquely representable as the sum of a potential and a positive L -harmonic function is known as the *Riesz representation theorem*. Hence, to get an integral representation for positive L -superharmonic functions, only the L -harmonic part is left. The corresponding measures are supported on the *Martin boundary* which is defined exactly for this purpose. Its identification in terms of more common geometric boundaries is the subject of subsections 5.3 and 6.3.

Later we want to control L -superharmonic functions along Φ -chains. Here we shift the part of the defining measure supported in $M \setminus U_i$ onto ∂U_i without changing the function on U_i . This strategy is called sweeping or, due to its French origin (Poincaré, Cartan), **balayage**.

Concretely, for an L -superharmonic function $u \geq 0$ on M and a subset $A \subset M$ we consider

$$\mathcal{R}_u^A := \inf\{v \geq 0 \mid v \text{ is } L\text{-superharmonic on } M \text{ with } v \geq u \text{ on } A\} .$$

This is called the *reduit* (reduced). It enjoys the following properties we will need later:

- \mathcal{R}_u^A is L -harmonic outside of \bar{A} and equal to u in A .
- The *reduit* is always L -superharmonic.
- If A is relatively compact, \mathcal{R}_u^A is a potential.
- $\mathcal{R}_{\lambda u}^A = \lambda \mathcal{R}_u^A$ for a constant $\lambda \geq 0$, if u is a positive function.
- $\mathcal{R}_{u+v}^A = \mathcal{R}_u^A + \mathcal{R}_v^A$ for functions u, v .
- $\mathcal{R}_u^{A \cup B} \leq \mathcal{R}_u^A + \mathcal{R}_u^B$ for sets $A, B \subset M$.
- Denoting the *reduit* with respect to the adjoint operator L^* of L by ${}^*\mathcal{R}$, we have $\mathcal{R}_{G(\cdot, y)}^A(x) = {}^*\mathcal{R}_{G(x, \cdot)}^A(y)$ [Anc90, I.5.1, p. 19].
- If $G(\mu)$ is a potential, $\mathcal{R}_{G(\mu)}^A(x) = \int_M \mathcal{R}_{G(\cdot, y)}^A(x) d\mu(y)$ for any $x \notin \bar{A}$ [Her62, Théorème 22.4].

For general sets A , it may happen that \mathcal{R}_u^A is not lower semi-continuous, but it always admits a canonical regularization $\widehat{\mathcal{R}}_u^A$, the *balayée* (swept). For open sets or in general outside of \bar{A} the two concepts coincide. We can even recover the classical Perron solution u of the Dirichlet problem on a ball B with (say, continuous) boundary value f as

$$u(x) = \mathcal{R}_f^{\partial B}(x) .$$

Also useful in this context are global variants of the maximum principle.

Theorem 2.2 (Global Maximum Principle) [Her62, p. 429]

- (i) If u is L -superharmonic on an open set $V \subset M$, $u \geq 0$ on ∂V , and there is a potential p such that $u \geq -p$, then $u \geq 0$ on V .

(ii) Let p a potential, L -harmonic on an open set V and locally upper bounded near every point of ∂V . If $u \geq p$ on ∂V for some positive L -superharmonic function u , then $u \geq p$ in all of V .

Proof. For (i), the function \bar{u} defined as $\min(u, 0)$ on V and 0 on $M \setminus V$ is L -superharmonic and $\geq -p$. Now the supremum of the family $\{L$ -subharmonic functions $\leq \bar{u}\}$ is L -harmonic, $\geq -p$, and ≤ 0 , hence by the definition of potentials it is 0 which implies $u \geq 0$.

(ii) follows from (i) by considering the function $u - p$. □

3 Local Maximum Principles and Harnack Inequalities

We derive estimates for positive solutions and Green's functions of L_t on uniformly sized balls $B_R := B_{\sigma/\ell} := B_{\sigma/\ell}(0) \subset \mathbb{R}^n$ in the adapted charts. For notational convenience, in this chapter, all balls are measured in the Euclidean distance. The required assumptions on Locally defined, Adapted operators with a Green's function will be called

Assumption (LAG) We say that a differential operator L on $B_R(0) \subset \mathbb{R}^n$ fulfills assumption (LAG) for $R > 0$, $k > 0$ and $n \geq 2$ if

- L is k -adapted on $B_R(0)$ and
- there is a Green's function for L on $B_R(0)$.

Later all results can be transferred to the manifold using the bi-Lipschitz charts with only a small loss in constants. Weak coercivity will not yet be used explicitly, but the results and estimates apply to all operators L_t with $0 \leq t \leq \tau$, where the constants now depend on τ as well (via k), but not on t .

3.1 Maximum Principles

The following version of the weak maximum principle does not require positivity of the "potential" term c in L , the existence of a Green's function (or equivalently, a positive L -superharmonic function) is sufficient, but the result is weaker. It can easily be obtained from the version for $c \geq 0$ using the h -transform, see [Pin95], where the sign convention for L is different from ours.

Theorem 3.1 (Local Minimum-Zero Principle) [Pin95, Theorem 3.2.2, p. 81] *If L satisfies assumption (LAG), u is continuous on B_R , L -superharmonic on a domain $D \Subset B_R$ and $u|_{\partial D} \geq 0$, then $u \geq 0$ in all of D .*

With help of the uniform bounds on the coefficients, this can be upgraded to quantitative bounds in terms of the boundary values.

Lemma 3.2 (Local Almost-Maximum Principle) *If L satisfies (LAG), there are constants $0 < r_{\text{am}} < R$ and $m > 0$ depending only on k and n , such that any L -subharmonic function u on B_r , $r < r_{\text{am}}$, with $u|_{\partial B_r} \leq m$ satisfies $u \leq 1$ on B_r .*

Proof. In polar coordinates, L applied to a radial function $f(r)$ can be written as

$$Lf = -\alpha f''(r) + \frac{\alpha - \text{tr } a}{r} f'(r) + bf(r) + cf(r)$$

with $k^{-1} \leq \alpha := \frac{a_{ij}(x)x^i x^j}{r^2} \leq k$, $nk^{-1} \leq \text{tr } a \leq nk$ and $|b|, |c|$ bounded by k .

Consider the function $f(r) = \text{sech}(\beta r)$ with some $\beta > 0$. We want to have the following properties:

- $Lf \geq 0$ for $r < r_{\text{am}}$,
- $f \leq 1$, and
- $f|_{\partial B_r} \geq \text{sech}(\beta r_{\text{am}}) =: m$ for $r < r_{\text{am}}$.

Only the first property is not obvious and we need to tune the parameters r_{am} and β to achieve it. We can calculate

$$Lf = \left[\alpha \beta^2 (1 - \sinh^2(\beta r)) \text{sech}^2(\beta r) + (\text{tr } a - \alpha) \beta^2 \frac{\tanh(\beta r)}{\beta r} - b \beta \tanh(\beta r) + c \right] \text{sech}(\beta r) .$$

Here the first *two* terms behave as β^2 for large β and $\beta r < 1$. But in leading order in βr the α terms cancel, while the positive $\text{tr } a$ term survives. Using the uniform bounds on α and $\text{tr } a$ in terms of k and n , we first fix a small βr_{am} so that the β^2 terms are positively bounded below and then make β large to kill the remaining terms and get $Lf \geq 0$ for $r < r_{\text{am}}$ where r_{am} and β depend only on k and n .

Now we can apply the [local minimum-zero principle 3.1](#) to $f - u$, because $L(f - u) \geq 0$ and $(f - u)|_{\partial B_r} \geq 0$. This yields $u \leq f \leq 1$ on B_r . \square

With completely different methods we can see that Green's functions behave for small distances like the Euclidean Green's function

$$G_{\text{Eucl}}(x, y) = \begin{cases} \frac{1}{2\pi} \log |x - y| , & \text{if } n = 2 \\ \frac{1}{(n-2) \text{Vol } S^{n-1}} \frac{1}{|x-y|^{n-2}} , & \text{if } n \geq 3 . \end{cases}$$

Lemma 3.3 (Local Bound for Some Green's Function) *For L satisfying (LAG) there are constants $r_{\text{gb}} > 0$, $\tilde{q} \geq 1$ depending only on k and n and some (not necessarily minimal) Green's function $\tilde{G}(x, y)$ such that*

$$\tilde{q}^{-1} G_{\text{eucl}}(x, y) \leq \tilde{G}(x, y) \leq \tilde{q} G_{\text{eucl}}(x, y) \quad \text{for } x, y \in B_{r_{\text{gb}}} .$$

Proof. We sketch the argument for $a_{ij} = \delta_{ij}$ and $n \geq 3$. The general case is merely notationally more involved and can be found in [Mir70, pp. 61–63].

As distributions, we have $L_x G_{\text{Eucl}}(x, y) = \delta(x - y) + R(x, y)$, where $R(x, y)$ has a singularity at $x = y$ of order $\mathcal{O}(r^{-(n-1)})$. This means that we can choose $r_{\text{gb}} > 0$ so that $\int_{B_{r_{\text{gb}}}} |R(x, y)| dy < 1/2$ for $x \in B_{r_{\text{gb}}}$ because the integral scales as $\mathcal{O}(r_{\text{gb}})$ and all involved constants depend only on the bounds on the coefficients k . This in turn implies the finiteness of all iterates R^n where the product of kernels is defined as $R^2(x, y) = \int_{B_{r_{\text{gb}}}} R(x, z)R(z, y)dz$. The bound ensures also the summability of

$$\tilde{G} := G_{\text{Eucl}} \sum_{i=0}^{\infty} (-R)^n ,$$

again as products of kernels. Application of L yields

$$L\tilde{G} = (\delta + R) \sum_{i=0}^{\infty} (-R)^n = \sum_{i=0}^{\infty} (-R)^n - \sum_{i=1}^{\infty} (-R)^n = \delta ,$$

so that \tilde{G} is indeed a Green's function. $\tilde{G} - G_{\text{Eucl}}$ has a singularity at $x = y$ of quantitatively lower order than G_{Eucl} , hence the explicit bounds. \square

Corollary 3.4 (Local Bound for the Minimal Green's Function) *If L fulfills (LAG), there are constants $q > 0$ and $0 < r_{\text{mgbi}} < r_{\text{mgbo}} := \min(r_{\text{am}}, r_{\text{gb}})$ depending only on k and n such that the minimal Green's function $g(x, y)$ of $B_{r_{\text{mgbo}}}$ (i.e., vanishing on the boundary $\partial B_{r_{\text{mgbo}}}$) satisfies*

$$q^{-1} G_{\text{eucl}}(x, y) \leq g(x, y) \leq q G_{\text{eucl}}(x, y) \quad \text{for } x, y \in B_{r_{\text{mgbi}}} .$$

Proof. For fixed $y \in B_{r_{\text{mgbo}}/2}$, g can be represented as $g(x, y) = \tilde{G}(x, y) - u(x)$ where u is the solution of the Dirichlet problem $Lu = 0$ in $B_{r_{\text{mgbo}}}$ and $u(x) = \tilde{G}(x, y)$ on $\partial B_{r_{\text{mgbo}}}$. By the [local bound for some Green's function 3.3](#) we know

$$u(x) \leq \frac{\tilde{q}}{|x - y|^{n-2}} \leq \frac{\tilde{q}}{(r_{\text{mgbo}}/2)^{n-2}} \quad \text{for } x \in \partial B_{r_{\text{mgbo}}}$$

(and similarly for $n = 2$). Thus we can apply the [local almost-maximum principle 3.2](#) to get $u \leq \tilde{q}/(m(r_{\text{mgbo}}/2)^{n-2})$, whence

$$g(x, y) \geq \frac{\tilde{q}^{-1}}{|x - y|^{n-2}} - \frac{\tilde{q}}{m(r_{\text{mgbo}}/2)^{n-2}} \quad \text{for } x \in B_{r_{\text{mgbo}}} .$$

From this equation, $q \geq \tilde{q}$ and r_{mgbi} can be determined to obtain the lower bound. For the upper bound, note that \tilde{G} is positive and thus $g \leq \tilde{G}$ by the [local minimum-zero principle 3.1](#). \square

3.2 Harnack Inequalities

This standard result can be found e.g. in [GT98, Theorem 8.20], but we include a proof here because it is an easy consequence of the previous results.

Theorem 3.5 (Harnack Inequalities) *For L satisfying (LAG), there is an $H(k, n, R) \geq 1$ such that if $u > 0$ is L -harmonic on $B_r(x_0) \subset B_R$, for some $0 < r \leq R$, then*

$$H^{-1}u(x_0) \leq u(x) \leq Hu(x_0) \quad \text{for any } x \in B_{r/2}(x_0) .$$

Proof. We start with the case $r \leq r_{\text{mgbi}}$.

The potential $\mathcal{R}_u^{B_r(x_0)}$ (on the total space $B_{r_{\text{mgbo}}}$) can be represented as $\mathcal{R}_u^{B_r(x_0)} = g(\mu_u)$ for some positive measure μ_u with support in $\partial B_r(x_0)$. On $B_r(x_0)$, $\mathcal{R}_u^{B_r(x_0)}$ coincides with u and we have $u = g(\mu_u)$. Now for $y \in B_{r/2}(x_0)$, we can apply both estimates from [Corollary 3.4](#) “local bound for the minimal Green's function” to get

$$\begin{aligned} u(y) &= \int_{\partial B_r(x_0)} g(y, z) d\mu_u(z) \leq \int_{\partial B_r(x_0)} \frac{q}{(r/2)^{n-2}} d\mu_u(z) \\ &\leq 2^{n-2} q^2 \int_{\partial B_r(x_0)} g(x_0, z) d\mu_u(z) = 2^{n-2} q^2 u(x_0) . \end{aligned}$$

The other inequality is analogous.

For the case of larger r , we can employ the Harnack chains introduced in [subsection 2.1](#) with radius r_{mgbi} . With every ball the estimate collects the constant for that case, but the total number of balls needed is bounded above by R/r_{mgbi} , so we can just use the resulting power of $2^{n-2}q^2$ as H in every case. \square

The Harnack chain tactic just demonstrated will be used in the following to get estimates for even larger distances. The result are exponential bounds (with fixed constants) for the

growth of L -harmonic functions—that is not especially satisfying, but a good starting point for improvement.

This Harnack inequality always needs a little more space around the balls where it holds, but for instance on bounded domains, it would be useful to have a similar result near points on the boundary. A blueprint is the following classical result.

Theorem 3.6 (Boundary Harnack Inequality on a Disc) *There exist constants $A, C > 1$ such that for any point $\xi \in \partial B_1(0) \subset \mathbb{R}^2$ and $0 < R < 1$ the following is true: For any two harmonic functions $u, v > 0$ (with respect to the Laplacian) on $B_{A \cdot R}(\xi) \cap B_1(0)$ that vanish along $B_{A \cdot R}(\xi) \cap \partial B_1(0)$,*

$$\frac{u(x)}{v(x)} \leq C \frac{u(y)}{v(y)} \text{ for all } x, y \in B_R(\xi) \cap B_1(0) .$$

In the non-boundary version, the appearance of another function v is obscured by the fact that the constant function 1 is (nearly) harmonic. Furthermore, the restriction to functions vanishing on the relevant part of the boundary is unavoidable, yet in combination with balayage techniques there are powerful applications as we will see after the proof of a much more general boundary Harnack inequality in [subsection 5.2](#).

4 Global Results from Resolvent Equations and Bounded Geometry

Now we use that our manifold M has Bounded geometry and L is Adapted and weakly Coercive, more precisely:

Assumption (BAC) We say that the pair (M, L) fulfills assumption (BAC) for $\sigma > 0$, $\ell \geq 1$, $k > 0$, $n \geq 2$ and $\tau > 0$ if L is a differential operator on a connected complete noncompact Riemannian manifold M^n such that

- M is of (σ, ℓ) -bounded geometry,
- L is k -adapted in the bounded geometry charts, and
- L is weakly coercive with generalized principal eigenvalue $\tau =: 2\theta$.

This gives us global growth estimates for the minimal Green's function G of L . The basic idea is to combine the local estimates we derived in the last chapter with the resolvent equation for Green's functions viewed as operators.

While the boundedness of geometry and adaptedness of L allow to carry over all results from [section 3](#) with constants now depending on k, n, σ, ℓ and τ , as explained in the beginning of that section, the weak coercivity assumption is the most important new ingredient in this section. Note that we do not yet make use of Gromov hyperbolicity.

4.1 Resolvent Equation

For a closed operator L defined on a dense set of a Banach space the set $\rho(L)$ of all λ , so that the *resolvent* $R_\lambda = (L - \lambda \text{Id})^{-1}$ exists and is continuous, is called the *resolvent set*. $\rho(L)$ known to be an open set. For $\lambda, \mu \in \rho(L)$ we have the resolvent equation

$$R_\lambda \circ R_\mu = (\lambda - \mu)^{-1} \cdot (R_\lambda - R_\mu).$$

One of the main applications of this identity in the context of elliptic operators is the comparison of solutions of $R_\lambda w = 0$ with those of $R_\mu w = 0$. We will use this idea in the special

case of the minimal Green's functions G^t corresponding to the operators $L^t = L - t\text{Id}$ on M , viewed as resolvents. We include the simple proof of the resolvent equation in our context and note some of its consequences.

Lemma 4.1 (Resolvent Equation) *Assume (M, L) satisfies assumption (BAC). For any $0 \leq t < \tau$, the minimal Green's function G^t , viewed as an operator on the space of positive Radon measures μ with $G^t\mu < \infty$, satisfies*

$$G^t = G + tG \circ G^t .$$

Since all involved operators are positive, this yields the inequalities

$$G \leq G^t \quad \text{and} \tag{1}$$

$$G \circ G^t \leq \frac{1}{t}G^t . \tag{2}$$

In particular, these results hold in the cases of characteristic functions of bounded measurable sets or Dirac measures.

Proof. Consider an increasing sequence of relatively compact, smoothly bounded open sets (U_i) with $\bigcup U_i = M$. On each U_i , the corresponding Dirichlet Green's function G_i (i.e., $G_i(\cdot, y)|_{\partial U_i} \equiv 0$) satisfies the resolvent equation in the form

$$G_i(x, y) = G_i^t(x, y) - t \int_{U_i} G_i(x, z)G_i^t(z, y)dz$$

because the right-hand side fulfills the properties of a Green's function (application of L yields δ_y) and has the correct boundary behavior. To see that the integral is finite for $x \neq y$, notice that any Green's function is integrable near its pole by the same arguments as in the proof of [Corollary 3.4](#) "local bound for the minimal Green's function".

For $i \rightarrow \infty$, the G_i and G_i^t converge to G and G^t , respectively, uniformly on compact sets away from the pole because they are increasing and bounded above by the minimal Green's function. The same is true if the first argument is fixed because then one has the Green's function for the adjoint operator. It is easy to see that the equation survives in the limit $i \rightarrow \infty$.

By integration and Fubini's theorem one obtains the resolvent equation for arbitrary Radon measures μ , as soon as $G^t\mu < \infty$ (and thereby $G\mu < \infty$). \square

Recall from [subsection 2.2](#) that unlike L_t , for $t < \tau$, the operator L_τ is no longer weakly coercive. In turn, we observe from (2) that using resolvents loses its strength when t approaches 0. This suggests to work with L_θ , for

$$\theta := \tau/2.$$

4.2 Behavior of Green's functions

We combine the resolvent equation with [Harnack inequalities 3.5](#) to derive growth estimates of minimal Green's functions.

Proposition 4.2 (Bound for the Green's Function) *Given (M, L) satisfying (BAC), there is a constant $c_1(\sigma, \ell, k, n, \tau) \geq 1$ such that for the minimal Green's function of L ,*

$$\begin{aligned} c_1^{-1} &\leq G(x, y) && \text{if } d(x, y) \leq \sigma, \text{ and} \\ &G(x, y) \leq c_1 && \text{if } d(x, y) = \sigma . \end{aligned}$$

Proof. The lower bound is directly obtained from the [local bound for the minimal Green's function 3.4](#) because we have $G(\cdot, y) \geq g(\cdot, y)$ for any Green's function g on a smaller domain and iterated application of the [Harnack inequalities 3.5](#) (a number depending on σ , ℓ and r_{mgbi}).

For the upper bound, consider the ball $B := B_{\sigma/3}(x)$. The function $G^\theta(\chi_B) = \int_B G^\theta(\cdot, \zeta) d\zeta$ is bounded on B , where χ_B denotes the characteristic function of B , because we can apply the lower bound $1 \leq c_1 \cdot G(\zeta, z')$ to some z' with $d(x, z') = 2\sigma/3$ and arrive at

$$\int_B G^\theta(z, \zeta) d\zeta \leq c_1 \int_M G^\theta(z, \zeta) G(\zeta, z') d\zeta \leq \frac{c_1}{\theta} G^\theta(z, z')$$

for all $z \in B$. In the second inequality we used (2) for the Dirac function in z' . By continuity of $G^\theta(\cdot, z')$ on the compact \bar{B} this is uniformly bounded.

Thus we can integrate the [resolvent equation 4.1](#) applied to χ_B over B and get

$$\infty > \int_M \chi_B G^\theta(\chi_B) = \int_M \chi_B G(\chi_B) + \theta \int_M \chi_B G(G^\theta(\chi_B)) .$$

When G^* denotes the minimal Green's function for the adjoint operator L^* this means

$$0 < \int_M \chi_B G(\chi_B) = \int_M G^\theta(\chi_B) [\chi_B - \theta G^*(\chi_B)] .$$

Hence, there must be an $x' \in B$ such that $(\chi_B - \theta G^*(\chi_B))(x') > 0$ and therefore

$$\int_B G(\zeta, x') d\zeta = G^*(\chi_B)(x') < 1/\theta$$

which in turn shows the existence of an $x'' \in B$ with $d(x', x'') > \sigma/12$ and

$$G(x'', x') \leq \theta^{-1} (\text{Vol}(B \setminus B_{\sigma/12}(x'')))^{-1} .$$

Now the bounded geometry assumption assures that the volume can be bounded below by a constant $V_{\sigma, \ell, n}$ depending only on σ , the Lipschitz constant ℓ and the dimension n .

From the bounded geometry constraint, we observe that a Harnack chain of at most $\lceil 24 + \pi\ell \rceil$ balls $B_r(p_i)$, $i = 1, \dots, \lceil 24 + \pi\ell \rceil$, of radius $r = \sigma/12$ with $d(p_i, p_{i+1}) < r/2$ and $d(p_i, x'') > r$ suffices to link x' with either x or y . We get similar chains starting from x'' to y or x , respectively. Applying the [Harnack inequalities 3.5](#) on each of these balls and multiplying the Harnack constants step by step to get H^i on $B_r(p_i)$ we get the assertion for $c_1 = H^{2 \cdot \lceil 24 + \pi\ell \rceil} \cdot \theta^{-1} V_{\sigma, \ell}^{-1}$. \square

Note that we could now easily get more explicit growth estimates for G by comparison with the [local bound for the minimal Green's function 3.4](#), but for the following these coarse estimates are sufficient.

4.3 Relative Maximum Principles

We start with an elementary comparison result which already visualizes the central effect of weak coercivity: with the same boundary conditions, compared to L_θ -harmonic functions, L -harmonic functions *sag*, imagine a rope² or a rubber blanket where you apply less and less tension. To see this, we start with a paraboloid that fits in between.

²This analogy is quite precise, a suspended rope forms a hyperbolic cosine, a solution of the shifted one-dimensional Laplace equation.

Lemma 4.3 *Under the local assumptions (LAG), there is a constant $m(R = \sigma/\ell, k, n) > 0$ such that we can find a smooth function f defined on $B_{R/2}(0)$ with*

$$f(x) = -m, \quad f|_{\partial B_{R/4}(0)} \geq 0 \quad \text{and} \quad Lf > -1.$$

Proof. In polar coordinates, L applied to a radial function $f(r)$ can be written as

$$Lf(r) = -af''(r) + \frac{b}{r}f'(r) + cf(r)$$

with functions a, b, c bounded by a constant depending only on k, n and R .

With the ansatz $f(r) = s(r^2 - \rho)$ we have $Lf(r) = -2as + 2bs + cs(r^2 - \rho)$ and first choose $\rho > 0$ sufficiently small to assure $f \geq 0$ outside of $B_{R/4}(0)$ and then make $s > 0$ small to achieve $Lf \geq -1$. This only uses the bounds on coefficients such that $f(0) = -s\rho =: -m < 0$ depends only on k, n and R . \square

Proposition 4.4 (Relative Maximum Principle, Local Version) *Under assumption (LAG) for L and $L_\theta = L - \theta \text{Id}$, assume further we have two functions u and $\bar{u} > 0$ on $B_R = B_R(0)$, u L -subharmonic ($Lu \leq 0$) and \bar{u} L_θ -harmonic on B_R with $\bar{u}|_{\partial B_{R/4}} \geq u|_{\partial B_{R/4}}$. Then there is a constant $\tilde{\eta} = \tilde{\eta}(R, k, n, \tau) \in (0, 1)$ such that*

$$u(0) \leq \tilde{\eta} \bar{u}(0).$$

Proof. Consider the function $h(z) := \bar{u}(z) + \theta f(z) - \inf_{w \in B_{R/4}} \bar{u}(w) - u(z)$ where f is the function from the previous lemma on $B_{R/4}$. On $B_{R/4}$, we have $L\bar{u} = L_\theta \bar{u} + \theta \bar{u} \geq \theta \bar{u}$, $Lf > -1$ and therefore

$$Lh(z) > \theta \left(\bar{u}(z) - \inf_{w \in B_{R/4}} \bar{u}(w) \right) \geq 0.$$

The boundary condition together with $f|_{\partial B_{R/4}} \geq 0$ yields $h|_{\partial B_{R/4}} \geq 0$ and we can apply the [local minimum-zero principle 3.1](#) for L to see $h \geq 0$ in the interior of $B_{R/4}$ and especially $h(0) \geq 0$, with $f(0) = -m$ we have

$$u(0) = \bar{u}(0) - m \inf_{w \in B_{R/4}} \bar{u}(w) \leq (1 - m\theta H^{-1}) \bar{u}(0).$$

The harmonicity on the full ball B_R was used only in the last step to apply the [Harnack inequalities 3.5](#). \square

Applied globally, this describes the relative growth of L -harmonic versus L_θ -harmonic functions.

Proposition 4.5 (Relative Maximum Principle, Global Version) *If (M, L) satisfies (BAC), there is a constant $\eta = \eta(\sigma, \ell, k, n, \tau) \in (0, 1)$ such that the following holds: Assume we have two functions u and \bar{u} defined on $B_{r+3}(x)$ for some $x \in M$ and $r \geq \sigma$, u L -subharmonic and \bar{u} L_θ -harmonic on $B_{r+3}(x)$ and $\bar{u}|_{\partial B_r(x)} \geq u|_{\partial B_r(x)}$. Then*

$$u(x) \leq \eta^r \bar{u}(x).$$

Proof. For integer multiples r of $\sigma/4\ell$, this is proven by inductively applying [the local version](#) in adapted charts along a chain of intersecting balls of length proportional to $r/(\sigma/\ell)$. On each of them we may apply the same Harnack inequality as described in the proof of [4.2](#) above, so that we can choose η as a function of σ, ℓ, H and $\tilde{\eta}$. \square

For Green's functions, we get the following variants. We do not use them in the following arguments but they are worth being mentioned since they give some non-trivial constraints on the Green's functions from our standard assumption (BAC).

Corollary 4.6 (Exponentially Stronger Decay) *Under assumptions (BAC), there are constants $A(\sigma, \ell, k, n, \tau) > 0$ and $\alpha_1(\sigma, \ell, k, n, \tau) > 0$ such that*

$$G(x, y) \leq A e^{-\alpha_1 d(x, y)} G^\theta(x, y) \quad \forall x, y \in M .$$

From this we get the following growth estimate for Green's functions using the resolvent equation.

Proposition 4.7 (Exponential Decay) *Under assumptions (BAC), for suitable constants $B(\sigma, \ell, k, n, \tau) > 0$ and $\alpha_2(\sigma, \ell, k, n, \tau) > 0$ we have*

$$G(x, y)G(y, x) \leq B e^{-\alpha_2 d(x, y)} \quad \text{for } d(x, y) > 2\sigma .$$

Proof. Let $x' \in M$ such that $d(x, x') = \sigma$. Then employing the resolvent equation 4.1 we have

$$\begin{aligned} G(x, y)G(y, x) &\leq G(x, y) H^3 G(y, x') \leq \frac{H^5}{\text{Vol}(B_{\sigma/2}(y))} \int_{B_{\sigma/2}(y)} G(x, z)G(z, x') dz \\ &\leq c \int_M G(x, z)G(z, x') dz \stackrel{(1),(2)}{\leq} \frac{c}{\theta} G^\theta(x, x') \leq \frac{cc_1^\theta}{\theta} \end{aligned}$$

where we used the Harnack inequalities 3.5 for L^* and the bound for the Green's function 4.2 applied to L_θ which itself satisfies the assumptions, but it may lead to a weaker constant c_1^θ .

For the very same reason, we can do all of the above with G^θ instead of G to get the uniform boundedness of $G^\theta(x, y)G^\theta(y, x)$, again with slightly worse constants. Combined with Corollary 4.6 “exponentially stronger decay” we have proved the assertion. \square

In the case of a self-adjoint operator $L = L^*$ the Green's function is symmetric and the proposition says that it decays exponentially with the distance. This result does not use Gromov hyperbolicity and holds also e.g. in Euclidean space. The difference from the familiar Euclidean Laplacian's Green's function is owed to weak coercivity.

5 Hyperbolicity and Boundary Harnack Inequalities

Now we additionally invest the hyperbolicity of the underlying space. The property we employ is that in Gromov hyperbolic spaces any two of their points can be connected by well-controlled Φ -chains, cf. subsection 2.1. The results we prove are more general. They hold with respect to any individual Φ -chain even if the space carries essentially only this one Φ -chain as in the example $(\mathbb{R} \times \mathbb{R}^{n-1}, (1 + |y|^2)^2 \cdot g_{\mathbb{R}} + g_{\text{Euc}})$.

We exploit this to derive the main results for the potential theory on these spaces, the particular behavior of Green's function along Φ -chains, building on the results for bounded geometries we derived in the last few sections. From this we infer the boundary Harnack inequalities which largely dominate the potential theory of our elliptic operators.

Our general assumptions (BAC) on the manifold M and the elliptic operator L remain the same as in the previous section: M is complete with bounded geometry and L is adapted and weakly coercive. The additional assumptions, that is, the presence of a Φ -chain (depending on the function Φ) or even of an underlying hyperbolic geometry (with constant δ and coming with a universal function $\Phi = \Phi_\delta$), are stated directly in the results.

5.1 Global Behavior: Φ -Chains

The following result describes the key feature of Φ -chains in the potential theory of our elliptic operators.

Theorem 5.1 (Green's Functions Along Φ -Chains) *Under assumptions (BAC), there is a suitable constant $c(\sigma, \ell, k, n, \tau, \Phi) > 1$ such that for any Φ -chain with track points x_1, \dots, x_m we have for the minimal Green's functions*

$$c^{-1}G(x_m, x_j)G(x_j, x_1) \leq G(x_m, x_1) \leq cG(x_m, x_j)G(x_j, x_1), \quad j = 2, \dots, m-1.$$

At the heart of the argument we employ the pairing of two at first sight entirely unrelated geometric and analytic properties: the existence of Φ -chains on M and the weak coercivity of L . The idea is that, on the one hand, Φ -chains allow to find balls of arbitrary large radii in $U_{i-1} \setminus U_{i+1}$ centered in ∂U_i within a uniformly upper bounded distance to the track points. On the other hand, the [relative maximum principle 4.5](#) shows that on these balls we can improve estimates we have diminished from a previous application of a Harnack inequality. This makes the following result the main step in the proof of the Theorem.

Proposition 5.2 (Growth Recovery Along Φ -Chains) *With assumptions (BAC), for any given Φ -chain with track points x_1, \dots, x_m we have for the minimal Green's functions*

$$G(z, x_1) \leq cG^\theta(z, x_j)G(x_{j+1}, x_1) \quad \text{for } z \in \partial U_{j+1}, \quad (3)$$

for some constant $c(\sigma, \ell, k, n, \tau, \Phi) > 0$ independent of the length j of the Φ -chain.




Figure 3: Growth Recovery Mechanism

Proof. The argument is by induction over the length j . For $j = 1$, the result follows from $G^\theta \geq G$, inequality (1) and the [lower bound for the Green's function 4.2](#), $c_1 G(x_2, x_1) \geq 1$, from where we can take the first guess for the constant $c = c_1$ and note that c_1 depends only on σ, ℓ, k, τ and Φ . For the induction step we first assume we have proved the weaker assertion that there is a constant c_j so that (3) holds for any Φ -chain of length j . Then we can apply the Harnack inequalities for L and L_θ^* to get a constant $c'(\sigma, \ell, k, \tau, \Phi) \geq 1$, independent of j , such that

$$G(z, x_1) \leq c'c_j G^\theta(z, x_{j+1})G(x_{j+2}, x_1) \quad \text{for } z \in \partial U_{j+1}. \quad (4)$$

By the [global maximum principle 2.2](#) this inequality extends to $z \in \bar{U}_{j+1}$. Now we invest the weak coercivity of L and the properties of the Φ -chains to improve this inequality.

Towards this end, we first apply the [relative maximum principle 4.5](#) to the (L -superharmonic) function $G^\theta(\cdot, x_{j+1})$ and its greatest L -harmonic minorant u on some ball $B_R(x)$ which we can represent as $u = \mathcal{R}_{G^\theta(\cdot, x_{j+1})}^{\partial B_R(x)}$, the redit always taken w.r.t. L . For $R = \ln(1/c')/\ln \eta$ and $B_{R+3}(x) \subset U_{j+1}$, the relative maximum principle yields

$$u(x) \leq \frac{1}{c'} G^\theta(x, x_{j+1}) .$$

In turn, the definition of a Φ -chain shows that there is some $\Delta(\Phi, R) > 0$ such that $B_{R+3}(x) \subset U_{j+1}$, for $x \in \partial U_{j+2}$, as soon as $d(x, x_{j+2}) \geq \Delta$. Then, we have from (4)

$$\begin{aligned} G(x, x_1) &= \mathcal{R}_{G(\cdot, x_1)}^{\partial B_R(x)}(x) \leq c' c_j \mathcal{R}_{G^\theta(\cdot, x_{j+1})}^{\partial B_R(x)}(x) G(x_{j+2}, x_1) \\ &= c' c_j u(x) G(x_{j+2}, x_1) \leq c_j G^\theta(x, x_{j+1}) G(x_{j+2}, x_1) . \end{aligned}$$

On the other hand, we get universal estimates for $x \in \partial U_{j+2}$ with $d(x, x_{j+2}) < \Delta$: Since $d(x, x_{j+2}) < \Delta$, there are constants $c'', c''' \geq 1$ only depending on Δ, Φ, H and c_1 , such that $G(x, x_1) \leq c'' G(x_{j+2}, x_1)$ by Harnack inequalities and $G^\theta(x, x_{j+1}) \geq (c''')^{-1}$ by the [bounds for the Green's function 4.2](#) and Harnack inequalities. The result is

$$G(x, x_1) \leq c'' c''' G^\theta(x, x_{j+1}) G(x_{j+2}, x_1) \quad \text{for } x \in \partial U_{j+2} \text{ with } d(x, x_{j+2}) < \Delta .$$

Everything combined, we can choose $c = \max\{c_1, c'' c'''\}$ and outside a tube of radius Δ around the track points the constant can be kept in every induction step while on the inside we have a universal constant. \square

Proof of Theorem 5.1 “Green's functions along Φ -chains”. The first inequality is rather easy: For $x \in \partial B_\sigma(x_j)$ we have

$$G(x, x_j) G(x_j, x_1) \leq c_1 G(x_j, x_1) \leq c_1 H G(x, x_1)$$

by [Proposition 4.2 “bound for the Green's function”](#) and the [Harnack inequalities 3.5](#). Since the left hand side is a potential and the right hand side is L -superharmonic, this inequality extends to $M \setminus B_\sigma(x_j)$ and in particular to x_m by the [global maximum principle 2.2](#).

For the second inequality we use repeatedly [Proposition 5.2 “growth recovery along \$\Phi\$ -chains”](#) and the [resolvent equation 4.1](#):

$$\begin{aligned} G(x_m, x_1) &= \mathcal{R}_{G(\cdot, x_1)}^{\partial U_j}(x_m) && | x_1 \notin U_j \\ &\stackrel{(3)}{\leq} c \mathcal{R}_{G^\theta(\cdot, x_j)}^{\partial U_j}(x_m) G(x_{j+1}, x_1) \\ &= c \mathcal{R}_{G(\cdot, x_j) + \theta G(G^\theta(\cdot, x_j))}^{\partial U_j}(x_m) G(x_{j+1}, x_1) && | \text{ res.eq.} \\ &\leq c \left(G(x_m, x_j) + \theta \int_M \mathcal{R}_{G(\cdot, z)}^{\partial U_j}(x_m) G^\theta(z, x_j) \, d\text{Vol}(z) \right) G(x_{j+1}, x_1) \end{aligned} \quad (5)$$

At this point we can again employ the first step (3), but now for the reversed Φ -chain x_m, \dots, x_1 with $M \setminus U_m, \dots, M \setminus U_1$ and for the adjoint operator L^* , namely

$$G(x_m, z) \leq c G^\theta(x_{j+2}, z) G(x_m, x_{j+1}) \quad \text{for } z \in M \setminus U_{j+1} .$$

This holds on all of $M \setminus U_{j+1}$ by the [global maximum principle 2.2](#). Since $x_j \in M \setminus U_{j+1}$, this can be directly applied to $G(x_m, x_j)$. For the second summand in (5), we have $\mathcal{R}_{G(\cdot, z)}^{\partial U_j}(x_m) =$

${}^*\mathcal{R}_{G(x_m, \cdot)}^{\partial U_j}(z) \leq G(x_m, z)$ for $z \in \partial U_j \subset M \setminus U_{j+1}$ (denoting the reduit with respect to L^* by ${}^*\mathcal{R}$, cf. subsection 2.3), but the upper bound ${}^*\mathcal{R}_{G(x_m, \cdot)}^{\partial U_j}(z) \leq cG^\theta(x_{j+2}, z)G(x_m, x_{j+1})$ is valid for *all* $z \in M$ by the definition of the reduit since the right hand side is positive and L^* -superharmonic in z . Thus,

$$G(x_m, x_1) \leq c^2 G(x_m, x_{j+1}) G(x_{j+1}, x_1) \left(G^\theta(x_{j+2}, x_j) + \theta \int_M G^\theta(x_{j+2}, z) G^\theta(z, x_j) \, d\text{Vol}(z) \right).$$

The large bracket is universally bounded by Proposition 4.2 “bound for the Green’s function”, the Harnack inequalities 3.5, and the inequalities (1) and (2) following from the resolvent equation 4.1 for $t = \frac{3}{2}\theta$. \square

Now we assume that M is a δ -hyperbolic space, then we can choose $\Phi = \Phi_\delta$ and recall that there are Φ -chains along geodesics in M . Since Φ_δ is determined from δ , the Φ -dependence of the estimates now reduces to a δ -dependence.

Corollary 5.3 (Green’s Function Along Hyperbolic Geodesics) *Assuming (BAC) for (M, L) , M δ -hyperbolic, let $x, y, z \in M$ such that y lies on geodesic connecting x and z with $d(x, y), d(y, z) > 22\delta$. Then there is a constant $c(\sigma, \ell, k, n, \tau, \delta) > 1$ such that*

$$c^{-1}G(x, y)G(y, z) \leq G(x, z) \leq cG(x, y)G(y, z).$$

5.2 Boundary Harnack Inequality

Using Theorem 3.6 “boundary Harnack inequality on a disc” as a blueprint, we want to formulate a boundary Harnack inequality near points on the Gromov boundary of a δ -hyperbolic space.

As a replacement for balls in the classical version of the boundary Harnack inequality, we need some characterization of neighborhoods of a point at infinity. This is made precise by the notion of Φ -neighborhood bases.

Definition 5.4 (Φ -Neighborhood Basis) We call two open subsets $V \supset W$ of the space M the **Φ -neighborhoods with hub** $h \in M$, if $\overline{W} \subset V$, $B_{\Phi_0}(h) \subset V \setminus \overline{W}$ and any two points $p \in \partial V$ and $q \in \partial W$ can be joined by a Φ -chain that has h as a track point. We call an infinite family of nonempty open $\mathcal{N}_i \subset M$, $i = 1, 2, 3, \dots$, with $\bigcap_i \mathcal{N}_i = \emptyset$ a **Φ -neighborhood basis**, if \mathcal{N}_i and \mathcal{N}_{i+1} are Φ -neighborhoods with hub p_i , for every i .

Just as metric balls are, besides their rôle as neighborhood bases, the basic playground for Harnack inequalities, these Φ -neighborhoods are their counterpart in *boundary* Harnack inequalities.

In δ -hyperbolic spaces, every point in the Gromov boundary has a canonical neighborhood basis that is also a Φ_δ -neighborhood basis. Namely, as shown in [BHK01, Proposition 8.10], we have

Lemma 5.5 (Φ_δ -Neighborhood Basis) *For a δ -hyperbolic manifold M , $z \in \partial_G M \subset \overline{M}^G$ and $\gamma : \mathbb{R} \rightarrow M$ a geodesic so that z is its endpoint for $t \rightarrow -\infty$ we set*

$$\mathcal{N}_i^\delta := \{x \in X \mid \text{dist}(x, \gamma([300 \cdot i, +\infty))) < \text{dist}(x, \gamma((-\infty, 300 \cdot i]))\}$$

Then the \mathcal{N}_i^δ are a Φ_δ -neighborhood basis and their closures $\overline{\mathcal{N}_i^\delta} \subset \overline{M}^G$ a neighborhood basis of $z \in \partial_G M$.

For non-self-adjoint operators (which will be important for the applications in section 6) it is not always possible to find L -harmonic functions that vanish at infinity, because even minimal Green’s functions might diverge. Hence we introduce a more general notion which can be thought of as a minimal growth condition. This will be further explained in Proposition 5.15 “ L -vanishing and Martin boundary” below.

Definition/Proposition 5.6 (L-Vanishing) We say that a positive L -superharmonic function u **L-vanishes** towards infinity on a domain $V \subset M$ if one of the following equivalent conditions is satisfied:

- (i) There is a positive L -superharmonic function w , such that $u/w \rightarrow 0$ at infinity, i.e., for every $\varepsilon > 0$ there is a compact set $K \subset M$ with $u/w < \varepsilon$ on $V \setminus K$.
- (ii) There is a potential p such that $p \geq u$ on V .
- (iii) The redit \mathcal{R}_u^V is a potential.³

Proof. (i) \Rightarrow (iii): Assume there is a positive L -harmonic function h , such that $\mathcal{R}_u^V \geq h > 0$ on M . For some fixed $\varepsilon > 0$, choose a compact set K with $u < \varepsilon w$ on $V \setminus K$. By the properties of the redit, we even have

$$\varepsilon w \geq \mathcal{R}_u^{V \setminus K} \geq \mathcal{R}_u^V - \mathcal{R}_u^{V \cap K} \geq h - \mathcal{R}_u^{V \cap K}$$

on all of M . Now $\mathcal{R}_u^{V \cap K}$ is a potential since $V \cap K$ is relatively compact in M , $\varepsilon w - h$ is L -superharmonic and $\varepsilon w \geq u = \mathcal{R}_u^V \geq h$ on $\partial(V \setminus K) = \partial(M \setminus (V \setminus K)) \subset M$, thus we can apply the [global maximum principle 2.2](#) to see $\varepsilon w \geq h$ on all of M . Since ε was arbitrary, $h = 0$.

(iii) \Rightarrow (ii): Choose $p = \mathcal{R}_u^V$.

(ii) \Rightarrow (i): It suffices to show that a potential p satisfies the condition everywhere. Towards this end, consider the functions $\mathcal{R}_p^{M \setminus \overline{B_R(x_0)}}$ on balls around an arbitrary basepoint $x_0 \in M$. They converge to zero for $R \rightarrow \infty$ because the limit is L -harmonic and $\leq p$. Let (x_j) be a countable dense set in M . We may choose a sequence (R_i) such that $\mathcal{R}_p^{M \setminus \overline{B_{R_i}(x_0)}}(x_j) \leq 2^{-i}$ for all $j \leq i$. Then the function $w = \sum_i \mathcal{R}_p^{M \setminus \overline{B_{R_i}(x_0)}}$ is finite on a dense set, L -superharmonic and $p/w \leq 1/i$ outside of $B_{R_i}(x_0)$. \square

Note that all potentials such as the minimal Green's function are L -vanishing on $V = M$ and hence on all open sets because the property of L -vanishing is conserved on subsets as can be easily seen using condition (i).

On bounded sets, L -vanishing *at infinity* is trivially true for any L -superharmonic function. The concept is also useless for unbounded sets shrinking to a small set, e.g. a single point, when approaching infinity. It becomes significant for $V = W \cap M$ where $W \subset \overline{M}^G$ is open with non-empty intersection $W \cap \partial_G M$. For more elaborate criteria in the context of Martin theory see [Proposition 5.15 "L-vanishing and Martin boundary"](#).

On Φ -neighborhoods we can now formulate the following central result.

Theorem 5.7 (Boundary Harnack Inequality) Assume (M, L) satisfies assumptions [\(BAC\)](#). Let $V \supset W$ be Φ -neighborhoods with hub h and u, v two positive L -superharmonic functions that are L -harmonic and L -vanishing on V , then there is a constant $H_B = H_B(\sigma, \ell, k, n, \tau, \Phi)$ such that

$$\frac{u(x)}{u(y)} \leq H_B \frac{v(x)}{v(y)} \quad \text{for any } x, y \in W .$$

Proof. By [Definition/Proposition 5.6 "L-vanishing"](#), the redit \mathcal{R}_u^V is a potential and therefore admits a representation as

$$\mathcal{R}_u^V = \int_{\partial V} G(\cdot, z) d\nu(z)$$

³[\[Anc87\]](#) uses the first definition, but it is easier to employ the last.

for some Radon measure ν . On \overline{W} , \mathcal{R}_u^V agrees with u and therefore we have

$$u(x) = \int_{\partial V} G(x, z) d\nu(z) \leq c \int_{\partial V} G(x, h) G(h, z) d\nu(z) = c G(x, h) u(h) \quad \text{for } x \in \partial W$$

using the assumption that $x \in \partial W$ and $z \in \partial V$ can be connected by a Φ -chain through h and [Theorem 5.1 “Green’s functions along \$\Phi\$ -chains”](#). The other inequality from there gives

$$v(x) \geq c^{-1} G(x, h) v(h) \quad \text{for } x \in \partial W$$

and both inequalities extend to W by the [global maximum principle 2.2](#) because \mathcal{R}_u^V and $G(\cdot, h)$ respectively are potentials. We can combine them to obtain

$$\frac{u(x)}{v(x)} \leq c^2 \frac{u(h)}{v(h)} \quad \text{for } x \in W .$$

Interchanging the rôles of u and v yields the result with $H_B = c^4$. □

In most applications, u and v are either globally L -harmonic functions or minimal Green’s functions with pole outside of V .

In the case of a δ -hyperbolic manifold the size of the smaller neighborhood can be explicitly quantified using [Lemma 5.5 “ \$\Phi_\delta\$ -neighborhood basis”](#). We get

Corollary 5.8 (Hyperbolic Boundary Harnack Inequality) *If (M, L) satisfies (BAC) and M is δ -hyperbolic, there is some positive constant $H_B(\sigma, \ell, k, n, \tau, \delta) > 1$ such that two positive L -superharmonic functions u, v that are L -harmonic and L -vanishing on a Φ_δ -neighborhood \mathcal{N}_i^δ of $\xi \in \partial_G M$ satisfy*

$$\frac{u(x)}{u(y)} \leq H_B \frac{v(x)}{v(y)} \quad \text{for any } x, y \in \mathcal{N}_{i+1}^\delta .$$

5.3 The Martin Boundary

In the previous section we have already formulated *boundary* Harnack inequalities on the complete manifold M although they neither needed nor referred to any concrete boundary. The natural boundary concept related to these boundary inequalities we introduce here is the Martin boundary. These ideal boundaries can equally be defined on non-complete manifolds like Euclidean domains.

The Martin boundary is determined from both the geometry of M and the analysis of L . A basic problem in Martin theory is the characterization of this boundary with a minimal amount of analytic input. The distinguished result in our case is that when we also assume Gromov hyperbolicity, then the Martin boundary actually only depends on the geometry. It is homeomorphic to the Gromov boundary $\partial_G M$ of M .

We first recall some basic notions from Martin theory, cf. [\[Anc90\]](#) or [\[Pin95, 7.1\]](#).

Martin Boundary For a non-compact Riemannian manifold M and a linear second order elliptic operator L on M with a minimal Green’s function $G : M \times M \rightarrow (0, \infty]$ as well as a basepoint $p \in M$, we consider the space S of sequences s of points $p_i \in M$, $i = 1, 2, \dots$ with

- $p_i \in M$ has no accumulation points in M ,
- $K_{p_i} := G(\cdot, p_i) / G(p, p_i) \xrightarrow{n \rightarrow \infty} K_s$ compactly, for some function K_s .

The **Martin boundary** $\partial_M(M, L)$ is the quotient of S modulo the relation $s \sim s^*$ if $K_s \equiv K_{s^*}$. For $\zeta \in \partial_M(M, L)$, this function is written K_ζ and called the **Martin kernel**.


The Martin boundary does not depend on the choice of the basepoint p . The Harnack inequality and elliptic theory show that each $K_\zeta \in \partial_M(M, L)$ is a positive solution of $Lu = 0$ on M . This also shows that the convex set $S_L(M)$ of positive solutions of $Lu = 0$ on M with $u(p) = 1$ is compact in the topology of compact convergence. In turn, $\partial_M(M, L)$ is a compact subset of $S_L(M)$.

The **Martin topology** on $\overline{M}^M := M \cup \partial_M(M, L)$ is the topology of compact convergence on the space of associated Martin functions $\{K_y \mid y \in \overline{M}^M\}$.

Then the usual topology is induced on $M \subset \overline{M}^M$, $\partial_M(M, L)$ is closed and \overline{M}^M is compact. The space \overline{M}^M is called the **Martin compactification**. It is easy to see that \overline{M}^M is metrizable, cf. [BJ06, Ch. I.7] or [Hel14, Ch. 12] for further details.

To motivate the idea of Martin integrals we recall the following classical result, cf. [Cho69, Ch. 6]:

Proposition 5.9 (Minkowski's Theorem) *Each point in a convex set $K \subset \mathbb{R}^n$ is a convex combination of the extremal points of K .*



Convex domains and Extremal points

Each point in a triangle is a **unique** convex sum of its **affine independent** vertices. For general convex sets, inner points usually have infinitely many different sum representations of its extremal points. For the set G , we have

$$\begin{aligned} x &= 0.5 x_1 + 0.5 (0.2 x_2 + 0.8 x_4) \\ &= 0.6 x_2 + 0.4 x_5 \end{aligned}$$

Figure 4: Minkowski's Theorem in \mathbb{R}^2

The **Martin integral** is essentially an extension of this result to the case of the convex set $S_L(M)$. The extremal elements of $S_L(M)$ form a subset $\partial_M^0(M, L) \subset \partial_M(M, L)$ of the Martin boundary one can think of as the vertices of an infinite dimensional simplex spanning $S_L(M)$. A positive solution u of $Lu = 0$ on M with $u(p) = 1$ is extremal if and only if u is a minimal solution. Here we call u **minimal** if for any other solution $v > 0$, $v \leq u$, we have $v \equiv c \cdot u$, for some constant $c > 0$. Therefore $\partial_M^0(M, L) \subset \partial_M(M, L)$ is also called the **minimal Martin boundary**.

The Choquet integral representations in [Cho69, Ch. 6] give the following general version of the Martin representation theorem, cf. [Pin95, 7.1]:

Proposition 5.10 (Martin Integral) *For any positive solution u of $Lu = 0$ on M , there is a unique Radon measure μ_u on $\partial_M^0(M, L)$ so that*

$$u(x) = \int_{\partial_M^0(M, L)} K_\zeta(x) d\mu_u(\zeta) .$$

Conversely, for any Radon measure μ on $\partial_M^0(M, L)$,

$$u_\mu(x) = \int_{\partial_M^0(M, L)} K_\zeta(x) d\mu(\zeta)$$

defines a positive solution of $Lu_\mu = 0$ on M .

Although this already looks like a classical contour integral the result is not yet truly satisfactory. Unlike the classical case the boundary $\partial_M^0(M, L)$ depends not only on the underlying space but also on the analysis of the operator L . A natural question is whether one could get rid of this dependence. In general the answer is no. In 6.6 we will see some by no means exotic examples. However, we will now see that in our case of operators on δ -hyperbolic spaces, this is actually possible. This is a remarkable particularity not even valid for such simple spaces as in [Example 5.14 “Ideal Boundaries of \$\mathbb{H}^m \times \mathbb{H}^n\$ ”](#).

Back to hyperbolic spaces In the situation at hand, Φ -neighborhood bases are essentially neighborhood bases of minimal Martin boundary points.

Theorem 5.11 (Characterization of Minimal Martin Points) *Assume (BAC) holds. Let (\mathcal{N}_i) be a Φ -neighborhood basis with hubs p_i . Denoting the interior of the closure of $\mathcal{N}_i \subset M \subset \overline{M}^M$ in the Martin compactification \overline{M}^M of M by $\tilde{\mathcal{N}}_i$, there is exactly one Martin boundary point ζ in $\bigcap \tilde{\mathcal{N}}_i$. The resulting K_ζ is characterized as the only positive L -harmonic function L -vanishing on every $M \setminus \mathcal{N}_i$ except for scalar multiples. In particular, this Martin point is minimal.*

Proof. By the Harnack inequalities, the sequence $K_{p_i} = \frac{G(\cdot, p_i)}{G(p, p_i)}$ has a subsequence compactly converging to some L -harmonic function K_ζ representing a Martin boundary point ζ . K_ζ is L -vanishing on every $M \setminus \mathcal{N}_i$ because by the [boundary Harnack inequality 5.7](#), every K_{p_j} for $j \geq i$, and hence the limit, is upper bounded by the potential $H_B K_{p_i}$ on $M \setminus \mathcal{N}_i$.

Now assume there is another positive L -harmonic function u that is L -vanishing on $M \setminus \overline{\mathcal{N}}_i$ for every i , w.l.o.g. $u(p) = 1$. Applying the [boundary Harnack inequality 5.7](#) we see $H_B^{-1} K_\zeta \leq u \leq H_B K_\zeta$ on M . Thus $\eta := \inf u/K_\zeta \leq 1$ is positive. By the strong maximum principle [[Pin95](#), Theorem 3.2.6, p.84], the L -harmonic function $u - \eta K_\zeta \geq 0$ has to be positive everywhere, else it would be identical zero. In the former case we can again apply the [boundary Harnack inequality 5.7](#) to K_ζ and $u - \eta K_\zeta$ to get $(\eta + (1 - \eta)H_B^{-1})K_\zeta \leq u$ which contradicts the definition of η , unless $\eta = 1$ and $u = K_\zeta$. \square

We have the following characterization of the Martin boundary in case we have enough Φ -chains:

Corollary 5.12 (Identifying the Martin Boundary) *If under assumptions (BAC) in a given compactification \overline{M} of M (i.e., \overline{M} is compact and $M \subset \overline{M}$ dense) every boundary point admits a neighborhood basis of the form $\overline{\mathcal{N}}_i \subset \overline{M}$ for some Φ -neighborhood basis (\mathcal{N}_i) , it is canonically homeomorphic to the Martin compactification \overline{M}^M .*

Proof. [Theorem 5.11 “characterization of minimal Martin points”](#) yields an injective map from \overline{M} to \overline{M}^M . It is continuous because for every sequence (y_i) in \overline{M} converging to $\zeta \in \overline{M} \setminus M$ the corresponding Martin functions K_{y_i} converge to the unique Martin function that is L -vanishing on all $M \setminus \mathcal{N}_i$ for some Φ -neighborhood basis (\mathcal{N}_i) of ζ , that is K_ζ . Thus, by elementary properties of compactifications, it is already a homeomorphism. \square

From this and [Lemma 5.5 “ \$\Phi_\delta\$ -neighborhood basis”](#) we get the following principal potential theoretic result on Gromov hyperbolic manifolds.

Corollary 5.13 (Gromov Boundary and Martin Boundary) *Assume that M is Gromov hyperbolic and (BAC) holds. Then the Gromov and Martin boundaries of M are canonically homeomorphic and every Martin boundary point is already minimal,*

$$\partial_G M \cong \partial_M(M, L) \cong \partial_M^0(M, L).$$

In particular, some function $u > 0$ on M solves $Lu = 0$ if and only if there is a (unique) Radon measure μ_u on $\partial_{\mathbb{G}}M$ such that

$$u(x) = \int_{\partial_{\mathbb{G}}M} K_{\zeta}(x) d\mu_u(\zeta).$$

The validity of such a simple identification of the Martin boundary with a geometric boundary, which one may possibly expect from a naive guess, actually is a rare exception. If we only slightly violate the hyperbolicity constraint we can get a completely different and rather inscrutable outcome with many non-minimal Martin boundary points.

Example 5.14 (Ideal Boundaries of $\mathbb{H}^m \times \mathbb{H}^n$) The product space of two classical hyperbolic spaces $\mathbb{H}^m \times \mathbb{H}^n$, $m, n \geq 2$, has bounded geometry but it is no longer Gromov hyperbolic since it contains flat planes obtained as products of pairs of geodesics in the two factors. In [GW93] we find a thorough discussion of the case of the Laplace operator. There exist positive functions h with $-\Delta h = \lambda \cdot h$ if and only if $\lambda \leq \lambda_0$ where λ_0 is the generalized principal eigenvalue. For $\lambda < \lambda_0$, the operator $-\Delta - \lambda \text{Id}$ is adapted and weakly coercive. The boundary at infinity (a natural generalization of Gromov boundary) is homeomorphic to S^{n+m-1} , cf. [BH99, II.8.11(6), p. 266]. In turn, for the minimal Martin boundary of $-\Delta - \lambda \text{Id}$, for $\lambda < \lambda_0$, we have

$$\partial_{\mathbb{M}}^0(\mathbb{H}^m \times \mathbb{H}^n, -\Delta - \lambda \text{Id}) = S^{m-1} \times S^{n-1} \times I_{\lambda},$$

where I_{λ} is a closed interval with a natural parameterization which depends on λ and which degenerates to a single point when $\lambda \rightarrow \lambda_0$ [GW93, p. 21]. The full Martin boundary contains two additional pieces

$$\partial_{\mathbb{M}}(\mathbb{H}^m \times \mathbb{H}^n, -\Delta - \lambda \text{Id}) = (S^{m-1} \times \mathbb{H}^n) \cup (S^{m-1} \times S^{n-1} \times I_{\lambda}) \cup (\mathbb{H}^m \times S^{n-1}) / \sim. \quad (6)$$

The gluing maps for \sim are described in [GW93, p. 27]. We observe that not only the boundary at infinity does not coincide with $\partial_{\mathbb{M}}^0(\mathbb{H}^m \times \mathbb{H}^n, -\Delta - \lambda \text{Id})$ but the details of the partition of the full Martin boundary (6) depend on λ . \square

Now that we have the right notion for a potential theoretic boundary at infinity, there are some new formulations for L -vanishing. Note that these are slightly different from the formulation in Definition/Proposition 5.6 “ L -vanishing” because there it was only possible to refer to L -vanishing *at infinity of open sets V in M* , i.e., on the Martin boundary points in $\overline{V} \cap \partial_{\mathbb{M}}M \subset \overline{M}^{\mathbb{M}}$.

Proposition 5.15 (L -Vanishing and Martin Boundary) *Assume (BAC) holds and every Martin boundary point has a Φ -neighborhood basis, e.g., M is Gromov hyperbolic. For an open subset $\Xi \subset \partial_{\mathbb{M}}M$ of the Martin boundary and a positive L -harmonic function u on M the following are equivalent:*

- (i) u is L -vanishing on any open set $V \subset M$ with $\overline{V} \cap \partial_{\mathbb{M}}M \subset \Xi$ in the Martin compactification.
- (ii) On any open set $V \subset M$ with $\overline{V} \cap \partial_{\mathbb{M}}M \subset \Xi$ in the Martin compactification, the following property holds: Each positive L -harmonic function v on V with $v \geq u$ on $\partial V \cap M$ satisfies $v \geq u$ on V .
- (iii) The Martin measure μ_u associated to u is supported outside Ξ , i.e., $\mu_u(\Xi) = 0$.

Proof. (i) \Rightarrow (ii) If u is L -vanishing on V , \mathcal{R}_u^V is a potential and $\mathcal{R}_u^V = u$ on V . Then the global maximum principle 2.2 gives exactly the desired property.

(iii) \Rightarrow (i) Each Martin function K_{ζ} is L -vanishing outside ζ and, therefore, the Martin integral representing u L -vanishes outside the support of μ_u .

(ii) \Rightarrow (iii) Assume $\mu_u(\Xi) \neq 0$, then there is a compact $K \subset \partial_M M$ and an open $W \subset \overline{M}^M$ such that $K \subset W \cap \partial_M M \Subset \Xi$, $V := W \cap M$ satisfies the condition in (ii), and $\mu_u(K) \neq 0$ (since the Radon measure μ_u is inner regular). Therefore it is enough to consider the case where $u \equiv u_K := \int_K K_\zeta d\mu_u(\zeta)$.

We compare u_K with the minimal Green's function $G(\cdot, p)$ with pole $p \in M \setminus V$. Recalling the argument of (iii) \Rightarrow (i) we know that u_K is L -vanishing on an open neighborhood N of $M \setminus V$. By compactness of $\overline{M \setminus V}$ in the Martin compactification, $M \setminus V$ can be covered by finitely many Φ -neighborhoods contained in N and a compact subset of M . Then the **boundary Harnack inequality 5.7** shows that there is a $C > 0$ such that $C \cdot G(\cdot, p) \geq u_K$ on $M \setminus V$ and especially on ∂V . But then from (ii) it follows that $C \cdot G(\cdot, p) \geq u_K$ on V , hence on all of M , hence $u_K \equiv 0$ because $G(\cdot, p)$ is a potential. \square

In the next section we will see that the integral representation we have obtained in the Gromov hyperbolic case naturally extends classical contour integrals for (positive) harmonic functions on the subclass of Euclidean domains which are *uniform*.

6 Geometry and Analysis on Uniform Domains

Here we consider domains sharing basic geometric and analytic properties with smoothly bounded ones. Examples are canonical deformations of such domains to Gromov hyperbolic manifolds of bounded geometry but also the validity of Poincaré-Sobolev type inequalities and embedding theorems. We start with the definition and examples of so-called uniform domains. Then we explain how one can transfer Ancona's theory from hyperbolized uniform domains back to original Euclidean domains to derive new analytic results and to recover classical integral formulas.

6.1 Uniform Domains

The path connectedness of an open set is a topological condition. For finer geometric and analytic investigations one seeks for a quantitative form of path connectedness. One of the nowadays central notions is described in the following definition. We refer to [Aik12] for an instructive overview and comparison of other regularity concepts for domains.

Uniform Domains *A domain $D \subset \mathbb{R}^n$ is called a **uniform domain**, more precisely, a c -uniform domain, if there is a $c \geq 1$ such that any two points $p, q \in D$ can be joined by a **c -uniform curve**. That is a rectifiable path $\gamma : [a, b] \rightarrow D$, for some $a < b$, from p to q so that the following conditions are satisfied:*

- **Quasi-geodesic:** $l(\gamma) \leq c d(p, q)$.
- **Twisted double cones:** $\min\{l(\gamma|_{[a,t]}), l(\gamma|_{[t,b]})\} \leq c \operatorname{dist}(\gamma(t), \partial D)$ for any $t \in [a, b]$.

Note that being c -uniform is a scaling invariant condition: whenever D is c -uniform, $\lambda \cdot D$, $\lambda > 0$, is also c -uniform.

Examples 6.1 (Uniform and Non-Uniform Domains) We start with some types of Euclidean domains which are uniform:


- Any bounded domain with smooth or at least Lipschitz boundary is uniform.
- Non-compact rotationally symmetric domains bounded with profile functions of at least linear growth are uniform. As an explicit example, choose $f(t) = c_1 \cdot t + c_2$, for constants $c_i > 0$ and let $F : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, $n \geq 3$, be given by $F(x) := f(|x|)$. Consider the domain

$$D_f := \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid |x_1| < F(x_2, \dots, x_n)\}.$$

To explain the idea of how to define the quasi-geodesics and twisted double cones joining, e.g., the pairs of points $p_k^\pm = (0, \dots, 0, \pm k)$, $k \geq 1$, we note that for $k = 1$ we can choose any such twisted double cone $\subset D_f$ along a half circle in the $n - 1, n$ -plane joining p_1^- and p_1^+ . Then the quasi-geodesic and twisted double cone scaled by k serve for the p_k^- and p_k^+ . Here it is essential that f grows at least linearly to ensure that the twisted double cone remains in D_f .

- Bounded domains with certain types of fractal boundaries, like the Snowflake in \mathbb{R}^2 or the complement of the Sierpinski gasket in \mathbb{R}^3 [ALM03], are uniform domains.

Non-Uniform Domains



Uniform Domains




Figure 5: Typical (Non-)Uniform Domains

However, even completely harmless domains can be non-uniform:

- The difference of the cube $(-1, 1)^n \subset \mathbb{R}^n$ minus the ball $B_1^n(0) \subset \mathbb{R}^n$: $(-1, 1)^n \setminus B_1(0)$ is not a uniform domain, since we cannot reach points arbitrarily near to the boundary point $(0, \dots, 0, 1) \in \partial((-1, 1)^n \setminus B_1(0))$ by twisted cones, for some common constant $c > 0$.
- The cylinder $B_1^{n-1}(0) \times \mathbb{R} \subset \mathbb{R}^n$ and similarly $B_1^k(0) \times \mathbb{R}^{n-k} \subset \mathbb{R}^n$, for $1 \leq k \leq n - 1$ are non-uniform. For the pairs of points $p_k^\pm = (0, \dots, 0, \pm k)$, $k \geq 1$, there are only twisted double cones reaching from p_k^+ to p_k^- , for $c = c(k) \leq 1/k$.
- Similarly, the domain $\mathbb{R}^3 \setminus \mathbb{R} \times \mathbb{Z}^2$ is not uniform.

- As in the earlier example of non-compact rotationally symmetric domains D_f we choose profile functions but this time of sublinear growth like $f(t) = c_1 \cdot \sqrt{t} + c_2$, for constants $c_i > 0$. Then, as in the last counterexample, D_f is non-uniform.

6.2 Quasi-Hyperbolic Geometry

On any domain $D \subset \mathbb{R}^n$ we can define the *quasi-hyperbolic metric* k_D introduced as an extension of the definition of the Poincaré metric on the unit disc to arbitrary domains. It is given by

$$k_D(x, y) := \inf \left\{ \int_{\gamma} 1/\bar{d} \mid \gamma \subset D \text{ rectifiable curve joining } x \text{ and } y \right\}$$

where we set $\bar{d} = \text{dist}(\cdot, \partial D)$. This corresponds to a conformal deformation of g_{Eucl} to the merely Lipschitz continuous $\bar{d}^{-2}g_{\text{Eucl}}$. The quasi-hyperbolic metric k_D always has bounded geometry since the chart

$$\text{Id} : (B_{\bar{d}(x)/2}(x), k_D) \rightarrow (B_{\bar{d}(x)/2}(x), \bar{d}^{-2}(x)g_{\text{Eucl}})$$

(with radii measured in Euclidean distance) is 2-bi-Lipschitz and we see that (D, k_D) has (σ, ℓ) -bounded geometry for some $\sigma, \ell > 0$ both independent of D .

In general, k_D need not to be (Gromov) hyperbolic, except D is a uniform domain. Due to geometric work, in particular of Bonk, Heinonen, Koskela [BHK01], we have:

Theorem 6.2 (Gromov Uniformization) [BHK01, Th. 1.11] *A bounded domain $D \subset \mathbb{R}^n$ is uniform if and only if (D, k_D) is both Gromov hyperbolic and the Euclidean boundary ∂D of D is naturally quasisymmetrically equivalent to the Gromov boundary $\partial_G(D, k_D)$. Moreover, when D is c -uniform, then (D, k_D) is δ_c -hyperbolic of (σ, ℓ) -bounded geometry, for some $\delta_c > 0$ depending only on c .*

A map $f : X \rightarrow Y$ between metric spaces (X, d_X) and (Y, d_Y) is called *quasi-symmetric* if f is not constant and if there is a homeomorphism $\eta : [0, \infty) \rightarrow [0, \infty)$, so that for any $x, a, b \in X$, $t \geq 0$, and $d_X(x, a) \leq t \cdot d_X(x, b)$ it follows that $d_Y(f(x), f(a)) \leq \eta(t) \cdot d_Y(f(x), f(b))$.

Since $\text{dist}(\cdot, \partial D)$ is Lipschitz continuous but not smooth, (D, k_D) is not a Riemannian manifold. But we need some smoothness to ensure the adaptedness of L , in particular to get uniform bounds on the coefficients of the adjoint operator L^* . Thus, in particular for analytic considerations, we will work with smoothed versions of \bar{d} . Fortunately, there is a well-controlled smoothing of \bar{d} :

Theorem 6.3 (Stein-Whitney Smoothing) [Ste70, VI.2.1, p.171] *There are constants $c_{\alpha} = c_{\alpha}(n)$ for every multiindex α such that on every open set $D \subset \mathbb{R}^n$ there is a C^{∞} function δ defined on D with*

$$c_0^{-1} \bar{d} \leq \delta \leq c_0 \bar{d} \quad \text{and} \quad \left| \frac{\partial^{\alpha} \delta}{\partial x^{\alpha}} \right| \leq c_{\alpha} \bar{d}^{1-|\alpha|} .$$

In particular, $(D, \bar{d}^{-2}g_{\text{Eucl}})$ and $(D, \delta^{-2}g_{\text{Eucl}})$ are quasi-isometric and, hence, $(D, \bar{d}^{-2}g_{\text{Eucl}})$ is Gromov hyperbolic if and only if $(D, \delta^{-2}g_{\text{Eucl}})$ is Gromov hyperbolic. Moreover, when $(D, \bar{d}^{-2}g_{\text{Eucl}})$ is δ -hyperbolic, for some $\delta > 0$, then $(D, \delta^{-2}g_{\text{Eucl}})$ is $a_n \cdot \delta$ -hyperbolic, for some $a_n \geq 1$ depending only on n .

6.3 Martin Theory for (Non-)Uniform Domains

Quasi-hyperbolic metrics can be used to transfer the potential theory on Gromov hyperbolic manifolds to understand the corresponding theory on uniform domains. Since there are examples of non-uniform domains which carry a rather complicated potential theory, this correspondence also gives us non-hyperbolic manifolds of bounded geometry with a much less transparent Martin theory than in the hyperbolic case.

On a uniform domain $D \subset \mathbb{R}^n$, we consider a Schrödinger operator $L^{\text{Eucl}} = -\Delta + V$ with a smooth potential V satisfying $|V| \leq a \cdot \delta^{-2}$, for some $a > 0$. That is, we even allow a controlled diverging behavior of V towards ∂D . The methods apply to more general operators, but we only want to explain the idea and this case is particularly important and notationally simple.

On $(D, \delta^{-2}g_{\text{Eucl}})$ we consider the operator $L = \delta^2 \cdot L^{\text{Eucl}}$. Then L is adapted in the bounded geometry charts. Note that L -harmonic functions are the same as L^{Eucl} -harmonic functions, but L is not self-adjoint anymore. At least the adjoint L^* is still adapted.

L is weakly coercive relative to $(D, \delta^{-2}g_{\text{Eucl}})$ provided L^{Eucl} has a positive first eigenvalue not only for L^{Eucl} but also for $\delta^2 \cdot L^{\text{Eucl}}$ in the original Euclidean metric [Anc86]:

Strong Barrier *There is a function $s > 0$ and some $\varepsilon > 0$ such that $L^{\text{Eucl}}s \geq \varepsilon \cdot \delta^{-2} \cdot s$.*

Example 6.4 (Regularity Versus Strong Barrier) The Laplace operator on bounded domains with smooth or Lipschitz boundary admits a strong barrier [Anc86, KK66]. In general, the strong barrier condition is not a consequence of the uniformity of D but of additional exterior conditions. An exterior twisted cone condition is sufficient [Aik12].

Under these mild conditions on L^{Eucl} the potential theory we established above readily transfers to uniform domains where we now use the sets \mathcal{N}_i^δ from Lemma 5.5 “ \mathcal{F}_δ -neighborhood basis”, defined relative $\delta^{-2}g_{\text{Eucl}}$, as a replacement for the concentric balls of Theorem 3.6 “boundary Harnack inequality on a disc” in the classical Euclidean setup.

Corollary 6.5 (Martin Theory on Uniform Domains) *Let $L^{\text{Eucl}} = -\Delta + V$ be an operator with $|V| \leq a \cdot \delta^{-2}$ satisfying a strong barrier condition on some c -uniform domain D , for some $a, \varepsilon, c > 0$. Then we have for any two L^{Eucl} -harmonic functions $u, v > 0$ on $\mathcal{N}_i^\delta \cap D$ both L^{Eucl} -vanishing⁴ towards $\mathcal{N}_i^\delta \cap \partial D$,*

$$u(x)/v(x) \leq C \cdot u(y)/v(y) \text{ for any two points } x, y \in \mathcal{N}_{i+1}^\delta \cap D ,$$

for some $C(a, \varepsilon, c, n) \geq 1$. Thus, the topological and the Martin boundary are homeomorphic and every Martin boundary point is minimal: $\partial D \cong \partial_M^0(D, L^{\text{Eucl}}) \cong \partial_M(D, L^{\text{Eucl}})$.

Proof. We first apply Theorem 6.2 “Gromov uniformization” to the uniform domain D , then we transform $\delta^{-2}g_{\text{Eucl}}$ into the smoothed $\delta^{-2}g_{\text{Eucl}}$ using the Stein-Whitney smoothing 6.3. We get an $\alpha_n \cdot \delta_c$ -hyperbolic manifold of (σ, ℓ) -bounded geometry with

$$\partial D \cong \partial_G(D, \delta^{-2}g_{\text{Eucl}}) \cong \partial_G(D, \delta^{-2}g_{\text{Eucl}}) .$$

Now we can study L , which is weakly coercive and adapted, via the hyperbolic boundary Harnack inequality 5.8 and Corollary 5.13 “Gromov boundary and Martin boundary”. We conclude that

$$\partial D \cong \partial_G(D, \delta^{-2}g_{\text{Eucl}}) \cong \partial_M((D, \delta^{-2}g_{\text{Eucl}}), L) \cong \partial_M^0((D, \delta^{-2}g_{\text{Eucl}}), L) .$$

⁴The definition of L^{Eucl} -vanishing towards $\mathcal{N}_i^\delta \cap \partial D$ is the same as in Definition/Proposition 5.6 “ L -vanishing” and Proposition 5.15 “ L -vanishing and Martin boundary” where we merely had an ideal boundary of M and hence called it L -vanishing towards infinity.

The Green's functions relative L^{Eucl} on (D, g_{Eucl}) and L on $(D, \delta^{-2}g_{\text{Eucl}})$ are related by $G(x, y) = \delta^{n-2}(y)G^{\text{Eucl}}(x, y)$, while the Martin functions and the solutions, along with the notion of L -vanishing, are the same. This means that the boundary Harnack inequalities carry over to D with L^{Eucl} and that $\partial_M^0((D, \delta^{-2}g_{\text{Eucl}}), L) \cong \partial_M^0(D, L^{\text{Eucl}})$ and $\partial_M((D, \delta^{-2}g_{\text{Eucl}}), L) \cong \partial_M(D, L^{\text{Eucl}})$. \square

Aikawa [Aik01, Aik04] has used a somewhat different approach to prove some remarkable refinements underlining the sharpness of these potential theoretic result.

- In [Aik01] he derives boundary Harnack inequalities for the Laplacian on arbitrary uniform domains even *without* imposing a strong barrier condition. However, in his result the Harnack constant depends on the domain D , whereas in the previous result it only depends on the two parameters ε and c .
- In [Aik04] we even find that $D \subset \mathbb{R}^n$ is a uniform domain if and only if the Laplacian satisfies boundary Harnack inequalities. As in Example 6.4 “regularity versus strong barrier” one also needs to assume an exterior regularity of D . Again an exterior twisted cone condition is sufficient [Aik12].

Here we summarize the three essentially **equivalent** properties of a Euclidean domain G discussed above: (i) G is uniform, (ii) G admits a canonical conformal hyperbolic deformation of bounded geometry and (iii) the boundary Harnack inequality for the Laplacian holds on G .




Figure 6: The three essentially **equivalent** properties of a Euclidean domain D

Examples 6.6 We mention two instructive examples where non-uniformity of a domain destroys the existence of boundary Harnack inequalities and where we concretely see how far the topological boundary may deviate from the Martin boundary of the Laplacian.

- In [Anc12] Ancona has described non-uniform Euclidean cones with only one topological point at infinity but with uncountably many minimal Martin boundary points at infinity.
- In [IP94] Ioffe and Pinsky have proved that for the non-uniform rotationally symmetric domains $D_f \subset \mathbb{R}^n$ from Examples 6.1 “uniform and non-uniform domains” the set of Martin boundary points at infinity is homeomorphic to S^{n-2} .

Finally we notice that this Martin theory on uniform domains reproves classical contour integral formulas, for instance, for harmonic functions. The Herglotz theorem [Her11, Dur83] states that a function $f > 0$ on the Euclidean unit disk (D, g_{Eucl}) is harmonic, $\Delta_{\text{Eucl}} f = 0$, if and only if there is a Radon measure μ_f on S^1 such that

$$f(x) = \int_{S^1} \frac{1 - |x|^2}{|x - y|^2} d\mu_f(y) .$$

For the Green’s function G for Δ_{Eucl} , and thus for the Green’s function of the associated hyperbolized manifold and operator, a direct computation shows that

$$K_\zeta(x) = \lim_{z \rightarrow \zeta} \frac{G(x, z)}{G(0, z)} = \frac{1 - |x|^2}{|x - \zeta|^2}, \text{ for any } \zeta \in S^1 .$$

But we also know from the above that we have a unique Martin integral representation:

$$f(x) = \int_{S^1} K_\zeta(x) d\mu_f(\zeta) .$$

μ_f is now understood as a measure on a certain ideal boundary of (D, g_{hyp}) , the Martin boundary, which in this case equals S^1 .

References

- [Aik01] H. AIKAWA, Boundary Harnack principle and Martin boundary for a uniform domain, *J. Math. Soc. Japan* **53** no. 1 (2001), 119–145. doi:10.2969/jmsj/05310119.
- [Aik04] H. AIKAWA, Potential-theoretic characterizations of nonsmooth domains, *Bull. of the Lond. Math. Soc.* **36** no. 4 (2004), 469–482. doi:10.1112/S002460930400308X.
- [Aik12] H. AIKAWA, Potential analysis on nonsmooth domains—Martin boundary and boundary Harnack principle, in *Complex Analysis and Potential Theory, CRM Proc. Lecture Notes* **55**, Amer. Math. Soc., Providence, RI, 2012, pp. 235–253.
- [ALM03] H. AIKAWA, T. LUNDH, and T. MIZUTANI, Martin boundary of a fractal domain, *Potential Analysis* **18** no. 4 (2003), 311–357. doi:10.1023/A:1021823023212.
- [Anc86] A. ANCONA, On strong barriers and an inequality of Hardy for domains in \mathbb{R}^n , *J. Lond. Math. Soc. (2)* **34** no. 2 (1986), 274–290.
- [Anc87] A. ANCONA, Negatively curved manifolds, elliptic operators, and the Martin boundary, *Ann. of Math. (2)* **125** no. 3 (1987), 495–536. doi:10.2307/1971409.
- [Anc90] A. ANCONA, Théorie du potentiel sur les graphes et les variétés, in *École d’été de Probabilités de Saint-Flour XVIII—1988, Lecture Notes in Math.* **1427**, Springer, Berlin, 1990, pp. 1–112. doi:10.1007/BFb0103041.
- [Anc12] A. ANCONA, On positive harmonic functions in cones and cylinders, *Rev. Mat. Iberoam.* **28** (2012), 201–230.

- [BHK01] M. BONK, J. HEINONEN, and P. KOSKELA, *Uniformizing Gromov Hyperbolic Spaces*, *Astérisque*, 270, 2001.
- [BJ06] A. BOREL and L. JI, *Compactifications of symmetric and locally symmetric spaces*, *Mathematics: Theory & Applications*, Birkhäuser, Boston, 2006.
- [BH99] M. R. BRIDSON and A. HAEFLIGER, *Metric Spaces of Non-Positive Curvature*, *Grundlehren der Mathematischen Wissenschaften* **319**, Springer, Berlin, 1999. doi:10.1007/978-3-662-12494-9.
- [Cho69] G. CHOQUET, *Lectures on analysis. Vol. II: Representation theory*, W. A. Benjamin, Inc., New York-Amsterdam, 1969.
- [Dur83] P. L. DUREN, *Univalent Functions*, *Grundlehren der Math. Wissenschaften* **259**, Springer-Verlag, New York, 1983.
- [GT98] D. GILBARG and N. S. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*, *Classics in Mathematics*, Springer, Berlin, 1998.
- [GW93] S. GIULINI and W. WOESS, The Martin compactification of the Cartesian product of two hyperbolic spaces, *J. Reine Angew. Math.* **444** (1993), 17–28. Available at <http://eudml.org/doc/153567>.
- [Hel14] L. HELMS, *Potential Theory*, Springer, London, 2014. doi:10.1007/978-1-4471-6422-7.
- [Her11] G. HERGLOTZ, Ber. über Potenzreihen mit positivem, reellen Teil im Einheitskreis, *Verh. Sächs. Akad. Wiss. Leipzig* (1911), 501–511.
- [Her62] R.-M. HERVÉ, Recherches axiomatiques sur la théorie des fonctions wurharmoniques et du potentiel, *Ann. Inst. Fourier (Grenoble)* **12** (1962), 415–571.
- [IP94] D. IOFFE and R. PINSKY, Positive harmonic functions vanishing on the boundary for the laplacian in unbounded horn-shaped domains, *Transactions AMS* **342** no. 2 (1994), 773–791.
- [KK66] J. KADLEC and A. KUFNER, Characterization of functions with zero traces by integrals with weight functions. I, *Časopis Pěst. Mat.* **91** (1966), 463–471. Available at <http://eudml.org/doc/20952>.
- [KB02] I. KAPOVICH and N. BENAKLI, Boundaries of hyperbolic groups, in *Combinatorial and Geometric Group Theory, Contemp. Math.* **296**, Amer. Math. Soc., Providence, RI, 2002, pp. 39–93. doi:10.1090/conm/296/05068.
- [Mir70] C. MIRANDA, *Partial Differential Equations of Elliptic Type*, second ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete* **2**, Springer, Berlin, 1970.
- [Pin95] R. G. PINSKY, *Positive Harmonic Functions and Diffusion*, *Cambridge Studies in Advanced Mathematics* **45**, Cambridge University Press, Cambridge, 1995. doi:10.1017/CBO9780511526244.
- [Sta65] G. STAMPACCHIA, Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus, *Ann. Inst. Fourier (Grenoble)* **15** no. 1 (1965), 189–258.
- [Ste70] E. M. STEIN, *Singular Integrals and Differentiability Properties of Functions*, *Princeton Mathematical Series, No. 30*, Princeton University Press, Princeton, N.J., 1970.

MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, GERMANY
 E-mail addresses: m.kemper@uni-muenster.de, j.lohkamp@uni-muenster.de